blue 🗑 of california

2 04 111	Gene Expression Profiling and Protein Biomarkers for Prostate				
2.04.111	Cancer Management				
Original Policy Date:	March 1, 2016	Effective Date:	January 1, 2023		
Section:	2.0 Medicine	Page:	Page 1 of 54		

Policy Statement

- I. The one-time use of Oncotype DX[®] Prostate, Decipher[®] (prostate biopsy) or Prolaris[®] gene expression profiling tests may be considered **medically necessary** to guide management of prostate cancer when **all** of the following have been met:
 - A. Needle biopsy discloses localized (i.e., less than T3) adenocarcinoma of prostate (with no clinical evidence of metastasis or lymph node involvement, i.e., NOMO)
 - B. Individual falls into **one** of the following Risk categories as outlined by the National Comprehensive Cancer Network (NCCN) guidelines (see Policy Guidelines)*:
 - C. Very Low risk
 - D. Low risk
 - E. Favorable Intermediate risk
 - F. No documentation of significant co-morbidities, advanced age, or other factors that suggest the individual might have a life expectancy of less than 10 years
 - G. No documentation the individual is not a candidate for either active surveillance or definitive therapy (e.g., external beam radiotherapy, brachytherapy, or radical prostatectomy)
 - H. No documentation the individual has received prior treatment for prostate cancer
- II. The one-time use of the Decipher or Prolaris gene expression profiling tests may be considered **medically necessary** in patients with unfavorable intermediate and high risk disease and no documentation of significant co-morbidities, advanced age, or other factors that suggest the individual might have a life expectancy of less than 10 years
- III. The one-time use of Decipher Prostate RP (genomic classifier) may be considered medically necessary to inform adjuvant treatment if adverse features are found after radical prostatectomy or during workup for radical prostatectomy with PSA persistence or recurrence (category 2B for recurrence).
- IV. The use of gene expression analysis other than listed above, and protein biomarkers to guide management of prostate cancer is considered **investigational** in all situations.

NOTE: Refer to Appendix A to see the policy statement changes (if any) from the previous version.

Policy Guidelines

Testing to guide referrals for prostate biopsy (before cancer has been diagnosed) are discussed in the Blue Shield of California Medical Policy: Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer.

Table PG1: National Comprehensive Cancer Network (NCCN) Risk Categories for Prostate Cancer (Version 4.2018)¹³

Risk Category	Tumor Characteristics
Very Low Risk	Clinical Stage: T1c AND
	 Gleason score less than or equal to 6/grade group 1 AND
	PSA less than 10 ng/ml AND
	• Fewer than 3 prostate biopsy fragments/cores positive, less than or equal to
	50% cancer in each fragment/core AND

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 2 of 54

Risk Category	Tumor Characteristics
	• PSA density less than 0.15 ng/ml/g
Low Risk	Clinical Stage: T1 to T2a AND
	 Gleason score less than or equal to 6/grade group 1 AND
	• PSA less than 10 ng/ml
Favorable	Clinical Stage: T2b to T2c OR
Intermediate Risk	 Gleason score 3+4=7/grade group 2 OR
	PSA 10 ng/ml to 20 ng/ml
	AND
	 Percentage of positive biopsy cores less than 50%
Unfavorable	Clinical Stage: T2b to T2c OR
Intermediate Risk	 Gleason score 3+4=7/grade group 2 or Gleason score 4+3=7/grade group 3 OR
	PSA 10 ng/ml to 20 ng/ml
High Risk	Clinical Stage: T3a OR
	 Gleason score 8/grade group 4 or Gleason score 4+5=9/grade group 5 OR
	PSA greater than 20 ng/ml
Very High Risk	Clinical Stage: T3b to T4 OR
	Primary Gleason pattern 5 OR
	Greater than 4 cores with Gleason score 8-10/grade group 4 or 5

PSA: Prostate Specific Antigen

Note: TI and T2 have not spread outside the prostate. T3 and T4 have grown outside the prostate (seminal vesicles or beyond). T2a is half or less of only one side (left or right); T2b is more than one half but still only on one side; T2c is on both sides of the prostate (both left and right).

*As long as the patient at least meets the criteria for favorable intermediate risk, the testing can be approved.

Coding

The following CPT code is specific to the Prolaris® test:

• **81541**: Oncology (prostate), mRNA gene expression profiling by real-time RT-PCR of 46 genes (31 content and 15 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a disease-specific mortality risk score

The following CPT code is specific to the Oncotype DX[®] Prostate test:

• **0047U**: Oncology (prostate), mRNA, gene expression profiling by real-time RT-PCR of 17 genes (12 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a risk score

If the test uses an algorithm for analysis of the results of testing for multiple genes and the results are reported as a type of score, the following CPT code should be reported:

• **81599:** Unlisted multianalyte assay with algorithmic analysis

The following CPT code represents the Decipher Biosciences Decipher Prostate test:

• **81542**: Oncology (prostate), mRNA, microarray gene expression profiling of 22 content genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as metastasis risk score

If no algorithm is performed and the results are not reported as a score, the following CPT code may be reported:

• 81479: Unlisted molecular pathology procedure

Description

Gene expression profile analysis and protein biomarkers have been proposed as a means to riskstratify patients with prostate cancer to guide treatment decisions. These tests are intended to be used either on prostate needle biopsy tissue to guide management decisions for active surveillance or therapeutic intervention, to guide radiotherapy use after radical prostatectomy (RP), or to guide medication selection after progression in metastatic castration-resistant prostate cancer.

Related Policies

• Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer

Benefit Application

Benefit determinations should be based in all cases on the applicable contract language. To the extent there are any conflicts between these guidelines and the contract language, the contract language will control. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

Some state or federal mandates (e.g., Federal Employee Program [FEP]) prohibits plans from denying Food and Drug Administration (FDA)-approved technologies as investigational. In these instances, plans may have to consider the coverage eligibility of FDA-approved technologies on the basis of medical necessity alone.

Regulatory Status

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). Prolaris® (Myriad Genetics), Oncotype DX® Prostate and Oncotype DX AR-V7 Nuclear Detect (Genomic Health), Decipher gene expression profiling test (Decipher Corp), and the ProMark™ protein biomarker test (Metamark Genetics) are available under the auspices of the CLIA. Laboratories that offer laboratory-developed tests must be licensed by the CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration (FDA) has chosen not to require any regulatory review of these tests.

In November 2015, the FDA's Office of Public Health Strategy and Analysis published a report suggesting FDA oversight of laboratory-developed tests.^{15,} The FDA argued that many tests need more FDA oversight than the regulatory requirements of the CLIA. The CLIA standards relate to laboratory operations but do not address inaccuracies or unreliability of specific tests. Prolaris is among the 20 case studies in the document cited as needing FDA oversight. The report asserted that patients are potentially receiving inappropriate prostate cancer care because there is no evidence that results from the test meaningfully improve clinical outcomes.

Rationale

Background Prostate Cancer

Prostate cancer is the second most common noncutaneous cancer diagnosed among men in the U.S. Autopsy studies in the era before the availability of prostate-specific antigen (PSA) screening have identified incidental cancerous foci in 30% of men 50 years of age, with incidence reaching 75% at age 80 years.¹

Localized prostate cancers may appear very similar clinically at diagnosis.^{2,} However, they often exhibit diverse risk of progression that may not be captured by clinical risk categories (e.g., D'Amico criteria) or prognostic tools based on clinical findings, including PSA titers, Gleason grade, or tumor stage.^{3,4,5,6,7,} In studies of conservative management, the risk of localized disease progression based on prostate cancer-specific survival rates at 10 years may range from 15%^{8,9,} to 20%^{10,} to perhaps 27% at 20-year follow-up.^{11,} Among older men (ages ³70 years) with low-risk disease, comorbidities typically supervene as a cause of death; these men will die with prostate cancer present, rather than from cancer itself. Other very similar appearing low-risk tumors may progress unexpectedly rapidly, quickly disseminating and becoming incurable.

Risk Stratification in Newly Diagnosed Disease

In the U. S., most prostate cancers are clinically localized at diagnosis due in part to the widespread use of PSA testing. Clinicopathologic characteristics are used to stratify patients by risk based on the extent of the primary tumor (T category), nearby lymph node involvement (N category), metastasis (M category), PSA level and Gleason score. The National Comprehensive Cancer Network and American Urological Association risk categories for clinically localized prostate cancer are similar, derived from the D'Amico criteria and broadly include low-, intermediate-, or high-risk as follows as well as subcategories within these groups:^{12,13,}

- Low: T1-T2a and Gleason score $\leq 6/G$ leason grade group 1 and PSA level ≤ 10 ng/mL;
- Intermediate: T2b-T2c or Gleason score 3+4=7/Gleason grade group 2 or Gleason score 4+3=7/Gleason grade group 3 or PSA level 10-20 ng/mL;
- High: T3a or Gleason score 8/Gleason grade group 4 or Gleason score 9-10/Gleason grade group 5 or PSA level >20 ng/mL.

Risk stratification is combined with patient age, life expectancy, and treatment preferences to make initial therapy decisions.

Monitoring After Prostatectomy

All normal prostate tissue and tumor tissue are theoretically removed during radical prostatectomy (RP), so the serum level of PSA should be undetectable following RP. Detectable PSA post-RP indicates residual prostate tissue and presumably persistent or recurrent disease. Prostate-specific antigen is serially measured following RP to detect early disease recurrence. The National Comprehensive Cancer Network recommends monitoring serum PSA every 6 to 12 months for the first 5 years and annually thereafter.^{12,} Many recurrences following RP can be successfully treated. The American Urological Association recommends that biochemical recurrence be defined as a serum PSA of 0.2 ng/mL or higher, which is confirmed by the second determination with a PSA level of 0.2 ng/mL or higher.^{14,}

Castration-Resistant Prostate Cancer

Androgen deprivation therapy (ADT) is generally the initial treatment for patients with advanced prostate cancer. Androgen deprivation therapy can produce tumor response and improve quality of life but most patients will eventually progress on ADT. Disease that progresses while the patient is on ADT is referred to as castration-resistant prostate cancer. After progression, continued ADT is generally used in conjunction with other treatments. Androgen pathways are important in the progression of castration-resistant prostate cancer. Several drugs have been developed that either inhibit enzymes involved in androgen production or inhibit the androgen receptor, such as abiraterone and enzalutamide. Taxane chemotherapy with docetaxel or cabazitaxel may also be used after progression. Immunotherapy (sipuleucel-T) or radium 223 are options for select men.

Literature Review

This review was informed by a TEC Assessment (2014) addressing disease detected on needle biopsy, and has been supplemented by a TEC Assessment (2015) addressing high-risk disease after

prostatectomy. The Blue Cross Blue Shield Association Medical Advisory Panel also reviewed the evidence in September 2017.

Evidence reviews assess whether a medical test is clinically useful. A useful test provides information to make a clinical management decision that improves the net health outcome. That is, the balance of benefits and harms is better when the test is used to manage the condition than when another test or no test is used to manage the condition.

The first step in assessing a medical test is to formulate the clinical context and purpose of the test. The test must be technically reliable, clinically valid, and clinically useful for that purpose. Evidence reviews assess the evidence on whether a test is clinically valid and clinically useful. Technical reliability is outside the scope of these reviews, and credible information on technical reliability is available from other sources.

Initial Management Decision: Active Surveillance versus Therapeutic Intervention

The divergent behavior of localized prostate cancers creates uncertainty whether to treat immediately or follow with active surveillance.^{16,17,} With active surveillance, the patient will forgo immediate therapy and continue regular monitoring until signs or symptoms of disease progression are evident, at which point curative treatment is instituted.^{18,} A patient may alternatively choose potentially curative treatment upfront.^{19,} Surgery (i.e., radical prostatectomy [RP]) or external-beam radiotherapy (EBRT) is most commonly used to treat patients with localized prostate cancer. Complications most commonly reported with RP or EBRT and with the greatest variability are incontinence (0%-73%) and other genitourinary toxicities (irritative and obstructive symptoms); hematuria (typically \leq 5%); gastrointestinal and bowel toxicity, including nausea and loose stools (25%-50%); proctopathy, including rectal pain and bleeding (10%-39%); and erectile dysfunction, including impotence (50%-90%).^{13,} In a population-based retrospective cohort study using administrative hospital data, physician billing codes, and cancer registry data, Nam et al (2014) estimated the 5-year cumulative incidence of admission to hospital for a treatment-related complication following RP or EBRT to be 22% (95% confidence interval [CI], 21.7% to 22.7%).^{20,} In the Prostate Testing for Cancer and Treatment (ProtecT) trial (2016), active surveillance, immediate RP, and immediate EBRT for the treatment of clinically localized prostate cancer were compared in 1643 men identified through prostate-specific antigen (PSA) testing.^{21,} About 90% of the participants had a PSA level less than 10 ng/mL; two-thirds were Gleason score 6 and 20% were Gleason score 7; all were clinical stage T1c or T2. The mean age was 62 years. At a median of 10-year follow-up, prostate cancer-specific survival was high and similar across the 3 treatment groups: 98.8% (95% CI, 97.4% to 99.5%) in active surveillance, 99.0% (95% CI, 97.2% to 99.6%) in the surgery group, and 99.6% (95% CI, 98.4% to 99.9%) in the radiotherapy (RT) group. Surgery and RT were associated with lower incidences of disease progression and metastases compared with active surveillance. Approximately 55% of men in the active surveillance group had received a radical treatment by the end of follow-up. Similarly, very high prostate cancer-specific survival and metastasis-free survival outcomes were reported by large, prospective cohorts of active surveillance patients in the U.S. and Canada.^{22,23,}

The Prostate Cancer Intervention versus Observation Trial (PIVOT) randomized 731 men in the U. S. with localized prostate newly diagnosed cancer to RP or observation. The patients were 40% low-risk, 34% intermediate-risk and 21% high- risk. Results from PIVOT also concluded that RP did not prolong survival compared with observation through 12 years and 19.5 years of follow-up in the primary analyses including all risk groups.^{24,25,} However, among men with intermediate-risk tumors, surgery was associated with a 31% relative reduction in all-cause mortality compared with observation (hazard ratio [HR], 0.69; 95% CI, 0.49 to 0.98; absolute risk reduction, 12.6%).

An observational study by van den Bergh et al (2012), comparing sexual function of men with low-risk prostate cancer who chose active surveillance with men who received RT or RP, found that those who chose active surveillance were more often sexually active than similar men who received RP.²⁶. In a

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 6 of 54

2011 report of quality of life (QOL) for men in the Scandinavian Prostate Cancer Group Study Number 4, after a median follow-up of more than 12 years, distress caused by treatment-related side effects was reported significantly more often by men assigned to RP than by men assigned to watchful waiting.^{27,}

The American Urological Association (AUA), in joint guidelines (2017), has suggested that physicians recommend active surveillance for most men with low-risk localized prostate cancer but offer RP or RT to select low-risk, localized patients who have a high probability of progression on active surveillance.^{13,} The guidelines also suggested that physicians recommend RP or RT plus androgen deprivation therapy (ADT) to patients with intermediate-risk prostate cancer and that RT alone or active surveillance may also be offered to select patients with favorable intermediate-risk localized cancer.

Clinical Context and Test Purpose

In men with newly diagnosed low- or favorable intermediate risk clinically localized prostate cancer, the purpose of gene expression profiling (GEP) and protein biomarker testing is to inform a decision whether to undergo immediate therapy or to forgo immediate therapy and begin active surveillance. In individuals with newly diagnosed unfavorable intermediate- or high-risk clinically localized prostate cancer, the purpose of GEP and protein biomarker testing is to inform a decision between local therapy alone (radical prostatectomy or radiotherapy) and treatment intensification (radiotherapy plus androgen deprivation therapy).

The first question addressed in this evidence review is: Does GEP improve outcomes in newly diagnosed men with clinically localized prostate cancer, compared with clinicopathologic risk stratification or when used with clinicopathologic risk stratification? The specific questions differ by patient risk:

For newly diagnosed patients at low-risk, does GEP identify a group of patients who should receive immediate RP or RT instead of active surveillance?

For newly diagnosed patients at favorable intermediate-risk, does GEP identify a group of patients who can safely forgo immediate RP or RT and be followed with active surveillance?

For newly diagnosed patients at unfavorable intermediate- or high-risk, does GEP identify a group of patients who can safely forgo ADT?

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with newly diagnosed localized prostate cancer, who have not undergone treatment for prostate cancer, and who are deciding between therapeutic intervention and active surveillance, or between single and multimodal therapy.

Interventions

Gene expression profiling refers to the analysis of messenger RNA expression levels of many genes simultaneously in a tumor specimen and protein biomarkers.^{28,29,30,31,32,33}, Three GEP tests and 1 protein biomarker test are intended to stratify biologically prostate cancers diagnosed on prostate needle biopsy: Prolaris, Oncotype DX Prostate Cancer Assay, and Decipher Biopsy are GEP tests that use archived tumor specimens as the messenger RNA source, reverse-transcriptase polymerase chain reaction amplification, and the TaqMan low-density array platform. A protein biomarker test, ProMark is an automated quantitative imaging method to measure protein biomarkers by immunofluorescent staining in defined areas in intact formalin-fixed paraffin-embedded biopsy tissue to provide independent prognostic information to aid in the stratification of patients with prostate cancer to active surveillance or therapy.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 7 of 54

Comparators

Clinicopathologic risk stratification along with age/life expectancy and patient preference are currently being used to make decisions about prostate cancer management. Clinical characteristics (e.g., stage, biopsy Gleason grade, serum PSA level) and demographic characteristics (e.g., age, life expectancy) are combined to classify men according to risk. National Comprehensive Cancer Network (NCCN) and AUA have provided treatment recommendations based on risk stratification and life expectancy.^{12,34,} The Kattan et al (2003) nomogram was developed to predict the risk of indolent cancer in a low-risk population considering active surveillance.^{35,} The Cancer of the Prostate Risk Assessment (CAPRA) is a pretreatment nomogram that provides risk prediction of outcomes following RP developed from a cohort of RP patients.^{36,}

Outcomes

Beneficial outcomes resulting from a true test result are prolonged survival, improved QOL, and reduction in unnecessary treatment-related adverse events. Harmful outcomes resulting from a false test result are recurrence, metastases or death, and unnecessary treatments. The outcomes of interest are listed in Table 1. The primary survival outcome of interest is disease-specific survival because overall survival (OS) is very high in this group.

Outcomes	Details
Overall survival	10-year survival
Disease-specific survival	10-year prostate cancer-free survival; 10-year prostate cancer death rate; 10- year recurrence rate
Quality of life	See Chen et al (2014) ^{37,} for NCI-recommended health-related quality of life measures for localized prostate cancer
Treatment-related morbidity	Adverse events of radiotherapy, radical prostatectomy, or androgen- deprivation therapy

NCI: National Cancer Institute.

Ten-year outcomes are of interest due to the prolonged natural history of localized prostate cancer.

Study Selection Criteria

For the evaluation of clinical validity of the Prolaris, Oncotype DX Prostate, ProMark protein biomarker, and Decipher Biopsy tests, studies that meet the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a validation cohort independent of the development cohort;
- Included a suitable reference standard (10 year prostate cancer-specific survival or death rate)
- Patient/sample clinical characteristics were described
- Patient/sample selection criteria were described.

Prolaris

Prolaris is used to quantify expression levels of 31 cell cycle progression (CCP) genes and 15 housekeeper genes to generate a CCP score. This section reviews Prolaris for initial management decisions in newly diagnosed, localized cancer.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Three studies reporting clinical validity related to newly diagnosed men with clinically localized prostate cancer are summarized in Table 2.

Study	Design	Dates	Sites	Ν	Population
Cuzick et al (2012) ^{38,}	Retrospective cohort from prospective registry	1990-1996	6 U.K. registries; not screen-detected	349	Clinically localized; 66% Gleason score 6-7; 46% PSA level ≤25 ng/mL
Cuzick et al (2015) ^{39,}	Retrospective cohort from prospective registry	1990-2003	3 U.K. registriesª; not screen-detected	761	Clinically localized; 74% Gleason score ≤7, mean PSA level 21 ng/mL
Lin et al (2018) ^{40,}	Validation cohort: Subset of Cuzick et al (2015) Clinical testing cohort: Consecutive men with biopsies submitted for testing to manufacturer	1990-2003 2013-2016	3 U.K. registries ^a ; not screen-detected NA; manufacturer database	585 19,215	 See Cuzick et al (2015) Median PSA level, 5.6 ng/mL (IQR, 44-7.6 ng/mL) NCCN risk: Low, 57% Favorable intermediate, 20% Intermediate, 17% High 7%
					 nigh, /%

Table 2.	Clinical	Validitv	Studies	Assessina	Prolaris	for Informing	a Initial	Manaaemen	t Decisions

QR: interquartile range; NA: not available; NCCN: National Comprehensive Cancer Network; PSA: prostatespecific antigen.

^a No overlap in population with Cuzick et al (2012).Cuzick et al (2012) examined the Prolaris prognostic value for prostate cancer death in a conservatively managed needle biopsy cohort.^{38,} Cell cycle expression data were read blind to all other data. Patients were identified from 6 cancer registries in Great Britain and were included if they had clinically localized prostate cancer diagnosed by needle biopsy between 1990 and 1996; were younger than 76 years at diagnosis; had a baseline PSA measurement; and were conservatively managed. Potentially eligible patients who underwent RP, died, showed evidence of metastatic disease within 6 months of diagnosis, or received hormone therapy before diagnostic biopsy were excluded. The original biopsy specimens were retrieved and centrally reviewed by a panel of expert urologic pathologists to confirm the diagnosis and, where necessary, to reassign Gleason scores.^{41,} Of 776 patients diagnosed by needle biopsy and for which a sample was available to review histology, needle biopsies were retrieved for 527 (68%), 442 (84%) of which had adequate material to assay. From the 442 samples, 349 (79%) produced a CCP score and had a complete baseline and follow-up information, representing 45% of 776 patients initially identified. The median follow-up time was 11.8 years. Ninety deaths from prostate cancer occurred within 2799 person-years. The primary, unadjusted analysis found a 1-unit increase in CCP score associated with a 2-fold increase (HR=2.02) in the risk of dying from prostate cancer (see Table 3). In a multivariate model including CCP, Gleason score, and PSA level, the adjusted HR for a 1-unit increase in CCP score was 1.65. However, changes in HRs may not reflect meaningful changes in absolute risk. As is shown in Table 4, Kaplan-Meier analyses of the 10 year risk of prostate cancer death are stratified by CCP score groupings. It appears that there might be a large change in risk for scores below 2 compared with above 2, but no CIs are reported so it is impossible to draw conclusions. Measures that would suggest improved discriminatory ability (e.g., area under the curve [AUC] or reclassification) compared with an existing nomogram were not reported in Cuzick et al (2012). The authors did not provide evidence that the test could correctly reclassify men initially at high-risk to lower risk to avoid overtreatment, or conversely, correctly reclassify those initially at low-risk to high-risk to avoid undertreatment.

Cuzick et al (2012) examined the Prolaris prognostic value for prostate cancer death in a conservatively managed needle biopsy cohort.^{38,} Cell cycle expression data were read blind to all other data. Patients were identified from 6 cancer registries in Great Britain and were included if they had clinically localized prostate cancer diagnosed by needle biopsy between 1990 and 1996; were younger than 76 years at diagnosis; had a baseline PSA measurement; and were conservatively managed. Potentially eligible patients who underwent RP, died, showed evidence of metastatic disease within 6 months of diagnosis, or received hormone therapy before diagnostic biopsy were

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 9 of 54

excluded. The original biopsy specimens were retrieved and centrally reviewed by a panel of expert urologic pathologists to confirm the diagnosis and, where necessary, to reassign Gleason scores.^{41,} Of 776 patients diagnosed by needle biopsy and for which a sample was available to review histology, needle biopsies were retrieved for 527 (68%), 442 (84%) of which had adequate material to assay. From the 442 samples, 349 (79%) produced a CCP score and had a complete baseline and follow-up information, representing 45% of 776 patients initially identified. The median follow-up time was 11.8 years. Ninety deaths from prostate cancer occurred within 2799 person-years.

The primary, unadjusted analysis found a 1-unit increase in CCP score associated with a 2-fold increase (HR=2.02) in the risk of dying from prostate cancer (see Table 3). In a multivariate model including CCP, Gleason score, and PSA level, the adjusted HR for a 1-unit increase in CCP score was 1.65. However, changes in HRs may not reflect meaningful changes in absolute risk. As is shown in Table 4, Kaplan-Meier analyses of the 10 year risk of prostate cancer death are stratified by CCP score groupings. It appears that there might be a large change in risk for scores below 2 compared with above 2, but no Cls are reported so it is impossible to draw conclusions. Measures that would suggest improved discriminatory ability (e.g., area under the curve [AUC] or reclassification) compared with an existing nomogram were not reported in Cuzick et al (2012). The authors did not provide evidence that the test could correctly reclassify men initially at high-risk to lower risk to avoid overtreatment, or conversely, correctly reclassify those initially at low-risk to high-risk to avoid undertreatment.

Cuzick et al (2015) examined 3 U.K. cancer registries from 1990 to 2003 to identify men with prostate cancer who were conservatively managed following needle biopsy, with follow-up through December 2012.^{39,} The authors stated that the samples did not overlap with Cuzick et al (2012). Men were excluded if they had undergone RP or RT within 6 months of diagnosis. A combination of the CCP and CAPRA scores (called the combined clinical cell cycle risk [CCR] score) was used to predict prostate cancer death. There were 989 men who fit eligibility criteria; CCP scores were calculable for 761 (77%), and combined CCP and clinical variables were available for 585 (59%). Median age at diagnosis was 70.8 years, and the median follow-up was 9.5 years. The prostate cancer mortality rate was 17% (n=100), with 29% (n=168) dying from competing causes. Higher CCP scores were associated with increased 10-year risk of prostate cancer mortality (see Table 5): 7% (CCP score <0), 15% (CCP score 0-1), 36% (CCP score 1-2), and 59% (CCP score >2). For the CCR score, the HR for 10-year prostate cancer mortality increased to 2.17 (95% CI, 1.83 to 2.57). The C statistic for the CAPRA score was 0.74; adding the CCP score increased the C statistic to 0.78 (no CIs for the C statistic were reported). Estimates with CIs for 10 year death rates for the CCR score are provided in a figure and given in Table 5 based on digitizing the figure. Note that the predictions appear to cross 100% for CCR of about 6. Treatment changes after 6 months were documented in only part of 1 of the 3 cohorts; at 24 months, 45% of the men in this cohort had undergone RT or prostatectomy.

Lin et al (2018)^{40,} validated a CCR cutoff of 0.8 using a subset of 585 conservatively managed men from the Cuzick (2015) cohort. Of the 585 men, 60 had CCR scores of 0.8 or less. Among the 284 men who were at low- or intermediate-risk by NCCN criteria, 59 had CCR scores of 0.8 or less. The text reports that the estimated 10-year prostate cancer mortality risk was 2.7% for men with CCR scores below the threshold and 3.3% (95% CI, 1.9% to 5.7%) at the threshold in the full cohort, and 2.3% below the threshold and 2.9% (95% CI, 1.3% to 6.7%) at the threshold in the cohort that excluded high-risk men. However, the Kaplan-Meier curves show an estimated prostate cancer mortality at 10 years of 0% for men with CCR of 0.8 or less in both cohorts. The Kaplan-Meier curve estimated prostate cancer mortality at 10 years for men with CCR greater than 0.8 was 20% in the full cohort and 9% in the cohort excluding high-risk men (see Table 5; precision estimates not provided).

Tward et al (2021) reported the association of the CCR score with 10-year risk of metastasis and progression in men with unfavorable intermediate- or high-risk prostate cancer. However, this study did not meet inclusion criteria for this review because it did not provide survival outcomes.^{42,}

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 10 of 54

Table 3. Univariate and Multivariate Associations Between CCP and Death From Prostate Cancer

Study	Ν	Unadjusted	Multivariate
		HRº (95% CI)	HRº (95% CI)
Cuzick et al (2012) ^{38,}	349	2.02 (1.62 to 2.53)	1.65 (1.31 to 2.09)°
Cuzick et al (2015) ^{39,}	585	2.08 (1.76 to 2.46)	1.76 (1.47 to 2.14) ^b

CCP: Cell Cycle Progression; CI: confidence interval; HR: hazard ratio.

^a Adjusted for Gleason score and prostate-specific cancer level.

^b Adjusted for Cancer of the Prostate Risk Assessment.

^c For a 1-unit increase in CCP.

Table 4. Kaplan-Meier Estimates of Prostate Cancer Death at 10 Years by CCP Score Groupings in the Cuzick Validation Studies^c

	Cuzick et	t al (2012) ^{38,}	Cuzick et al (2015) ^{39,}		
CCP Score	Ν	10-Year Death Rate, %ª	Ν	10-Year Death Rate, %ª	
≤ 0	36	19.3	194	7	
0 to <i>≤</i> 1	133	19.8	251	15	
1 to ≤ 2	114	21.1	110	36	
2 to <i>≤</i> 3	50	48.2	30 ^b	59	
>3	16	74.9			

CCP: Cell Cycle Progression.

^a Confidence intervals were not reported.

^b Grouped CCP score >2.

 $^{\circ}$ No overlap in populations with Cuzick et al (2012) and Cuzick et al (2015).

Table 5. Predicted Risk of Prostate Cancer Death at 10 Years by CCR Score Groupings

Cuzick et al (2015) ^{39,}		Lin et al (2018) ^{40,} Using Data From Cuzick et al (2015) ^{39,}			
Clinical Cell Cycle Risk Score	Ν	10-Year Death Rate (95% Cl), %ª	CCR Score	N	10-Year Death Rate (95% Cl), % ^d	
-1	NR	1.0 (0.2 to 1.8)				
0		2.2 (0.7 to 3.4)	≤0.8	Full ^b : 60 Modified ^c : 59	Full: 0 (Cl NR) Modified: 0 (Cl NR)	
1		4.5 (2.3 to 7.0)	>0.8	Full ^b : 525 Modified ^c : 225	Full: 19.9. (CI NR) Modified: 8.7 (CI NR)	
2		9.9 (6.4 to 13.0)				
3		20.2 (16.2 to 24.1)				
4		43.1 (34.1 to 51.2)				
5		73.5 (59.4 to 92.8)				
6		109.7 (82.0 to 120.8)				

CCR: combined clinical cell cycle risk; CI: confidence interval; NR: not reported.

^a Estimated from digitizing a figure.

^b Including all men from the validation cohort (»52% high-risk).

^c Excluding high-risk men in the validation cohort.

^d Based on the Kaplan-Meier plots.

Lin et al (2018) also reported reclassification of men using the CCR score threshold (0.8) in a group of 19,215 consecutive patients whose biopsies were sent for Prolaris testing between 2013 and 2016 (see Table 6).^{40,} According to the table of clinicopathologic features of patients, 14,685 of the 19,215 men had a low or favorable intermediate-risk by NCCN risk classification. However, in the reclassification table and the text describing the table (see Table 6), the authors said that only 8177 of the 19,215 men met NCCN criteria for active surveillance based on low/favorable intermediate-risk clinicopathologic features. It is not clear why fewer men were categorized as meeting NCCN low/favorable intermediate criteria for the purposes of demonstrating reclassification and, therefore, it is not clear how many of the 14685 men at low- or intermediate-risk by NCCN criteria would have been reclassified using the CCR threshold.

Table 6. Reclassification of NCCN Risk Stratification Criteria for Active Surveillance With the CCR Score^a

NCCN Risk Group	CCR Score ≤0.8	CCR Score >0.8	Total
Met NCCN criteria for active surveillance ^b	7463	714	8177 ^b
Did not meet NCCN criteria for active	5758	52809	11038 ^b
surveillance ^b			
Total	13221	5994	19215

CCR: combined clinical cell cycle risk; NCCN: National Comprehensive Cancer Network.

^a Adapted from Lin et al (2018).^{40,}

^b Sample sizes here do not match the number of men reported to be low and favorable intermediate vs. intermediate and high-risk.

The purpose of the limitations tables (see Tables 7 and 8) is to display notable limitations identified in each study.

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Cuzick et al (2012) ^{38,}	4. Not screen selected; higher risk than intended use	1. Thresholds not described		4. Reclassification not provided	
Cuzick et al (2015) ^{39,}	4. Not screen selected; higher risk than intended use	1. Thresholds not described			
Lin et al (2018) ^{40,}	Note. Validation cohort is from Cuzick (2015)				

Table 7. Study Relevance Limitations

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

Table 8. Study Design and Conduct Limitations

Study	Selection ^a	Blinding ^b	Delivery of Test ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
Cuzick et al (2012) ^{38,}	1. Unclear if all men meeting criteria were included				2,3. 349 of 776 had sufficient data for inclusion	1. Cls not reported for KM estimates at 10 y for CCP
Cuzick et al (2015) ^{39,}	1. Unclear if all men meeting criteria were included				2,3. 585 of 989 had sufficient data for inclusion	1. CIs not reported for KM estimates at 10 y for CCP
Lin et al (2018) ^{40,}	Note. Used data from Cuzick (2015) for validation cohort					1. Cls not reported for KM estimates at 10 y for CCR

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

CCP: Cell Cycle Progression; CCR: combined clinical cell cycle risk; CI: confidence interval; KM: Kaplan-Meier. ^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

^bBlinding key: 1. Not blinded to results of reference or other comparator tests.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 12 of 54

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication. ^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

In summary, Table 3 displays the association between CCP score adjusted for CAPRA; Table 4 shows the risk of death by groups of CCP score; and Table 5 shows predicted risk of death by CCR score, which is the combined CCP and CAPRA score. The CCR score is most relevant because it appears in the sample report provided by the manufacturer. Table 3 demonstrates an association between CCP and the risk of death on the relative scale but does not necessarily indicate that there is a difference in absolute risk that would be meaningful for clinical decision making. Table 4 displays the estimated absolute risk of death for the CCP score but notably absent are CIs that would help in interpretation. However, given the data provided, several concerns arise. Even the lowest risk group shown in Cuzick et al (2012) has a 10-year death rate of 20%, which may be explained by the population characteristics (i.e., not PSA screen-selected, a third with Gleason >7 score and half with PSA level >25 ng/mL); however, a death rate of 20% is unlikely to be low enough to forgo immediate treatment.^{38,} Table 4 does not include the death rates by CCR score; however, the predicted 10-year prostate cancer death rates by CCR score were provided in a figure in Cuzick et al (2015). The predicted 10-year risk for CAPRA alone compared with CCR was provided in a dot plot in Cuzick et al (2015). The authors stated that CCR identified 11 men with a CAPRA score of 2 (indicating an estimated 10-year mortality rate of 4%) who "had a higher risk" based on CCR score. From the dot plot, it appears that for these 11 men, the 10-year mortality rate estimated by CCR score ranged from just greater than 4% to about 8%. The authors also indicated that for 31 men with CAPRA score of 3 (corresponding to the 10-year risk of death rate of 5.7%), the risk as estimated by CCR was less than 4.0% from the plot the CCR estimated risk appears to range from about 2.5% to 4% for those 31 men. It is not clear that either of these reclassifications would change the estimated mortality enough to alter treatment decisions. Using data from Cuzick et al (2015) and a CCR cutoff of 0.8, Lin et al (2018) estimated that the 10-year death rate for men with low to favorable intermediate-risk was 0% in men with CCR score of 0.8 or less and 9% for men with CCR score greater than 0.8, but precision estimates were not provided.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials (RCTs).

No studies that directly supported the clinical utility of Prolaris.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Three decision-impact studies have assessed the potential impact of Prolaris on physicians' treatment decisions in patients.^{43,44,45,} The authors of these studies³/₄Crawford et al (2014),^{43,} Shore et al (2014),^{44,} and Shore et al (2016)^{45,} have suggested that their findings supported the "clinical utility" of the test, based on whether the results would lead to a change in treatment. Pathology results were

not reported for these studies. Given the lack of established clinical validity and no reported outcomes, it is uncertain whether any treatment changes were clinically appropriate.

In trying to construct a chain of evidence from clinical validity to clinical utility, there are several obstacles to drawing conclusions. First, as noted in the clinical validity section, it is not clear if the test provides incremental value over the CAPRA score for decision making. In the example of reclassification given by Cuzick et al (2015), 11 men with a CAPRA estimated 10-year mortality risk rate of 4% were reclassified as having higher 10-year mortality estimated by CCR score with risk ranging from just greater than 4% to about 8%, and 31 men with CAPRA 10-year mortality risk rate of 5.7% were reclassified as having lower estimated risk by CCR of about 2.5% to 4%.⁴² It is not clear that these reclassifications would change treatment decisions.

Given that the PIVOT trial supported RP for the intermediate-risk group, showing a 30% relative and 12% absolute benefit for OS, in order to be suitable for clinical decision making, the test would have to identify a lower risk group of intermediate-risk men with very high negative predictive value (NPV) for survival with tight CIs. Because it is not clear how the Cuzick et al (2012) or Cuzick et al (2015) results would apply specifically to intermediate-risk men, it is not clear whether the test could be used to identify intermediate-risk men who can delay RP or RT.

Health Quality Ontario (2017) reported on a health technology assessment including a systematic review of the literature assessing the clinical utility of the Prolaris CCP.^{46,} The literature search identified Crawford et al (2014)^{43,} and Shore et al (2016).^{45,} Reviewers concluded that the GRADE rating of the quality of evidence was very low and that there was no evidence on clinical outcomes of patients whose treatment was informed by CCP results.

Section Summary: Prolaris

In a cohort of men conservatively managed following needle biopsy, Cuzick et al (2012) suggested that the CCP score alone was more prognostic than either PSA level or Gleason score for tumor-specific mortality at 10-year follow-up based on HRs.^{38,} Comparison with CAPRA score was not provided in Cuzick et al (2012). Cuzick et al (2015) found that discrimination improved somewhat by adding the CCP score to the CAPRA score, as reflected in the C statistic. ^{39,} Ten-year mortality rates based on CCP were inconsistent within Prolaris risk categories across Cuzick et al (2012) and Cuzick et al (2015). Numerical summaries of mortality rates for the CCR were provided in a figure in Cuzick (2015). The men included in the U.K. registries were not screen-selected, and a large proportion of the men in the validation studies were not low- or intermediate-risk.

No direct evidence is available to support the clinical utility of Prolaris for improving the net outcomes of patients with localized prostate cancer. The chain of evidence is also incomplete. The ProtecT trial showed 99% 10-year disease-specific survival in all 3 treatment groups: active surveillance, RT, and RP including predominately low-risk but also some intermediate-risk men. American Urolocial Association has recommended active surveillance in low-risk men. The low mortality rate estimated with tight precision makes it unlikely that a test intended to identify a subgroup of low-risk men with a net benefit from immediate treatment instead of active surveillance would find such a group.

The PIVOT trial preplanned subgroup analysis showed a reduction in mortality for RP compared with observation for men with intermediate-risk; AUA has recommended RT or RP for such men. For intermediate-risk men, a test designed to identify men who can receive active surveillance instead of RP or RT would need to show very high NPV for disease-specific mortality at 10 years and improvement in prediction compared with existing tools used to select such men. To forgo evidence-based beneficial treatment, there would have to be a very high standard of evidence for the clinical validity of the test.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 14 of 54

Oncotype DX Prostate

The Oncotype DX Prostate assay includes 5 reference genes and 12 cancer genes that represent 4 molecular pathways of prostate cancer oncogenesis: androgen receptor, cellular organization, stromal response, and proliferation. The assay results are combined to produce a Genomic Prostate Score (GPS), which ranges from 0 to 100. Higher GPS scores indicate more risk.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Five studies reporting clinical validity are summarized in Table 9. One publication by Klein et al (2014) compiled results for 3 cohorts: 2 in test development including a contemporary (1997-2011) group of patients in a prostatectomy study (n=441; Cleveland Clinic database, 1987-2004) and a biopsy study (n=167; Cleveland Clinic database, 1998-2007); the third was an independent clinical validation study cohort (n=395; University of California, San Francisco [UCSF] Database, 1998-2011).^{47,} A second study, Cullen et al (2015), evaluated men with NCCN clinically very low- to intermediate-risk undergoing prostatectomy.^{48,} The third study, Whalen et al (2016), evaluated men in a clinical practice setting.^{49,} The study by van Den Eeden et al (2018) included men from a cancer registry^{50,} and the study by Salmasi et al (2018) included men from an institutional database.^{51,}

Study	Design	Dates	Sites	Ν	Population
Klein et al (2014) ^{47,}	Case-cohort from prospective registry ^a	1998-2011	UCSF	395	Clinically localized; clinical stage T1/T2; PSA level ≤20 ng/mL, Gleason score ≤7; 3% African American
Cullen et al (2015) ^{48,}	Retrospective cohort from prospective longitudinal study	1990-2011	U.S. military centers	382	Clinically localized; clinical stage T1/T2; PSA level \leq 20 ng/mL, Gleason score \leq 7; 20% African American
Whalen et al (2016) ^{49,}	Prospective observational cohort (median follow-up, 5.2 y)	2013-2014	Mount Sinai Hospital	50	Clinically localized; clinical stage T1/T2; PSA level ≤20 ng/mL, Gleason score ≤7
Van Den Eeden et al (2018) ^{50,}	Retrospective cohort from registry (median follow-up, 9.8 y)	1995-2010	Kaiser Permanente Northern California	259	Prostate cancer who underwent RP within 12 mo of diagnosis, NCCN risk: very low, 3%; low, 21%; intermediate, 67%; high, 9%; 11% African American
Salmasi et al (2018) ^{51,}	Retrospective cohort from institutional database	2010-2016	UCLA	134	NCCN very low, low- or intermediate-risk prostate cancer treated with RP; 11% African American

Table 9. Clinical Validity Studies Assessing Oncotype DX Prostate

NCCN: National Comprehensive Cancer Network; PSA: prostate-specific antigen; RP: radical prostatectomy; UCSF: University of California, San Francisco.

 $^{\alpha}$ Only the validation sample cohort is listed. 52,

Results from the clinical validation study and prostatectomy study by Klein et al (2014) provided information on the potential clinical validity of this test.^{47,} The cohorts included men with a mix of low-to low-intermediate clinical risk characteristics using NCCN or AUA criteria. The Klein (2014) clinical validation study (see Table 9) was prospectively designed, used masked review of prostatectomy pathology results, and as such met the Reporting Recommendations for Tumor Marker Prognostic

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 15 of 54

Studies guidelines for biomarker validation.^{53,} The prostatectomy study used a case-cohort design to select a 1:3 ratio of recurrent to nonrecurrent patients. Favorable pathology was defined as freedom from high-grade or non-organ-confined disease. In the prostatectomy study, the ability of the GPS to stratify patients further within AUA groupings was related to the clinical recurrence-free interval in regression-to-the-mean estimated survival curves. Results of the Klein et al (2014) validation study showed that the GPS could refine the stratification of patients within specific NCCN criteria groupings, as summarized in Table 10. Proportions were estimated from a plot of GPS versus the percent likelihood of favorable pathology.^{47,}

NCCN Risk Level	Estimated Mean Likelihood of Favorable Tumor Pathology					
	NCCN Criteria, %	GPS + NCCN Criteria, Range, %				
Very low	»84	63-91				
Low	»76	55-86				
Intermediate	»56	29-75				

Table ⁻	10.	Recl	assificatio	on of	fΡ	rostc	Ite	Cancer	Risk	C	ategori	es \	Nith	On	coty	/pe	DX	Pros	state	Э
													_	_						

Adapted from the Klein et al (2014) validation study.^{47,}

GPS: Genomic Prostate Score; NCCN: National Comprehensive Cancer Network.

The actual number of patients correctly or incorrectly reclassified across all 3 categories cannot be ascertained from the data provided. The results would suggest that the combination of GPS plus clinical criteria can reclassify patients on an individual basis within established clinical risk categories. Extrapolation of this evidence to a true active surveillance population, for which the majority in the study would be otherwise eligible, is difficult because all patients had elective RP within 6 months of diagnostic biopsy.

The Klein et al (2014) prostatectomy study, although used to identify genes to include in the GPS, provided estimates of clinical recurrence rates stratified by AUA criteria^{54,} compared with rates after further stratification according to the GPS from the validation study. The survival curves for clinical recurrence reached nearly 18 years based on the dates individuals in the cohort were entered into the database (1987-2004). The reclassifications are summarized in Table 11. The GPS groups are grouped by tertiles defined in the overall study. Absolute rates and precision estimates of clinical recurrence by GPS low-, intermediate-, and high-risk groups were not reported. These data would suggest the GPS can reclassify patient risk of recurrence based on a specimen obtained at biopsy. However, the findings do not necessarily reflect a clinical scenario of predicting disease progression in untreated patients under active surveillance.

Table 11. Reclassification of Prostate Cancer 10 Year Clinical Recurrence Risk With Oncotype D	Х
Prostate	

Overall 10-Year Risk (AUA Risk Level)	10-Year Risk (GPS Low- Risk Group), %	10-Year Risk (GPS Intermediate-Risk Group), %	10-Year Risk (GPS High- Risk Group), %
3.4% (low)	2.0	3.4	7.0
9.6% (intermediate)	2.8	5.1	14.3
18.2% (high)	6.2	9.2	28.6

Adapted from the Klein et al (2014) prostatectomy study.^{47,}

AUA: American Urological Association; GPS: Genomic Prostate Score.

A retrospective cohort study by Cullen et al (2015) included men with NCCN-defined very low through intermediate-risk prostate cancer undergoing RP within 6 months of diagnosis.^{48,} The sample was obtained from men enrolled in the Center for Prostate Disease Research longitudinal study at 2 U.S. military medical centers. A Gleason score of 4 or 5 with the non-organ-confined disease was considered adverse pathology. Biopsies were available for 500 (57.9%) of 864 eligible patients; 382 (44.2% of eligible) with both adequate tissues for gene expression analysis and available RP pathology were included in the analysis. Selected patients were older (61.0 years vs. 59.7 years, p=.013) and had both higher Gleason scores (p<.001) and NCCN risk classification (29.8% vs 32.9% intermediate, p=.035). Median follow-up was 5.2 years and biochemical recurrence (BCR) occurred in

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 16 of 54

62 (15.4%). Estimates of 5-year BCR by GPS score are shown in Table 12. Adverse pathology was noted in 163 (34%) men. In an analysis adjusted for baseline characteristics, the GPS was associated with BCR-free survival and adverse pathology following RP (see Table 13). The GPS improved the C statistic for adverse pathology over NCCN risk alone from 0.63 to 0.72 (CIs not reported). Comparisons with other predictors such as CAPRA or Gleason score alone were not reported. Study implications were limited by the low proportion of eligible men in the analysis and differences between excluded and included men.

Whalen et al (2016) prospectively evaluated the correlation between GPS and final pathology at RP in a clinical practice setting.^{49,} Eligible men were 50 years of age and older with more than 10 years of life expectancy, PSA levels of 20 ng/mL or less, stage cTIc-cT2c newly diagnosed, untreated prostate cancer, who met NCCN classifications as very low-risk, low-risk, or low-intermediate risk. Men were enrolled from May 2013 to August 2014 at an academic medical center. Genomic Health reclassified patients' cancers as "less favorable," "consistent with," or "more favorable" than what would have been predicted by their NCCN risk group. Adverse pathology at RP was defined as any pT3 stage and primary Gleason grade of 4 or any-pattern 5. Fifty patients had RP pathology, and the reclassification results for these participants are discussed here; 21 (42%) met the definition of adverse pathology. The NCCN risk classification categorized 2 (4%) patients as very low-risk, 34 (68%) as low-risk, and 14 (28%) as a low-intermediate risk. Twenty-three (46%) patients were reclassified using GPS and the percentage with adverse pathology for the reclassification is shown in Table 14, as derived from data provided in the text. Confidence intervals were not provided.

Van Den Eeden et al (2018) reported on a retrospective study using a stratified cohort sampling design including 279 of 6184 men who were diagnosed with prostate cancer within a registry between 1995 and 2010 and underwent RP within 12 months of diagnosis, with a median follow-up of 9.8 years.^{50,} Characteristics are shown in Table 9. In an analysis adjusted for NCCN risk classifications, the GPS was associated with BCR-free survival, distant metastasis, and prostate cancer death following RP (see Table 13). Ten-year prostate cancer death by GPS score was displayed in a figure stratified by NCCN risk classification, which provides some information on potential reclassification. Ten-year prostate cancer death appears to be close to zero for men who are NCCN low-risk regardless of GPS score, indicating little useful reclassification of NCCN low-risk men based on GPS. For NCCN intermediate-risk, the risk of prostate cancer death ranges from approximately 0 for a GPS of less than 40 to close to 40% for a GPS of 100. It is unclear how many men with GPS less than 40 were NCCN favorable intermediate-risk.

Salmasi et al (2018) reported on a retrospective cohort from a UCLA institutional database of men with NCCN very low-, low-, or intermediate-risk prostate cancer treated with RP between 2010 and 2016 who had undergone simultaneous 3 Tesla multiparametric magnetic resonance imaging fusion targeted and systematic biopsies within the 6-month period prior to RP (see Table 9). The association between GPS and adverse pathology is shown in Table 13. The authors also reported an AUC for a model including Gleason score, GPS, and highest Prostate Imaging Reporting and Data System score determined by magnetic resonance imaging was 0.79 (95% CI, 0.71 to 0.87). The AUC of other models had overlapping CIs; the AUC of a model with Gleason score and highest Prostate Imaging Reporting and Data System score was 0.69 (95% CI, 0.59 to 0.78); and another model including Gleason score and PSA level was 0.68 (95% CI, 0.58 to 0.78).

Genomic Prostate Score	Ν	5-Year Biochemical Recurrence (95% Confidence Interval), %ª
10	Not reported	5.1 (2.7 to 9.1)
20		8.5 (5.8 to 13.4)
30		14.2 (10.2 to 19.0)
40		22.9 (18.0 to 28.8)
50		35.2 (27.1 to 45.4)
60		53.8 (38.6 to 65.6)
70		71.8 (50.6 to 89.3)

Table 12. Estimates of 5 Year Biochemical Recurrence With Oncotype DX Prostate

Genomic Prostate Score N	5-Year Biochemical Recurrence (95% Confidence Interval), %ª
80	87.3 (64.2 to 98.0)

Adapted from Cullen et al (2015).^{48,} ^a Estimated from digitizing a figure.

Study	Outcome	Ν	Unadjusted	Multivariate
			Ratio (95% CI)	Ratio (95% CI)
Klein et al (2014) ^{47,} validation study	Adverse pathology	395	OR=2.1 (1.4 to 3.2)	1.9 (1.3 to 2.8)¤
Cullen et al (2015) ^{48,}	BCR	392	HR=2.9 (2.0 to 4.2)	2.7 (1.8 to 3.8) ^b
	Adverse pathology	392	HR=3.2 (2.1 to 5.0)	HR=2.7 (1.8 to 4.4) ^c
Whalen et al (2016) ^{49,}	Adverse pathology	50	NR	OR=1.4 (NR) ^d
Van Den Eeden et al (2018) ^{50,}	Distant metastasis	259	HR=2.8 (1.6 to 4.6)	HR= 2.3 (1.4 to 3.9)°
	Prostate-cancer death	259	HR=3.2 (1.8 to 5.7)	HR=2.7 (1.5 to 4.8)°
	BCR	259	HR=2.5 (1.6 to 3.9)	HR=2.1 (1.4 to 3.1)¤
Salmasi et al (2018) ^{51,}	Adverse pathology	134	OR=3.8 (2.1 to 7.4)	OR=2.9 (1.5 to 5.9) ^e

BCR: biochemical recurrence; CI: confidence interval; GPS: Genomic Prostate Score; HR: hazard ratio; NCCN: National Comprehensive Cancer Network; NR: not reported; OR: odds ratio.

^a Per 20-point increase in GPS; adjusted for NCCN risk group.

^b Per 20-point increase in GPS; adjusted for NCCN risk group and medical center.

^c Per 20-point increase in GPS; adjusted for NCCN risk group and age.

^d As a continuous variable, adjusted for age, prostate-specific antigen level, clinical Gleason score, and NCCN risk category.

^e Per 20-point increase in GPS; adjusted for Gleason score, magnetic resonance imaging score, and prostatespecific antigen level.

Table 14. Risk of Adverse Pathology With Oncotype DX Prostate

Overall AP Risk, %(NCCN Risk Level)	n	AP Risk, n (%)(GPS Less Favorable Group; n=5)	AP Risk, n (%)(GPS Consistent With Group; n=29)	AP Risk, n (%)(GPS More Favorable Group; n=18)
0% (very low)	2	-	0	-
32% (low)	34	5 (100)	6 (21)	0
71% (low-	14	-	10 (34)	0
intermediate)				

Adapted from Whalen et al (2016).49,

AP: adverse pathology; GPS: Genomic Prostate Score; NCCN: National Comprehensive Cancer Network.

Systematic Reviews

Brand et al (2016) combined the Klein et al (2014) and Cullen et al (2015) studies using a patientspecific meta-analysis.^{55,} The GPS was compared with the CAPRA score, NCCN risk group, and AUA risk group. Reviewers tested whether the GPS added predictive value for the likelihood of favorable pathology above the clinical risk assessment tools. The model including the GPS and CAPRA score provided the best risk discrimination; the AUC improved from 0.68 to 0.73 by adding the GPS to the CAPRA score. The AUC improved from 0.64 to 0.70 by adding the GPS to the NCCN risk group. The improvements were reported to be significant, but the CIs for AUC were not provided.

Tables 15 and 16 display notable limitations identified in each study. This information is synthesized as a summary of the body of evidence and provides the conclusions on the sufficiency of the evidence supporting the position statement.

Т	able	15.	Study	Relevance	Limitations

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Klein et al	4. All patients			1. Survival outcomes	
(2014) ^{47,} validation	had RP			not included	
study					

2.04.111	Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management
Page 18 of 54	4

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Cullen et al (2015) ^{48,}	4. All patients had RP		3. No comparison to other risk predictors	1. Survival outcomes not included	1. 10-y outcomes not provided
Whalen et al (2016) ^{49,}	4. All patients had RP			1. Survival outcomes not included	1. 10-y outcomes not provided
Van Den Eeden et al (2018) ^{50,}	4. All patients had RP				
Salmasi et al (2018) ^{51,}	4. All patients had RP			1. Survival outcomes not included	1. Follow-up duration unclear

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

RP: radical prostatectomy.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest. ^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

Table 16. Study Design and Conduct Limitations

Study	Selection ^a	Blinding ^b	Delivery of Test ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
Klein et al (2014) ^{47,} validation study						1. Cls for reclassification not provided
Cullen et al (2015) ^{48,}						1. Cls for AUC and reclassification not provided
Whalen et al (2016) ^{49,}						1. Cls for reclassification not provided
Van Den Eeden et al (2018) ^{50,}						
Salmasi et al						

(2018)^{51,}

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

AUC: area under the curve; CI: confidence interval.

^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

^b Blinding key: 1. Not blinded to results of reference or other comparator tests.

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 19 of 54

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

Did not identify any studies that directly supported the clinical utility of Oncotype DX Prostate.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Decision-impact studies have assessed the potential impact of Oncotype DX Prostate on physicians' and patients' treatment decisions.^{56,57,58,} As with the previously evaluated test, given the lack of established clinical validity and no reported outcomes, it is uncertain whether any treatment changes were clinically appropriate. Decision-impact studies have also indicated that men classified as low-risk by guidelines criteria, and thus meeting guidelines criteria for active surveillance, are more likely to receive active surveillance if they are *tested* with the Oncotype DX Prostate test.^{57,59,60,} These arguments would suggest that the test may be a useful behavioral modifier. However, comparison with educational or quality improvement initiatives designed to improve the uptake of active surveillance in low-risk men has not been provided.

Klein et al (2014)^{47,} reported a decision-curve analysis^{61,} that they proposed reflects the clinical utility of Oncotype DX Prostate. In this analysis, they compared the predictive impact of the GPS plus the CAPRA validated tool^{62,} with the CAPRA score alone on the net benefit for the outcomes of patients with high-grade disease (Gleason score >4+3), high-stage disease, and combined high-grade and high-stage disease. They reported that, over a range of threshold probabilities for implementing treatment, "...incorporation of the GPS would be expected to lead to fewer treatments of patients who have favorable pathology at prostatectomy without increasing the number of patients with adverse pathology left untreated." For example, at a threshold risk of 40% (e.g., a man weighing the harms of prostatectomy vs. the benefit of active surveillance at 4:6), the test could identify 2 per 100 men with a high-grade or high-stage disease at a fixed false-positive rate, compared with using the CAPRA score alone. Thus, an individual patient could use the findings to assess his balance of benefits and harms (net benefit) when weighing the choice to proceed immediately to curative RP with its attendant adverse sequelae, or deciding to enter an active surveillance program. The latter would have an immediate benefit realized by forgoing RP but might be associated with greater downstream risks of disease progression and subsequent therapies. However, no CIs were presented for the decision-curve analysis.

Section Summary: Oncotype DX Prostate

The evidence from 5 studies on clinical validity for Oncotype DX Prostate has suggested the GPS can reclassify a patient's risk of recurrence or risk of adverse pathology at RP based on a biopsy specimen.^{47,48,49,} One study provided a figure with data on the reclassification of disease-specific survival using NCCN and GPS.^{50,} Ten-year prostate cancer death appears to be close to zero for men who are NCCN low-risk regardless of GPS score, indicating little useful reclassification of NCCN low-risk men based on GPS. For NCCN intermediate-risk, the risk of prostate cancer death ranges from approximately 0 for a GPS of less than 40 to close to 40% for a GPS of 100. It is unclear how many of the men with a GPS less than 40 were NCCN favorable intermediate-risk. Moreover, generalizing this evidence to a true active surveillance population, for which most in the study would be otherwise eligible, is difficult because all patients had elective RP. Thus, the findings do not reflect a clinical

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 20 of 54

scenario of predicting the risk of 10 year disease-specific survival in untreated patients under active surveillance. Some publications also lacked precision estimates for important variables such as risk estimates for recurrence or AUC estimates.

No direct evidence of clinical utility was found. The chain of evidence is also incomplete. Klein et al (2014) decision-curve analyses have suggested the potential for the combined GPS and CAPRA score data to help patients make decisions based on relative risks associated with immediate treatment or deferred treatment (i.e., active surveillance). This would reflect the clinical utility of the test. However, it is difficult to ascribe possible clinical utility of Oncotype DX Prostate in active surveillance because all patients regardless of clinical criteria elected RP within 6 months of diagnostic biopsy. Moreover, the validity of using tumor pathology as a surrogate for cancer-specific death is unclear. Reports from validation studies lack precision estimates for important variables such as risk estimates for recurrence.

The ProtecT trial showed 99% 10-year disease-specific survival in all 3 treatment groups: active surveillance, RT, and RP, including predominately low-risk but also some intermediate-risk men. AUA has recommended active surveillance in low-risk men. The low mortality rate estimated with tight precision makes it unlikely that a test intended to identify a subgroup of low-risk men with a net benefit from treatment instead of active surveillance would find such a group.

The PIVOT trial preplanned subgroup analysis showed a reduction in mortality for RP compared with observation for men at intermediate-risk; AUA has recommended RT or RP for such men. For intermediate-risk men, a test designed to identify men who can receive active surveillance instead of RP or RT would need to show very high NPV for disease-specific mortality at 10 years and improvement in prediction compared with existing tools used to select such men. For these men to forgo evidence-based beneficial treatment, there would have to be a very high standard of evidence for the clinical validity of the test.

Decipher Biopsy

This section reviews Decipher for initial management decisions in men with newly diagnosed, localized prostate cancer.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Three retrospective cohort studies reporting the clinical validity of Decipher Biopsy in men with newly diagnosed, localized prostate cancer are summarized in Tables 17 and 18.

Table 17. Characteristics of Clinical Validity Studies Assessing Decipher for Initial Management

Study	Study Population	Design	Comparator	Outcome	Sites	Dates
Berlin et al (2018) ^{63,}	Intermediate-risk PCa treated with curative-intent dose-escalated image-guided RT without neoadjuvant, concomitant or adjuvant ADT	Retrospective cohort from registry	NCCN risk groups	BCR, metastasis (5y)	Tertiary care center, probably in Ontario	2005- 2011
Nguyen et al (2017) ^{64,}	Treated with first-line RP or first-line RT plus ADT, had adverse pathology at surgery (defined as either preoperative PSA >20 ng/mL, stage pT3 or margin-positive, or RP grade group \geq 4), the vast majority of whom had presented with intermediate- or high-risk PCa	Retrospective cohort from manufacturer database	NCCN risk groups; clinical nomogram (CAPRA)	Metastases (5 y)	7 tertiary referral clinics including Cleveland Clinic, Johns Hopkins	1987- 2014

Study	Study Population	Design	Comparator	Outcome	Sites	Dates
Tosoian	High-risk prostate cancer, defined	Retrospective cohort	NCCN risk	Metastases	11 centers	1995-
et al	as clinical stage T3a, Grade Group		groups;	(5y)		2005
(2021) ^{65,}	4-5, or PSA >20 ng/ml. Patients had		CAPRA			
	undergone RP or RT with ADT.					

ADT: Androgen deprivation therapy; BCR: biochemical recurrence; CAPR : Cancer of the Prostate Risk Assessment I; NCCN: National Comprehensive Cancer Network; PCa: prostate cancer; RP: radical prostatectomy; RT: radiotherapy.

The cumulative incidence of metastases at 5 years by risk group is shown in Table 18.

Table 18. Reported Prognostic Accuracies for Metastasis or PC Mortality of Decipher as a Continuous Score and Comparators

Study	Outcome	AHR/AOR (95% CI) for Association Between GC and Outcome	AUC (95% CI)		
			GC	Comparator	GC + Comparator
Berlin (2018) ^{63,}	Metastasis (5 y)	2.1 (1.2 to 4.2)	0.86 (NR)	0.54 (NR)ª	0.89 (NR)
Nguyen (2017) ^{64,}	Metastasis (5 y)	1.4 (1.1 to 1.8)	0.74 (0.63 to 0.83)	0.66 (0.53 to 0.77)ª	0.74 (0.66 to 0.82)°
Tosoian et al (2021) ^{65,}	Metastasis (5y)	1.33 per 0.1 unit (1.19 to 1.48)	NR	NCCN risk group: 0.46 (NR) CAPRA: 0.59 (NR)	GC + NCCN: 0.67 (NR) GC + CAPRA: 0.71 (NR)

AHR: adjusted hazard ratio; AOR: adjusted odds ratio; AUC: area under the curve; CI: confidence interval; GC: genomic classifier; NR: not reported; PCa: prostate cancer.

^a National Comprehensive Cancer Network risk categories.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No published studies on the clinical utility of the Decipher test were identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Section Summary: Decipher Biopsy

For individuals who have clinically localized untreated prostate cancer who receive Decipher Biopsy, the evidence includes retrospective cohort studies of clinical validity using archived samples in intermediate-risk and high-risk patients and no studies of clinical utility. Relevant outcomes include OS, disease-specific survival, QOL, and treatment-related morbidity. A test designed to identify intermediate-risk men who can receive active surveillance instead of RP or RT or high-risk men who can forego ADT would need to show very high NPV for disease-specific mortality at 10 years and improvement in prediction compared with existing tools used to select such men.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 22 of 54

ProMark Protein Biomarker Test

The ProMark assay includes 8 biomarkers that predict prostate pathology aggressiveness and lethal outcomes: *DERL1, PDSS2, pS6, YBX1, HSPA9, FUS, SMAD4*, and *CUL2*. The assay results are combined using predefined coefficients for each marker from a logistic regression model to calculate a risk score. A risk score is a continuous number between 0 and 1, which estimates the probability of "non-GS 6" pathology.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Blume-Jensen et al (2015) reported on a study of 381 biopsies matched to prostatectomy specimens used to develop an 8-biomarker proteomic assay to predict prostate final pathology on prostatectomy specimen using risk scores.^{66,}

Biomarker risk scores were defined as favorable if less than or equal to 0.33 and nonfavorable if greater than 0.80, with a possible range between 0 and 1 based on false-negative and false-positive rates of 10% and 5%, respectively. The risk score generated for each patient was compared with 2 current risk stratification systems³/₄NCCN guideline categories and the D'Amico system. Results from the study showed that, at a risk score of less than or equal to 0.33, the predictive values of the assay for favorable pathology in very low- and low-risk NCCN and low-risk D'Amico groups were 95%, 81.5%, and 87.2%, respectively, while the NCCN and D'Amico risk classification groups alone had predictive values of 80.3%, 63.8%, and 70.6%, respectively. The positive predictive value for identifying favorable disease with a risk score of less than or equal to 0.33 was 83.6% (specificity, 90%). At a risk score greater than 0.80, 77% had nonfavorable disease. Overall, 39% of the patients in the study had risk scores less than or equal to 0.33 or greater than 0.8, 81% of which were correctly identified with the 8-biomarker assay. Of the patients with intermediate-risk scores (>0.33 to \leq 0.8), 58.3% had favorable disease.

The performance of the assay was evaluated in a second blinded validation study of 276 cases (see Table 19), also reported in Blume-Jensen et al (2015), to validate the assay's ability to distinguish "favorable" pathology (defined as Gleason score on prostatectomy \leq 3+4 and organ-confined disease) from "nonfavorable" pathology (defined as Gleason score on prostatectomy \geq 4+3 or non-organ-defined disease). The second validation study separated favorable from nonfavorable pathology (AUC=0.68; 95% CI, 0.61 to 0.74).

Table 19. Clinical Validity of ProMark

Study	Designª	Outcome	Site	Ν
Blume-Jensen et al (2015) ^{66,}	Retrospective cohort ^a	Favorable pathology at RP	Montreal, QC	276ª
RP: radical prostatectomy.				

^a Only the validation sample cohort N.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No published studies on the clinical utility of the ProMark test were identified.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 23 of 54

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical utility of the ProMark test has not been established, a chain of evidence supporting the test's clinical utility cannot be constructed.

Section Summary: ProMark Protein Biomarker Test

Data are insufficient to establish the clinical validity or the clinical utility of the ProMark test.

Management Decision after Radical Prostatectomy

Clinical Context and Test Purpose

The purpose of GEP and protein biomarker testing in patients who have prostate cancer and who have undergone RP is to inform management decisions.

For example, the optimal timing of RT after RP is debated. Adjuvant RT may maximize cancer control outcomes; early salvage RT (at first evidence of a rising serum PSA level) can minimize overtreatment and still lead to acceptable oncologic outcomes.^{66,} Adjuvant RT in men with pT3 or margin-positive cancer has been compared with observation in RCTs; such comparisons have shown that adjuvant RT improves the biochemical and local control rates among patients with adverse pathology at RP.^{67,68,} Although the observation arms in these trials included men who received adjuvant therapy, the trials did not directly compare early salvage RT with immediate adjuvant RT because they included varying or unspecified thresholds for the initiation of salvage therapy RT.

Several observational analyses have shown conflicting conclusions whether adjuvant RT is favored over early salvage RT.^{67,69,68,} RCTs comparing adjuvant with early salvage RT are underway. Guidelines have recommended that adjuvant RT be offered to patients with adverse pathologic findings at RP, and salvage RT is offered to patients with PSA or local recurrence after RP.^{14,70,} However, many men treated with RT will never experience recurrence after surgery and therefore receive no benefit while experiencing harm from RT. Therefore, a test that could be used to identify men who meet criteria for adjuvant or early salvage RT but can safely receive observation instead would be useful.

Other post-RP clinical questions for which GEP or protein biomarker testing might be useful is in guiding systemic treatment (ADT and/or chemotherapy) in men receiving RT.

The second question addressed in this evidence review is: Does GEP or protein biomarker testing, compared with clinicopathologic risk stratification or when used with clinicopathologic risk stratification, improve outcomes in men following RP?

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals who have undergone RP for prostate cancer, and who are deciding on subsequent management such as adjuvant RT or no adjuvant RT. The Decipher results report says that "Decipher is intended for use in those patients who present with specific risk factors for the recurrence of prostate cancer after radical prostatectomy: (1) stage T2 disease with positive surgical margins, or (2) stage T3 disease, or (3) rising prostate-specific antigen (PSA) levels after initial PSA nadir."

Interventions

Prolaris, described in the previous section, is also intended to classify individuals who have undergone RP.

Decipher is a tissue-based tumor 22-biomarker GEP test intended to classify high-risk individuals who have undergone RP. The cutpoints 0.45 and 0.60 are used to categorize men using a low-, intermediate-, and high-risk genomic classifier (GC) on the Decipher test results report.

Comparators

Clinicopathologic risk stratification is currently being used to make decisions about prostate cancer management following RP. Clinical characteristics (e.g., stage, biopsy Gleason grade, serum PSA level, surgical margin, disease involvement) and demographic characteristics (e.g., age, life expectancy) are combined to classify men according to risk. As described previously, NCCN and AUA provide risk stratification guidelines.^{12,14}, The Stephenson nomogram^{71,72}, and Cancer of the Prostate Risk Assessment-Surgical (CAPRA-S) nomogram^{73,} can be used to predict outcomes after RP.

Outcomes

Beneficial outcomes resulting from a true test result are prolonged survival, improved QOL, and reduction in unnecessary treatment-related adverse events. Harmful outcomes resulting from a false test result are recurrence, metastases or death, and unnecessary treatments. The outcomes of interest are listed in Table 20.

Outcome	Details
Overall survival	10-year survival
Disease-specific survival	10-year prostate cancer-free survival; 10-year prostate cancer death rate; 10-year recurrence rate
Quality of life	See Chen et al (2014) ^{37,} for NCI-recommended health-related quality of life measures for localized prostate cancer
Treatment-related morbidity	Adverse events of radiotherapy or radical prostatectomy
NCI: National Cancor Instituto	

Table 20. Outcomes of Interest for Individuals After Radical Prostatectomy

NCI: National Cancer Institute.

Ten-year outcomes are of interest due to the prolonged natural history of prostate cancer and the low number of events observed.

Prolaris

Prolaris used for initial management decisions was described in the previous section. This section reviews Prolaris for management after RP.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Five studies reporting clinical validity in the post-RP management setting are summarized in Table 21. Four of these studies: Cuzick et al (2011),^{74,} Cooperberg et al (2013),^{52,},Bishoff et al (2014)^{75,}and Swanson et al (2021)^{76,} reported on post-RP patients. Koch et al (2016)^{77,} reported on post-RP patients with BCR. Freedland et al (2013)^{78,} reported on post-RT patients but is included in this section for completeness.

Table 21.	Clinical Validity	Studies Assessing	Prolaris for Post-l	RP or Post-RT	Management

		5			
Study	Design	Population	Dates	Sites	N
After					
prostatectomy					
Cuzick et al (2011) ^{74,}	Retrospective cohort from prospective registry	Clinical stage T1/T2; no neoadjuvant therapy; 71% PSA level ≤10 ng/mL, 96% Gleason score ≤7	1985-1995	Scott and White Clinic	366
Cooperberg et al (2013) ^{52,}	Retrospective cohort from	98% PSA level ≤20 ng/mL, 95% Gleason score ≤7; no neoadjuvant or adjuvant therapy	1994-2011	UCSF Registry	413

Study	Design	Population	Dates	Sites	Ν
	prospective registry				
Bishoff et al (2014) ^{75,}	Retrospective cohort from	Clinical stage T1/T2; median PSA level 5.5-7.2 ng/mL; between 91% and 94%	2005-2006	Martini Clinic	283
	medical records	Gleason score ≤7; between 3% and 19% with adjuvant therapy	1994-2005	Durham VAMC	176
			1997-2004	Intermoun tain Healthcar e	123
Koch et al (2016) ^{77,}	Retrospective cohort from medical records	Median PSA level 6.5-11 ng/mL; 64% Gleason score ≤7; no adjuvant RT	1995-2010	Indiana University SOM	47
Swanson et al (2021) ^{76,}	Retrospective cohort from prospective registry	46% considered to have a low risk of disease progression, 35% to have an intermediate risk, and 19% high risk according to CAPRA-S	1985-1997	Scott and White hospital (Temple, TX)	360
After external- beam radiotherapy					
Freedland et al (2013) ^{78,}	Retrospective cohort, source unclear	97% clinical stage T1/T2; Median PSA level 8 ng/mL; 88% Gleason score ∈7; 53% no concurrent hormone use; 57% African American	1991-2006	Durham VAMC	141

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 25 of 54

CAPRA-S: Cancer of the Prostate Risk Assessment Postsurgical; PSA: prostate-specific antigen; RP: radical prostatectomy; RT: radiotherapy; SOM: School of Medicine; UCSF: University of California, San Francisco; VAMC: Veterans Affairs Medical Center.

Cuzick et al (2011) examined the potential use of the Prolaris CCP test combined with a clinical score following RP, using a retrospective cohort of archived samples from a tumor registry.^{74,} The study also included a cohort of men with localized prostate cancer detected from specimens obtained during transurethral resection of the prostate, which is not a population of interest here, and so is not described. Men conservatively managed after RP between 1985 and 1995 were identified from a tumor registry (n=366 with CCP scores). The primary endpoint was time to BCR, and the secondary endpoint was prostate cancer death. Myriad Genetics assessed CCP scores blindly. The median age of patients was 68 years (median follow-up, 9.4 years). Gleason scores were 7 or lower in 96%, but margins were positive in 68%. Cancers were clinically staged as T3 in 34%; following RP, 64% was judged pathologic stage T3. CCP score was associated with BCR (see Table 15). Analyses of prostate cancer deaths in the RP cohort were problematic, due to only 12 (3%) deaths. The clinical score included PSA level, stage, positive surgical margins, and Gleason score. The AUC for BCR within 5 years in the RP cohort was 0.825 for the clinical score and 0.842 for the CCR score. Although the CCP increased the AUC by 2%, whether that improvement is clinically useful is unclear because reclassification data and analysis of net benefits are lacking.

Swanson et al (2021) published a reanalysis of 360 patients from the cohort first reported in Cuzick et al (2011).⁷⁶ After a median follow-up of 16 years, 163 (45%) of the cohort developed BCR, 41 (11%) developed metastatic disease, and 33 (9%) died from prostate cancer. The CCR score (a combination of CAPRA-S and the CCP molecular score) was prognostic of prostate cancer death, but the estimate was imprecise (HR per unit score, 3.40; 95%CI, 1.52 to 7.59). The study authors illustrated the added value of CCR for predicting disease-specific mortality by comparing predicted risk using CCR to risk predicted by a CAPRA-S-only model in a Kaplan-Meier curve; however, precision estimates were not presented.

Cooperberg et al (2013) evaluated the CCP score in an RP cohort and the incremental improvement over the CAPRA-S score for predicting BCR using a prospective-retrospective design (conforming to

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 26 of 54

a PRoBE study design).^{52,} A prognostic model was developed from the RP cohort described by Cuzick et al (2011).^{74,} The validation cohort was obtained from patients identified from the UCSF Urologic Oncology Database. Tissue sufficient to obtain a CCP score was available for 413 men (69% of the 600 eligible samples). Both UCSF and Myriad Genetics performed statistical analyses. In the validation cohort, 95% had Gleason scores of 7 or lower, 16% of samples had positive margins, 4% had seminal vesicle invasion, and 23% had extracapsular extension. BCR occurred in 82 (19.9%) men. The association with BCR is shown in Table 22. The AUC for BCR with CAPRA-S alone was 0.73, increasing to 0.77 for the combined CCR score.

Bishoff et al (2014) examined the prognostic ability of the CCP score in 3 cohorts: the Martini Clinic (n=283, simulated biopsies from formalin-fixed paraffin-embedded RP specimen), Durham Veterans Affairs Medical Center (n=176, diagnostic biopsies), and Intermountain Healthcare (n=123, diagnostic biopsies).^{75,} The combined analysis included all 582 patients. Gleason scores were 7 or lower in 93% of men. In the combined cohorts, a unit increase in the CCP score increased the adjusted HR for BCR by 1.47 (see Table 22). Metastatic events (n=12) were too few to draw conclusions.

Koch et al (2016) evaluated whether the CCP score could discriminate between systemic disease and local recurrence in patients with BCR after RP.^{77,} All 60 patients given RP as primary therapy at an academic medical center between 1995 and 2010 for whom samples were available and who had a BCR and either developed metastatic disease or received salvage EBRT with at least 2 years of follow-up were eligible for retrospective analysis. Data from 5 patients were excluded for failing to meet clinical eligibility requirements (no clarification provided) or because data were incomplete; sample blocks from 3 patients contained insufficient tumor for assay and data from 6 patients were excluded due to lack of "passing" CCP scores. Forty-seven patients were included in the analysis. Outcomes were classified into 3 categories: (1) metastatic disease (n=22), (2) nonresponse to salvage EBRT (n=14), and (3) durable response to salvage EBRT (n=11). Analyses were performed with a binary outcome (categories 1 and 2 combined). For each 1-unit change in the CCP score, the univariate odds ratio for metastatic disease or nonresponse was 3.72 (see Table 22). Multivariate analysis was performed; however, due to the very small number of participants in the durable response group, CIs were very wide.

Study	Outcomes	Median FU, y	Ν	Unadjusted	Multivariate
				Ratio (95% CI)	Ratio (95% CI)
Cuzick et al (2011) ^{74,}	BCR	9.4	366	HR=1.89 (1.54 to 2.31)	1.77 (1.40 to 2.22)¤
	Prostate cancer death		337	HR=2.92 (2.38 to 3.57)	2.56 (1.85 to 3.53) ^b
Cooperberg et al (2013)52,	BCR	7	413	HR=2.1 (1.6 to 2.9)	1.7 (1.3 to 2.4) ^c
Bishoff et al (2014) ^{75,}	BCR	5/7 ^f	582	HR=1.60 (1.35 to 1.90)	1.47 (1.23 to 1.76) ^d
Koch et al (2016) ^{77,}	Metastatic disease or	9.4	47	OR=3.72 (1.29 to 10.7)	10.4 (2.05 to 90.1) ^e
	nonresponse				
Swanson et al (2021) ^{76,}	Prostate cancer death	16	360	HR=2.11 (1.68 to 2.65)	3.40 (1.52 to 7.59) ^c

Table 22. Univariate and Multivariate Associations Between Prolaris CCP and Outcomes in Pos	st-
RP Clinical Validation Studies	

BCR: biochemical recurrence; CCP: Cell Cycle Progression; CI: confidence interval; FU: follow-up; HR: hazard ratio; OR: odds ratio; PSA: prostate-specific antigen; RP: radical prostatectomy.

^a Per 1-unit increase in CCP. Adjusted for PSA level, Gleason score, pathologic T stage and grade, positive surgical margins, extracapsular extension, bladder involvement, seminal vesicle involvement, positive lymph node, and age.

^b Per 1-unit increase in CCP. Adjusted for Gleason score, PSA level, Ki67, and cancer extent.

^c Per 1-unit increase in CCP. Adjusted for Cancer of the Prostate Risk Assessment-Surgical.

^d Per 1-unit increase in CCP. Adjusted for PSA level, Gleason score, and adjuvant treatment.

^e Per 1-unit increase in CCP. Adjusted for Gleason score, time from surgery to BCR, and PSA level.

^f Not reported for 3 cohorts.

Although not a study of management post-RP, Freedland et al (2013) described the prognostic ability of the CCP score for predicting BCR in men who received primary EBRT.^{79,} The retrospective data included 141 men diagnosed with prostate cancer who were treated with EBRT from 1991 to 2006,

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 27 of 54

with biopsy samples and follow-up of at least 3 years. Nineteen (13%) men experienced BCR by 5 years. The univariate HR for BCR for each 1-unit increase in CCP was 2.55 (95% Cl, 1.43 to 4.55). The multivariable HR for BCR associated with a 1-unit increase in CCP, including adjustment for pretreatment PSA level, Gleason, percent positive cores, and concurrent ADT, was 2.11 (95% Cl, 1.05 to 4.25).

Systematic Reviews

As described in the previous Prolaris section, results of an industry-sponsored systematic review and meta-analysis were reported.^{80,} Seven published studies were identified; all have been reviewed in the previous paragraphs (needle biopsy conservative management cohorts, postprostatectomy cohorts, and EBRT cohort). Including 4 validity studies^{74,52,75,78,} that reported outcomes of BCR in post-RP cohorts, the pooled estimate of the HR, calculated with random-effects meta-analytic methods, for BCR for a 1-unit increase in CCP score was 1.9 (95% CI, 1.6 to 2.3). Two studies reported outcomes for disease-specific mortality.^{38,74,} Since only one of those was a post-RP study, the pooled HRs are not relevant here. There was evidence of heterogeneity in both models; reviewers did not report any variables associated with heterogeneity.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

Did not identify any studies directly supporting the clinical utility of Prolaris.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Decision Curves

In a decision-curve analysis, Cooperberg et al (2013) found the CAPRA-S score superior to CCP alone (as well as treat-none or treat-all strategies) in men after prostatectomy.^{52,} A combined CCR predictor appeared only slightly better than CAPRA-S alone for thresholds of approximately 30% or more. For example, at a threshold of 30% (i.e., meaning a man would value the harm-to-benefit of treatment such as RT as 3:7), the CCR score would detect about 2 more men per 100 likely to experience BCR if the false-positive rate was fixed. However, the lack of CIs for the decision-curve analysis, together with the small difference, is consistent with an uncertain net benefit obtained by adding CCP to the CAPRA-S score. Also, it is not clear whether the group of patients identified as high-risk of experiencing BCR would have a net benefit from adjuvant instead of early salvage RT.

Section Summary: Prolaris

Five identified studies examined the clinical validity of Prolaris in men after RP using a BCR or systemic disease endpoint. Cuzick et al (2011) found that the CCP score offered little improvement in the AUC (2%) over clinicopathologic predictors and did not examine reclassification.^{74,} Cooperberg et al (2013) found the AUC for BCR improved from 0.73 (CAPRA-S alone) to 0.77 by adding CCP score.^{52,} Bishoff et al (2014)^{75,} and Koch et al (2016)^{77,} did not report any classification or discrimination measures. Koch et al (2016) was performed in patients who had a BCR following RP. Swanson et al (2021) published a reanalysis of 360 patients from the cohort first reported in Cuzick et al (2011).^{76,}After a median follow-up of 16 years 163 (45%) of the cohort developed biochemical recurrence, 41 (11%)

developed metastatic disease, and 33 (9%) died from prostate cancer. The CCR score was prognostic of prostate cancer death but the estimate was imprecise (HR per unit score, 3.40; 95% CI, 1.52 to 7.59). No direct evidence is available to support the clinical utility of Prolaris for improving net outcomes of patients with localized prostate cancer following RP. The chain of evidence is also incomplete. Decision-curve analysis did not provide convincing evidence of meaningful improvement in net benefit by incorporating the CCP score.

Prolaris CCP score may have an association with BCR, but disease-specific survival outcomes were reported in only one analysis. A larger number of disease-specific survival events and precision estimates for discrimination measures are needed.

Decipher Prostate RP

Decipher used for initial management decisions was described in the previous section. This section reviews Decipher for management after RP.

The Decipher test classifies as low-risk those patients who can delay or defer RT after prostatectomy, or as high-risk those who would potentially benefit from early radiation. The GC is a continuous risk score between 0 and 1, with higher risk scores indicating a greater probability of developing metastasis.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

The clinical validity of the Decipher test (GC) has been reported in multiple studies to predict metastasis, mortality, or BCR after RP in men with postoperative high-risk features like pathologic stage T2 with positive margins, pathologic stage T3 disease, or a rising PSA level (see Tables 23 and 24).^{81,82,79,83,84,85,86,87,88,89,90,91,}

			, ,			
Study	Study Population	Design	Comparator	Outcome	Sites	Dates
Feng et al (2021) ^{81,}	Recurrent disease after RP with a PSA of 0.2-4.0 ng/mL, pathologic T3 disease (tumor spread beyond the prostate) or T2 disease (tumor contained within the prostate) with a positive surgical margin and no evidence of nodal or metastatic disease	Ancillary study of specimens from an RCT	Standard clinicopathologic variables	Distant metastasis (primary), prostate cancer death and OS (secondary)	Multiple sites in US and Canada	1998- 2003
Spratt et al (2018) ^{92,}	Clinically localized PCa after RP; serious PSA levels post-RP documented; no neoadjuvant ADT; 31% with detectable PSA 8 wk post-RP	Retrospect ive cohort from registry	Clinicopathological risk factors (e.g., preop PSA, SM, RP grade group)	Metastases (5 y)	MD Anderson, Durham VA, Thomas Jefferson	1990- 2015
Karnes et al (2018) ^{93,}	Clinically localized PCa after RP; pathologic GS ≥7,	Retrospect ive cohort	Clinicopathologic risk factors (e.g., preop	PCa mortality (10 y)	Mayo Clinic, Johns Hopkins,	1987- 2010

Table 23. Characteristics of Clinical Validity Studies Assessing the Decipher Genomic Classifier

Study	Study Population	Design	Comparator	Outcome	Sites	Dates
	pT3, pN1, or margin- positive; no neoadjuvant treatment; ≥10 y follow-up for patient alive	from registry	PSA, EPE, GS); clinical nomogram (CAPRA-S)		Cleveland Clinic, Durham VA	
Freedland et al (2016) ^{89,}	Clinically localized PCa after RP; received postoperative SRT; pathologic node- negative disease; undetectable post- RP PSA; no neoadjuvant or adjuvant treatment; 32% African American	Retrospect ive cohort from registry	Clinicopathologic risk factors (e.g., preop PSA, EPE, GS); Clinical nomogram (Briganti, CAPRA-S)	Metastases	Durham VA, Thomas Jefferso n, Mayo Clinic	1991- 2010
Glass et al (2016) ^{90,}	Clinically localized PCa after RP; preop PSA >20 ng/mL, stage pT3, margin- positive, or pathologic GS ≥8; no neoadjuvant or adjuvant treatment; 2% African American	Retrospect ive cohort from registry	Clinical risk factors (age at diagnosis); Clinical nomogram (CAPRA-S)	Clinical recurrence (10 y)	Kaiser Permanente Northwest	1997- 2009
Ross et al (2016) ^{94,}	Clinically localized PCa after RP; CAPRA-S score ≥3, pathologic GS ≥7, post-RP PSA nadir <0.2 ng/mL, and sufficient tissue and clinical data; no nodal disease prior to surgery; no treatment before metastasis; 8% African American	Case cohort fro m registry	Clinicopathologic risk factors (e.g., preop PSA, EPE, GS); clinical nomogram (CAPRA-S, Eggener)	Metastases (10 y)	Johns Hopkins	1992- 2010
Ross (2016) ^{94,}	Clinically localized PCa after RP; stage pT3 or margin- positive; achieve PSA nadir after surgery; no node- positive; no neoadjuvant treatment; no hormone-only treatment prior to metastasis; no SRT for PSA >10 ng/mL	Retrospect ive cohort from registry	Clinical variables (e.g., ART, MRD-SRT, SRT, no-RT); clinical nomogram (CAPRA-S)	Metastasis (10 y)	Mayo Clinic, Johns Hopkins, Durham VA, Thomas Jefferson	1990- 2010
Cooperberg et al (2015) ^{84,}	Clinically localized PCa after RP; preop PSA >20 ng/mL, stage pT3b, or pathologic GS ≥8;	Case cohort fro m registry	Clinicopathologic risk factors (e.g., preop PSA, EPE, GS); clinical nomogram (CAPRA-S)	PCa mortality	CapSURE Registry	2000- 2006

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 29 of 54

Study	Study Population	Design	Comparator	Outcome	Sites	Dates
	no neoadjuvant treatment; achieve PSA nadir after surgery					
Den et al (2015) ^{82,}	Clinically localized PCa after RP; pT3 or margin-positive disease; received post-RP RT; no neoadjuvant treatment; no lymph node invasion	Retrospect ive cohort from registry	Clinicopathologic risk factors (e.g., preop PSA, EPE, GS); clinical nomogram (CAPRA-S)	Metastases	Thomas Jefferson, Mayo Clinic	1990- 2009
Klein et al (2015) ^{79,} ; Klein et al (2016) ^{91,}	Clinically localized PCa after RP; preop PSA >20 ng/mL, stage pT3, margin- positive or pathologic GS ≥8; pathologic node- negative disease; undetectable post- RP PSA; no neoadjuvant or adjuvant treatment; ≥5 y follow-up for censored patients; 8% African	Retrospect ive cohort from registry	Clinicopathologic risk factors (e.g., pre-op PSA, EPE, GS); clinical nomogram (Stephenson, CAPRA- S)	Metastases (5 y, 10 y)	Cleveland Clinic	1993- 2001
Den et al (2014) ^{83,}	Clinically localized PCa after RP; pT3 or margin-positive disease; received post-RP RT; no neoadjuvant treatment; 39% BCR; 13% African American	Retrospect ive cohort from registry	Clinicopathologic risk factors (e.g., preop PSA, EPE, GS); clinical nomogram (Stephenson, CAPRA- S)	BCR	Thomas Jefferson	1999- 2009
Ross et al (2014) ^{85,a} (BC R only)	Clinically localized PCa with BCR after RP; preop PSA >20 ng/mL, pathologic GS \geq 8, SVI or Mayo Clinic nomogram score \geq 10; no neoadjuvant treatment	Case cohort fro m registry	Clinicopathologic risk factors (e.g., preop PSA, EPE, GS); clinical nomogram (Stephenson, CAPRA- S)	Metastases (5 y)	Mayo Clinic	2000-2006
Erho et al (2013) ^{87,} (valid ation)	Clinically localized PCa after RP; 32% no evidence of disease post-RP within 7 y of follow- up; 34% BCR post- RP with no clinical metastasis within 5 y of BCR; 34% clinical metastasis within 5 y of BCR	Nested case- control from registry	Clinicopathologic risk factors (e.g., preop PSA, EPE, GS)	Metastases	Mayo Clinic	1987- 2001
Karnes et al (2013) ^{86,}	Clinically localized PCa after RP; preop	Case cohort	Clinicopathologic risk factors (e.g., preop	Metastases (5 y)	Mayo Clinic	2000- 2006

2.04.111	Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management
Page 30 of 5	4

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 31 of 54

Study	Study Population	Design	Comparator	Outcome	Sites	Dates
	PSA >20 ng/mL,	from	PSA, EPE, GS); clinical			
	pathologic GS ≥8,	registry	nomogram			
	SVI or Mayo Clinic		(Stephenson)			
	nomogram score					
	≥10; no neoadjuvan	t				
	treatment					

ART: adjuvant radiotherapy; CARPA-S: Cancer of the Prostate Risk Assessment Postsurgical; BCR: biochemical recurrence; EPE: extraprostatic extension; GS: Gleason Score; MRD: minimal disease residual; PCa: prostate cancer; preop: preoperative; RP: radical prostatectomy; RT: radiotherapy; SM: surgical margins; SRT: salvage radiotherapy; SVI: seminal vesicle invasion.

^a Appears to be subgroup with BCR from Karnes et al (2013).

Table 24. I	Reported Progno	stic Accuracies for Metastasis or PC Mortality of Decipher as a				
Continuou	Continuous Score and Comparators					
Study	Outcome	AHR/AOR (95% CI) AUC (95% CI)				

Study	Outcome	for Association Between GC and Outcome	AUC (95% CI)		
			GC	Comparator	GC + Comparator
Feng et al (2021) ^{81,}	MetastasisPCSMOS	 1.17 (1.05 to 1.832) p=.006 1.39 (1.20 to 1.63); p<.001 1.17 (1.06 to 1.29); p=.002 	NR	NR	NR
Spratt (2018) ^{92,} ;95% received RT	Metastasis	NR	0.86 (0.80 to 0.94)	0.69 (0.41 to 0.89) [⊾]	0.83 (0.70 to 1)
Karnes (2018) ^{93,}	PCa mortality	1.3 (1.2 to 1.5)	0.73 (0.67 to 0.78)	0.73 (0.68 to 0.78)	0.76 (0.71 to 0.82)
Freedland (2016) ^{89,}	Metastasis post-RT	1.6 (1.1 to 2.1)	0.85 (0.73 to 0.88)	0.65 (0.54 to 0.81) ^g	NR
Ross (2016) ^{94,}	Metastasis	1.3 (1.1 to 1.5)	0.76 (0.65 to 0.84)	0.77 (0.69 to 0.85) ^ь	0.87 (0.77 to 0.94)
Glass (2016) ^{90,}	Metastasis	1.5 (p=.011)	0.80 (0.64 to 0.92)	0.73 (0.49 to 0.95) ^c	0.84 (0.70 to 0.96)
Cooperberg (2015) ^{84,}	PCa mortality	1.8 (1.5 to 2.3)	0.78 (0.68 to 0.87)	0.75 (0.55 to 0.84) ^b	
Klein (2015) ^{79,} ; Klein (2016) ^{91,}	Metastasis 5 y Metastasis 10 y	1.5 (1.1 to 2.1); 1.7 (1.1 to 2.8)	0.77 (0.66 to 0.87); 0.80 (0.58 to 0.95)	0.75 (0.65 to 0.84) ^c ; 0.75 (0.64 to 0.87) ^h	0.79 (0.65 to 0.85) 0.88 (0.76 to 0.96)
Den (2015) ^{82,}	Metastasis post-RT	1.9 (p<.001)	0.78 (0.64 to 0.91)	0.70 (0.49 to 0.90) ^b	0.85 (0.79 to 0.93)
Ross (2014) ^{85,}	Metastasis	1.4 (p=.003)	0.82 (0.76 to 0.86)	0.70 (0.66 to 0.75)ª	0.75 (0.69 to 0.80)
Den (2014) ^{83,}	Metastasis	NR	0.70 (0.49 to 0.90) ^d	0.78 (0.64 to 0.91)	0.80 (0.68 to 0.93)
Erho (2013) ^{87,}	Metastasis	1.4 (p<0.001)	0.75 (0.70 to 0.81) ^e	0.69 (0.60 to 0.77) ^{a,e}	0.74 (0.65 to 0.82) ^{a,e}
Karnes (2013) ^{86,}	Metastasis	1.5 (p<0.001)	0.79 (0.68 to 0.87)	0.64 (0.55 to 0.72) ^{d,f}	

AHR: adjusted hazard ratio; AOR: adjusted odds ratio; AUC: area under the curve; Cl: confidence interval; GC: genomic classifier; NR: not reported; OS: overall survival; PCa: prostate cancer; PCSM: prostate-cancer specific mortality; RT: radiotherapy.

^a Clinical classifier includes Gleason score, extracapsular extension, positive surgical margins, seminal vesicle invasion, or lymph node involvement.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 32 of 54

^b Cancer of the Prostate Risk Assessment-Surgical.

- ^c Stephenson nomogram.
- ^d Only reported vs. single clinical predictors.
- ^e AUC CI obtained by digitizing figure.
- ^f Gleason score.
- ^g Briganti score.
- ^h National Comprehensive Cancer Network risk categories.
- With detectable PSA post-RP.

All studies were conducted retrospectively from registry data or clinical records. The development study had a nested case-control design.^{87,} The 5- and 10-year results of 1 study were published separately.^{79,91,} Four were case-cohort studies and 8 used retrospective cohorts. Nine studies were supported by GenomeDx (now Decipher Corp), which offers the Decipher test. The cutpoints used to classify men into low-, intermediate- and high-risk by GC score were updated in 2016. Only 1 study (Karnes et al [2018]^{93,}) has reported 10-year prostate cancer-specific survival after the update in the cutpoints.

Several studies,^{84,85,86,87,94,92,94,} including the test (validation) sample from the development study, examined men observed following RP and undergoing adjuvant or salvage RT. Median follow-up periods ranged from 6.4 to 16.9 years. The distributions of Gleason scores in the studies varied from 17.8% to 49.3% for those with Gleason scores of 8 or higher and from 0.4% to 15.1% for those with scores of 6 or lower. Extracapsular extension of the tumor ranged from 42.7% and 72.3% of men across studies.

Association between GC continuous score and metastasis or prostate cancer-specific mortality is shown in Table 25. The GC AUCs for predicting metastases are shown in Table 24. Among the 69 men developing metastases in Karnes et al (2013), of the 29 with Gleason scores of 7 or lower, 10 were correctly reclassified to the highest GC risk (score >0.6), but of the 40 men with Gleason scores of 8 or higher, 10 were incorrectly reclassified to the lowest GC risk group (score <0.4).^{86,}

The cumulative incidence of metastases by risk group is shown in Table 26. Three studies reported prostate cancer-specific mortality; only one of which included 10 year outcomes. Precision estimates were not provided. Values in the tables below may be estimated from figures when exact values were not provided in article text or tables.

Study	FU Time, v	N	Patient	s in Risk G	roup, %	Metasta	sis Rate, %	
			Low	Int	High	Low	Int	High
Feng et al (2021) ^{81,}	13	352	42	38	20	6.2	8.7	15.3
Spratt et al (2018) ^{92,}	10	561	46	28	26	0	3	23
Ross et al (2016) ^{94,}	5	422	57	27	16	7	10	22
Freedland et al (2016) ^{89,}	10	170	51	31	18	3	8	33
Glass et al (2016) ^{90,}	10	224	NR	NR	NR	0	3	
Ross et al (2016) ^{88,}	10	260	73	17	10	8	20	32
Klein et al (2015) ^{79,}								
Den et al (2015) ^{82,}	5	188	41	39	20	0	9	29
Den et al (2014) ^{83,}	5	139	21	38	41	0	5	17
Ross et al (2014) ^{85,}	5	85	NR	NR	NR	9	54	
Karnes et al (2013) ^{86,}	5	219	51	22	27	2	6	22

Table 25. Metastasis by GC Risk Group

FU: follow-up; GC: genomic classifier; Int: intermediate; NIIQRR: not reported. For prostate cancer mortality, compared with CAPRA-S, Cooperberg et al (2015) found that the GC improved reclassification somewhat-of the 19 men with CAPRA-S scores of 5 or lower, 12 were correctly reclassified to the highest GC risk, and 1 was incorrectly reclassified with a CAPRA-S score greater than 6 to low-risk; all men had CAPRA-S scores of 3 or

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 33 of 54

more.^{84,}Feng et al (2021) reported prostate specific mortality and OS according to GC category, but did not provide data on reclassification.^{81,}

Of note, Karnes et al (2018) reported the preferred outcome for this review (10-year prostate cancerspecific survival).^{93,} The authors found that adding the GC to CAPRA improved the AUC from 0.73 to 0.76 with highly overlapping CIs. The 10-year cumulative incidence of prostate cancer-specific mortality by CAPRA and GC risk categories are shown in Table 27. Samples sizes and precision estimates for the cross-tabulations were not provided.

Table 26. Prostate-Cancer-Specific Mortality by Genomic Classifier Risk Group

Study	FU, y	Ν	Patients i	n Risk Gr	oup, %	Prostate Ca	ncer Mor	tality
			Low	Int	High	Low	Int	High
Karnes et al (2018) ^{93,}	10	561	58	17	25	12	13	45
Cooperberg et al (2015) ^{84,}	5	185	54	22	24	6	3	30
Feng et al (2021) ^{81,}	13	352	42	38	20	0.7	2.4	9.8

FU: follow-up; Int: intermediate.

Table 27. Cross-Tabulation of 10 Year Cumulative Incidence of Prostate Cancer-Specific Mortality by GC and CAPRA

CAPRA-S Risk Category	Decipher GC Risk Category, %					
	Low/Intermediate (≤0.6)	High (>0.6)				
Low-risk (<6)	2.8 (CI NR)	18 (CI NR)				
High-risk (≥6)	5.5 (CI NR)	30 (CI NR)				

Adapted from Karnes et al (2018).^{93,}

CAPRA: Cancer of the Prostate Risk Assessment; CI: confidence interval; GC: genomic classifier; NR: not reported.

Tables 28 and 29 display notable limitations identified in each study. The limitations analysis focuses on 10-year prostate cancer-specific mortality outcomes (i.e., Karnes et al [2018]^{93,}).

Table 28. Study Relevance Limitations

Study	Populationa	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Karnes et al					
(2018) ^{93,}					

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

Table 29. Study Design and Conduct Limitations

Study	Selection ^a	Blinding ^b	Delivery	Selective	Data Completences	Statistical ^f
			of lest	Reporting	Completeness	
Karnes et	2. Unclear if included					1. Cls for prostate
al (2018) ^{93,}	men were					cancer-specific
	consecutive or					mortality by GC
	random samples of					low/high-risk and

Study	Selection ^a	Blinding ^b	Delivery of Test ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
	those meeting					reclassification not
	eligibility criteria					provided

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

CI: confidence interval.

^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

^b Blinding key: 1. Not blinded to results of reference or other comparator tests.

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication. ^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No studies reporting direct evidence were identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Decision Curves

Studies have included decision curves comparing the net benefit of different strategies using metastases or survival as the outcome (see Table 28).^{82,79,84,85,86,88,93,95,93,} In observational and RT samples from Karnes et al (2013)^{86,} and Ross et al (2014),^{85,} using a 15% to 25% range of thresholds for decision making (i.e., suspected probability of developing metastases) would be expected to identify correctly as few as no men or as many as 4 per 100 likely to experience metastases. This range of thresholds assumes several things: it assumes those making the decisions are relying on the GC result for adjuvant RT decisions, compared with treating based on the best comparator test, and it assumes no increase in false-positives. No CIs were provided for the net benefit estimates and uncertainty cannot be evaluated. In the 2 observation-only samples, although the GC improved the net benefit over a "treat none" strategy over 15% to 25% thresholds, it appeared to offer little over the comparator test (e.g., about 1 additional patient would be likely to experience metastases without an increase in false-positives).^{79,88}. In Ross et al (2014), the net benefit for CAPRA-S score exceeded that of the GC, with the net benefit of the GC plus CAPRA-S score being slightly better than the CAPRA-S score alone.^{88,} Finally, among men undergoing RT, decision curves suggested that the test would identify 3 or 4 men developing metastases per 100 tested at a fixed false-positive rate. Lobo et al (2015)^{95,} reported an individualized decision analysis comparing the GC with "usual care" using data from the cohorts in Karnes et al (2013) and Den et al (2014). The usual care probabilities of receiving each treatment were derived from the published literature. A 6% threshold for the GC score was used for GC-based treatment. Using the cohort from Karnes et al (2013), the estimated 10-year probability of metastasis or death was 0.32 (95% CI, 0.32 to 0.33) for usual care compared with 0.31 (95% CI, 0.30 to 0.32) for GC-based treatment. In the cohort from Den et al (2014), the estimated 10-year

probability of metastasis or death was 0.28 (95% CI, 0.27 to 0.29) for usual care compared with 0.26 (95% CI, 0.25 to 0.27) for GC-based treatment.

Study	Outcome	Range of Net Benefit versus	
		Treat None	Best Comparator
Spratt et al (2018) ^{92,}	Metastasis	-0.003 to 0.002	NR
Karnes et al (2018) ^{93,}	PC mortality	0.06 to 0.09	0.045 to 0.095
Ross et al (2016) ^{94,}	Metastasis	0.045 to 0.075	0.09 to 0.12
Freedland (2016) ^{89,}	Metastasis	0.01 to 0.045	0 to 0.02
Lobo et al (2015) ^{95,} with Karnes et al	Metastasis or death	NR	0.017
(2013) ^{86,} cohort			
Cooperberg et al (2015) ^{84,}	PCa mortality	0.003ª	NR
Klein et al (2015) ^{79,}	Metastasis	0.008 to 0.025	0.000 to 0.012
Den et al (2015) ^{82,}	Metastasis post-RT	0.02 to 0.03	-0.01 to 0.001
Lobo et al(2015) ^{95,} with Den et	Metastasis or death	NR	0.015
al(2014) ^{83,} cohort			
Ross et al (2014) ^{85,}	Metastasis	0.09 to 0.13	0.036 to 0.040
Karnes et al (2013) ^{86,}	Metastasis	0.009 to 0.020	-0.004 to 0.003
ND: not reported: DCa: prostate capes	DT. radiatharany		

NR: not reported; PCa: prostate cancer; RT: radiotherapy.

° For 25% threshold.

Changes in Management

Several studies have compared physician's treatment recommendations before and after receiving GC results.^{59,96,97,98,99,100,} Because the studies did not include information on outcomes and clinical validity has not been established, it is not known whether these treatment decisions represent a clinical improvement in management.

The Association Between the Genomic Classifier and Treatment Effects

Ross et al (2016) reported on results of a retrospective, comparative study of RT after RP for 422 men with pT3 disease or positive margins.^{94,} The men were from 4 cohorts previously described (Karnes et al [2013]^{86,}; Den et al [2014]^{83,}; Ross et al [2016]^{94,}; Freedland et al [2016]^{89,}). The 4 treatment groups were adjuvant RT (n=111), minimal residual disease salvage RT (n=70), salvage RT (n=83), and no RT (n=157). The primary endpoint was a metastasis. Thirty-seven men developed metastasis, and the median follow-up was 8 years. Both CAPRA-S (HR=1.39; 95% CI, 1.18 to 1.62) and Decipher (HR=1.28; 95% CI, 1.08 to 1.52) were independently associated with metastasis in multivariable analysis. There was no evidence that the treatment effect was dependent on genomic risk (interaction p=.16 for CAPRA-S, p=.39 for Decipher). Men with low CAPRA-S or low Decipher scores had a low-risk of metastatic events regardless of treatment selection, and men with high CAPRA-S or Decipher scores benefitted from adjuvant RT compared with the other treatments.

Section Summary: Decipher RP Prostate Cancer Classifier

Clinical validity has been evaluated in overlapping validation samples (including the development test set). The validation studies consisted of observational data obtained from registries or medical records with archived samples. Although each study evaluated different outcomes (i.e., metastasis, prostate cancer-specific mortality, BCR) in samples with different populations, all studies reported some incremental improvement in discrimination. Cls of AUC frequently overlapped between Decipher and comparators. Only 1 study (Karnes et al [2018]⁹³) reported 10-year disease-specific survival. Estimates with Cls of outcomes, particularly disease-specific mortality at 10 years, by GC low-, intermediate-, and high-risk are needed as well as reclassification analyses of prostate cancer-specific survival compared with comparators. Results did not consistently demonstrate meaningful improvement in reclassification-- possibly most importantly to lower risk categories. It is not clear whether the group of patients identified as low-risk using Decipher could be managed with an observation instead of adjuvant or early salvage RT.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 36 of 54

Management Decision in Castration-Resistant Prostate Cancer Clinical Context and Test Purpose

In men with metastatic castration-resistant prostate cancer (mCRPC), the purpose of protein biomarker assessment of circulating tumor cells (CTCs) is to inform a decision whether to administer androgen receptor signaling (ARS) inhibitors (e.g., abiraterone, enzalutamide), or a taxane (e.g., docetaxel).

Multiple approved therapeutic options exist for the treatment of men with mCRPC, which are given in conjunction with continued ADT. In particular, ARS inhibitors and taxane-based chemotherapy have both demonstrated effectiveness in prolonging survival but head-to-head comparisons of ARS inhibitors and taxanes in RCTs are lacking. Optimal sequencing of available treatments has also not been established. Guidelines have suggested that both ARS inhibitors and chemotherapy are appropriate for men with mCRPC who have sufficiently good performance status to tolerate chemotherapy as first-line treatment of mCRPC. In practice, sequencing depends on several factors such as sites and extent of disease, rates of progression, ease and convenience of administration, side effects, comorbidities, and patient preferences. However, unless a man has rapidly progressive, symptomatic disease, ARS inhibitors are generally used as first-line treatment of mCRPC because they are orally administered and have lower toxicity. After disease progression on first-line ARS inhibitor, men could then receive another ARS inhibitor or another systemic therapy, usually a taxane. A test that could inform the choice of second-line therapy would fill an unmet management need. The androgen receptor isoform encoded by splice variant 7 lacks the ligand-binding domain that is the target of the ARS inhibitors enzalutamide and abiraterone. Therefore, detection of androgen receptor splice variant 7 messenger RNA (AR-V7) in CTCs from men with mCRPC might be associated with a lack of response to enzalutamide and abiraterone but not with lack of response to taxanes. The question addressed in this section of the evidence review is: Does GEP testing improve the net health outcome in men with mCRPC compared with standard clinical care without AR-V7 testing? The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is men with mCRPC who have progressed on an ARS inhibitor (e.g., enzalutamide, abiraterone), have a good performance status (i.e., are able to tolerate chemotherapy), and who are deciding between a second ARS inhibitor or a taxane.

Interventions

The test being considered is the Oncotype DX AR-V7 Nuclear Detect. Detection of AR-V7 in men with progressive mCRPC is associated with resistance to the ARS inhibitors abiraterone and enzalutamide.^{101,} The Oncotype DX AR-V7 Nuclear Detect test is a liquid biopsy test that detects CTCs with nuclear expression of the AR-V7 truncated protein. The test reports a score of AR-V7-positive or -negative. Scher et al (2016) described the development of the test and results in the development cohort in which they observed longer OS for men taking taxanes compared with ARS inhibitors when AR-V7-positive CTCs were detected before therapy (HR=0.24; 95% CI, 0.10 to 0.57).^{102,} Scher et al (2017) explored whether expanding the AR-V7 scoring criteria to include both nuclear and cytoplasmic AR-V7 localization improved prediction in the same development cohort and concluded that the expanded "nuclear-agnostic" AR-V7 scoring criterion was less prognostic for men on ARS inhibitor therapy.^{103,}

Decisions about management of localized prostate cancer are typically made by patients, urologists, and oncologists in the secondary or tertiary care setting.

Comparators

Since there are no head-to-head comparisons of ARS inhibitors and taxanes in RCTs to determine optimal second- and subsequent-line therapies, in standard clinical care, physicians and men with mCRPC are making treatment decisions based on patient preference, disease characteristics, and comorbidities.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 37 of 54

Outcomes

Beneficial outcomes resulting from a true test result are prolonged survival, improved QOL, and reduction in unnecessary treatment-related adverse events. Harmful outcomes resulting from a false test result are unnecessary treatments and shortened survival. The primary survival outcome of interest is OS.

In a systematic review of randomized phase 3 trials of systemic therapies for CRPC, which included 23 trials (total N=13,909 men), the median OS was 19 months.^{96,} Outcomes with at least 1 year of follow-up of those surviving would be preferred.

Oncotype DX AR-V7 Nuclear Detect

Oncotype DX AR-V7 Nuclear Detectis used to detect nuclear-localized AR-V7 protein in CTCs of men with mCRPC who have failed first-line therapy and are considering additional ARS inhibitor therapy.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Two studies were not included in this assessment of clinical validity because they reported results in the developmental cohort.^{102,97,} Two published clinical validity studies met selection criteria.^{103,} Characteristics of the studies are provided in Table 31. Scher et al (2018) reported results of a blinded validation study including 142 samples from patients with histologically confirmed, progressing mCRPC from 3 centers in the U. S. and the United Kingdom from 2012 to 2016. The samples were collected prior to the administration of second-line or greater ARS inhibitors or taxanes. Armstrong et al (2019) reported results of the PROPHECY trial, a prospective validation study of AR-V7 detection in men with high-risk mCRPC starting abiraterone or enzalutamide treatment.

Study	Study Population	Design	Outcome Measure	Threshold for Positive Index Test	Blinding of Assessors
Scher et al (2018) ^{103,}	Men with progressing mCRPC undergoing change in therapy	Retrospective; unclear whether samples were consecutive or randomly chosen from eligible	OS (68 men with 12- mo follow-up, 15 men with 24 m follow-up, 6 men with 36-mo follow-up)	At least 1 CTC with an intact nucleus and nuclear-localized AR-V7 signal-to-noise ratio above a prespecified background intensity	Yes
Armstrong et al (2019) ^{98,}	Men with progressive, high-risk mCRPC initiating standard-of- care treatment with enzalutamide or abiraterone. Prior exposure to enzalutamide or abiraterone was permitted for men who were planning to receive the	Prospective, consecutive	PFS (primary)Response rates (PSA and radiographic) OS (secondary)	Johns Hopkins and Epic AR-V7 assays; results for both assays reported	Yes

Table 31. Characteristics of Clinical Validity Studies Assessing Oncotype DX AR-V7

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 38 of 54

Study	Study Population	Design	Outcome Measure	Threshold for Positive Index Test	Blinding of Assessors
	alternative				
	agent				

CTC: circulating tumor cell; mCRPC: metastatic castration-resistant prostate cancer; OS: overall survival; PFS: progression-free survival; PSA: prostate-specific antigen.

Results of the validation studies are shown in Table 32. In Scher et al (2018), median follow-up time in surviving men was not provided. Sixty-eight men were still in the risk set at 12 months. Numerically, men treated with ARS inhibitors had the longest OS if they were AR-V7-negative and had the shortest OS if they were AR-V7-positive. The unadjusted HR for OS for ARS inhibitors versus taxanes was statistically significantly greater than one (favoring ARS inhibitors) in the AR-V7-negative men, while there was no statistically significant difference in OS (but with an unadjusted HR favoring taxanes) in AR-V7-positive men. A test of interaction for AR-V7 status by treatment was not provided. The analysis was further stratified by a binary prognostic risk score (high vs. low) developed from the training cohort and including clinical biomarkers (see Table 31). However, the additional stratification resulted in the group that was AR-V7-positive and receiving ARS inhibitors including fewer than 10 men for both high- and low-risk. In Armstrong et al (2019), detection of AR-V7 in CTCs was associated with shorter PFS and OS.

Study	Initial N	Final N	Excluded Samples	AR-V7+, %	Median OS Therapy	(mo) by AR-	V7 and Ne	xt-Line
					AR-V7+, ARS Inhibitor	AR-V7+, Taxane	AR-V7-, ARS Inhibitor	AR-V7-, Taxane
Scher et al (2018) ^{103,}	248	142 (70 before ARS inhibitor tx, 72 before taxane tx)	144 (93 obtained before first-line tx, 24 duplicates, 23 second-line tx other than ARS inhibitor or taxane, 2 insufficient material, 2 missing clinical data)	24	7.3	14.3	19.8	12.8
HR (95% Cl); p ARS vs. taxane					AR-V7+0.6 (0.3 to 1.4);.25	AR-V7-1.7 (1.0 to 2.8);.05		
Interaction p					Not reported			
Armstrong et al (2019) ^{98,}	118	107	2 unevaluable (1%)	10	ARS inhibitor: 8.4 Taxane: NR	ARS inhibitor: 25.5 Taxane:NR		
HR (95% Cl); p ARS vs. taxane					Not reported	Not reported		
Interaction p			Changlishanani		Not reported			

Table 32. Results of Clinical Validity Studies Assessing Oncotype DX AR-V7

ARS: androgen receptor signaling; CI: confidence interval; HR: hazard ratio; NR: not reported; OS: overall survival; tx: treatment.

Table 33. Cross-Tabulation of AR-V7 Status and Clinical Risk Score

		Risk Score	Risk Score				
		High	Low	Total			
AR-V7 status	Positive	24	10	34			
	Negative	46	62	108			

Total	70	72	142	
Adapted from Scher et al (2018). ^{103,}				

Tables 34 and 35 display notable limitations identified in each study.

Table 34. Study Relevance Limitations of Clinical Validity Studies Assessing Oncotype DX AR-V7

Study	Populationa	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Scher et al					1. Median follow-up in surviving
(2018) ^{103,}					men not clear but overall <50%
					of men had 12-mo follow-up
A museline as a ball					

Armstrong et al (2019)^{98,}

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest. ^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

Table 35. Study Design and Conduct Limitations of Clinical Validity Studies Assessing Oncotype DX AR-V7

Study	Selection ^a	Blinding ^b	Delivery of Test ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
Scher et al (2018) ^{103,}	2. Unclear if original 248 samples included were consecutive or randomly chosen from eligible					1. Interaction p value not provided
Armstrong et al (2019) ^{98,}	2. Unclear if consecutive or convenience sample					

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

^b Blinding key: 1. Not blinded to results of reference or other comparator tests.

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 40 of 54

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No studies reporting direct evidence were identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of the Oncotype DX AR-V7 test has not been established, a chain of evidence supporting the test's clinical utility cannot be constructed.

Section Summary: Oncotype DX AR-V7 Nuclear Detect

Multiple, high-quality studies of the marketed version of the test (including current algorithms and cutoffs), in populations independent of the developmental cohort, that include the intended use population and have consistent and precise results are needed to characterize the performance characteristics.

One retrospective analysis of 142 men from the U. S. and the United Kingdom including men with progressing mCRPC undergoing a change in therapy is available. The median follow-up in surviving men is unclear, but, overall, 68 men had 12 months of follow-up, 15 men had 24 months of follow-up, and 6 men had 36 months of follow-up. Men treated with ARS inhibitors had the longest OS if they were AR-V7-negative (median, 19.8 months) and had the shortest OS if they were AR-V7-positive (median, 7.3 months). The unadjusted HR for OS was statistically significantly longer for ARS inhibitors compared with taxanes in the AR-V7-negative men (HR=1.7; 95% Cl, 1.0 to 2.8) but not in AR-V7-positive men (0.6; 95% Cl, 0.3 to 1.4). However, a test of interaction for AR-V7 status by treatment was not provided. In a prospective validation study of AR-V7 detection in 118 men with high-risk mCRPC starting abiraterone or enzalutamide treatment, the detection of AR-V7 in CTCs was associated with shorter PFS and OS.

Summary of Evidence

Initial Management Decision: Active Surveillance versus Therapeutic Intervention

For individuals who have clinically localized untreated prostate cancer who receive Prolaris, the evidence includes retrospective cohort studies of clinical validity using archived samples in patients of mixed risk categories. Relevant outcomes include overall survival (OS), disease-specific survival, quality of life (QOL), and treatment-related morbidity. For the low-risk group, the Prostate Testing for Cancer and Treatment trial showed 99% 10-year disease-specific survival in mostly low-risk patients receiving active surveillance. The low mortality rate estimated with tight precision makes it unlikely that a test intended to identify a subgroup of low-risk men with a net benefit from immediate treatment instead of active surveillance would find such a group. For the intermediate-risk group, the evidence of improved clinical validity or prognostic accuracy for prostate cancer death using Prolaris Cell Cycle Progression score in patients managed conservatively after a needle biopsy has shown some improvement in areas under the receiver operating characteristic curve over clinicopathologic risk stratification tools. There is limited indirect evidence for potential clinical utility. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have clinically localized untreated prostate cancer who receive Oncotype DX Prostate, the evidence includes case-cohort and retrospective cohort studies of clinical validity using archived samples in patients of mixed risk categories, and a decision-curve analysis examining indirect evidence of clinical utility. Relevant outcomes include OS, disease-specific survival, QOL, and treatment-related morbidity. Evidence for clinical validity and potential clinical utility of Oncotype DX Prostate in patients with clinically localized prostate cancer derives from a study predicting adverse

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 41 of 54

pathology after RP. The validity of using tumor pathology as a surrogate for the risk of progression and cancer-specific death is unclear. It is also unclear whether results from an RP population can be generalized to an active surveillance population. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have clinically localized untreated prostate cancer who receive Decipher Biopsy, the evidence includes retrospective cohort studies of clinical validity using archived samples in intermediate- and high-risk patients and no studies of clinical utility. Relevant outcomes include OS, disease-specific survival, QOL, and treatment-related morbidity. A test designed to identify intermediate-risk men who can receive active surveillance instead of RP or radiotherapy (RT) or high-risk men who can forego androgen deprivation therapy would need to show very high negative predictive value for disease-specific mortality at 10 years and improvement in prediction compared with existing tools used to select such men. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have clinically localized untreated prostate cancer who receive the ProMark protein biomarker test, the evidence includes a retrospective cohort study of clinical validity using archived samples and no studies of clinical utility. Relevant outcomes include OS, disease-specific survival, QOL, and treatment-related morbidity. Current evidence does not support improved outcomes with ProMark given that only a single clinical validity study is available. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Management Decision After Radical Prostatectomy

For individuals who have localized prostate cancer treated with RP who receive Prolaris, the evidence includes retrospective cohort studies of clinical validity using archived samples. Relevant outcomes include OS, disease-specific survival, QOL, and treatment-related morbidity. No direct evidence is available to support the clinical utility of Prolaris for improving net outcomes of patients with localized prostate cancer following RP. The chain of evidence is also incomplete. Decision-curve analysis did not provide convincing evidence of meaningful improvement in net benefit by incorporating the cell cycle progression (CCP) score. Evidence of improved clinical validity or prognostic accuracy for prostate cancer death using the Prolaris Cell Cycle Progression score in patients after prostatectomy has shown some improvement in areas under the receiver operating characteristic curve over clinicopathologic risk stratification tools. Although Prolaris CCP score may have an association with biochemical recurrence (BCR), disease-specific survival outcomes were reported in only 1 analysis. A larger number of disease-specific survival events and precision estimates for discrimination measures are needed. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have localized prostate cancer who are treated with RP and who receive the Decipher RP prostate cancer classifier, the evidence includes a study of analytic validity, prospective and retrospective studies of clinical validity using overlapping archived samples, decision-curve analyses examining indirect evidence of clinical utility, and prospective decision-impact studies without pathology or clinical outcomes. Relevant outcomes include OS, disease-specific survival, QOL, and treatment-related morbidity. The clinical validity of the Decipher RP genomic classifier has been evaluated in samples of patients with high-risk prostate cancer undergoing different interventions following RP. Studies reported some incremental improvement in discrimination. However, it is unclear whether there is consistently improved reclassification-particularly to higher risk categories-or whether the test could be used to predict which men will benefit from radiotherapy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Management Decision in Castration-Resistant Prostate Cancer

For individuals who have metastatic castration-resistant prostate cancer who receive the Oncotype DX AR-V7 Nuclear Detect, the evidence includes 1 prospective cohort study, 1 retrospective cohort

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 42 of 54

study of clinical validity using archived samples, and no studies of clinical utility. Relevant outcomes include OS, disease-specific survival, QOL, and treatment-related morbidity. Current evidence does not support improved outcomes with Oncotype DX AR-V7 Nuclear Detect, given that only 2 clinical validity studies meeting inclusion criteria were available. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Society of Clinical Oncology

In 2020, the American Society of Clinical Oncology (ASCO) published a guideline on molecular biomarkers in localized prostate cancer.^{104,} The guidelines state, "Currently, there are no strong data or expert guidelines to support active surveillance in otherwise healthy men with Grade Group 3 or higher cancer; therefore, we would consider the use of genomic biomarkers only in situations in which the assay result, when considered as a whole with routine clinical factors, is likely to affect a physician's recommendation or a patient's choice for surveillance versus treatment, but they should not be used routinely."

Specific recommendations included the following:

Molecular biomarkers to identify patients with prostate cancer who are most likely to benefit from active surveillance:

- Recommendation 1.1. Commercially available molecular biomarkers (i.e. Oncotype Dx Prostate, Prolaris, Decipher, and ProMark) may be offered in situations in which the assay result, when considered as a whole with routine clinical factors, is likely to affect management. Routine ordering of molecular biomarkers is not recommended (Type: Evidence based; Evidence quality: Intermediate; Strength of recommendation: Moderate).
- Recommendation 1.2. Any additional molecular biomarkers evaluated do not have sufficient data to be clinically actionable or are not commercially available and thus should not be offered (Type: Evidence based; Evidence quality: Insufficient; Strength of recommendation: Moderate).

Molecular biomarkers to diagnose clinically significant prostate cancer:

- Recommendation 2.1. Commercially available molecular biomarkers (i.e. Oncotype Dx Prostate, Prolaris, Decipher, and ProMark) may be offered in situations in which the assay result, when considered as a whole with routine clinical factors, is likely to affect management. Routine ordering of molecular biomarkers is not recommended (Type: Evidence based; Evidence quality: Intermediate; Recommendation: Moderate).
- Recommendation 2.2. Any additional molecular biomarkers evaluated do not have sufficient data to be clinically actionable or are not commercially available and thus should not be offered (Type: Evidence based; Evidence quality: Insufficient; Strength of recommendation: Moderate).

Molecular biomarkers to guide the decision of post prostatectomy adjuvant versus salvage radiation: Recommendation 3.1. The Expert Panel recommends consideration of a commercially available molecular biomarker (e.g., Decipher Genomic Classifier) in situations in which the assay result, when considered as a whole with routine clinical factors, is likely to affect management. In the absence of

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 43 of 54

prospective clinical trial data, routine use of genomic biomarkers in the postprostatectomy setting to determine adjuvant versus salvage radiation or to initiate systemic therapies should not be offered (Type: Evidence based; Evidence quality: Intermediate; Strength of recommendation: Moderate). Recommendation 3.2. Any additional molecular biomarkers evaluated do not have sufficient data to be clinically actionable or are not commercially available and thus should not be offered (Type: Evidence based; Evidence quality: Insufficient; Strength of recommendation: Moderate).

American Urological Association and American Society for Radiation Oncology

The American Urological Association and American Society for Radiation Oncology published guidelines on clinically localized prostate cancer.^{105,} The guidelines included the following statements on risk assessment:

- "Clinicians should use clinical T stage, serum PSA, Grade Group (Gleason score), and tumor volume on biopsy to risk stratify patients with newly diagnosed prostate cancer. (Strong Recommendation; Evidence Level: Grade B)"
- 2. "Clinicians may selectively use tissue-based genomic biomarkers when added risk stratification may alter clinical decision-making. (Expert Opinion)"
- 3. "Clinicians should not routinely use tissue-based genomic biomarkers for risk stratification or clinical decision-making. (Moderate Recommendation; Evidence Level: Grade B)"

The American Urological Association (2018) published guidelines for castration-resistant prostate cancer.^{106,} The guidelines do not mention AR-V7 assays.

National Comprehensive Cancer Network

The National Comprehensive Cancer Network guidelines for prostate cancer (v.1.2023) provide a table of tissue-based tests for prostate cancer prognosis.^{12,}

The guidelines include the following statements related to risk stratification:

- Patients with NCCN low, favorable intermediate, unfavorable intermediate, or high-risk disease and life expectancy ≥10 y may consider the use of the following tumor-based molecular assays: Decipher, Oncotype DX Prostate, and Prolaris.
- Decipher may be considered to inform adjuvant treatment if adverse features are found after radical prostatectomy and during workup for radical prostatectomy PSA persistence or recurrence (category 2B for the latter setting)

The panel also recommended that "the use of AR-V7 tests in circulating tumor cells can be considered to help guide selection of therapy in the post-abiraterone/enzalutamide metastatic castration-resistant prostate cancer setting."

National Institute for Health and Care Excellence

In 2019 (updated 2021), the National Institute for Health and Care Excellence updated its guidance on the diagnosis and management of prostate cancer.^{107,} The guidance did not address gene expression profile testing.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 36.

Table 36. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
Prolaris			
NCT03152448°	Two-Part Prospective Study to Measure Impact of Prolaris® Testing Added to Treatment Decision Following Biopsy in Newly Diagnosed Prostate Cancer Patients to Measure Prediction of Progression/Recurrence in Men Treated at VAMC	1511	Mar 2022
NCT03290508°	Long-Term Prospective Registry to Evaluate Treatment Decisions and Clinical Outcomes in Patients With Favorable Intermediate- Risk Localized Prostate Cancer Following Cell Cycle Progression (CCP) Testing (Prolaris® Test)	524	Jan 2022
NCT04404894ª	Long-Term Prospective Registry to Evaluate Treatment Decisions and Clinical Outcomes in Prostate Cancer Patients From Diverse Urology Practice Settings Following Prolaris® Testing	500	Nov 2029
Decipher			
NCT02723734	Validation Study on the Impact of Decipher Testing - VANDAAM Study	250	May 2024
NCT04396808	Genomics in Michigan to AdJust Outcomes in Prostate canceR (G- MAJOR): A Randomized Multi-center Study for Men With Newly Diagnosed Favorable Risk Prostate Cancer	900	Nov 2023
NCT05050084°	Parallel Phase III Randomized Trials of Genomic-Risk Stratified Unfavorable Intermediate Risk Prostate Cancer: De-Intensification and Intensification Clinical Trial Evaluation (GUIDANCE)	2050	Apr 2037
NCT04484818	A Phase III Double Blinded Study of Early Intervention After RADICAI ProstaTEctomy With Androgen Deprivation Therapy With or Without Darolutamide vs. Placebo in Men at Highest Risk of Prostate Cancer Metastasis by Genomic Stratification (ERADICATE)	810	May 2028
NCT04513717	Parallel Phase III Randomized Trials for High Risk Prostate Cancer Evaluating De-Intensification for Lower Genomic Risk and Intensification of Concurrent Therapy for Higher Genomic Risk With Radiation (PREDICT-RT*)	2478	Dec 2033
Prolaris or Decipher or Oncotype			
NCT04396808	Genomics in Michigan to AdJust Outcomes in Prostate canceR (G- MAJOR): A Randomized Multi-center Study for Men With Newly Diagnosed Favorable Risk Prostate Cancer	900	Nov 2023
NCT: national clir	nical trial		

^a Denotes industry-sponsored or cosponsored trial.

References

- 1. Dall'Era MA, Cooperberg MR, Chan JM, et al. Active surveillance for early-stage prostate cancer: review of the current literature. Cancer. Apr 15 2008; 112(8): 1650-9. PMID 18306379
- 2. Bangma CH, Roemeling S, Schroder FH. Overdiagnosis and overtreatment of early detected prostate cancer. World J Urol. Mar 2007; 25(1): 3-9. PMID 17364211
- 3. Johansson JE, Andren O, Andersson SO, et al. Natural history of early, localized prostate cancer. JAMA. Jun 09 2004; 291(22): 2713-9. PMID 15187052
- 4. Ploussard G, Epstein JI, Montironi R, et al. The contemporary concept of significant versus insignificant prostate cancer. Eur Urol. Aug 2011; 60(2): 291-303. PMID 21601982
- 5. Harnden P, Naylor B, Shelley MD, et al. The clinical management of patients with a small volume of prostatic cancer on biopsy: what are the risks of progression? A systematic review and meta-analysis. Cancer. Mar 01 2008; 112(5): 971-81. PMID 18186496
- 6. Brimo F, Montironi R, Egevad L, et al. Contemporary grading for prostate cancer: implications for patient care. Eur Urol. May 2013; 63(5): 892-901. PMID 23092544

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 45 of 54

- 7. Eylert MF, Persad R. Management of prostate cancer. Br J Hosp Med (Lond). Feb 2012; 73(2): 95-9. PMID 22504752
- 8. Eastham JA, Kattan MW, Fearn P, et al. Local progression among men with conservatively treated localized prostate cancer: results from the Transatlantic Prostate Group. Eur Urol. Feb 2008; 53(2): 347-54. PMID 17544572
- 9. Bill-Axelson A, Holmberg L, Ruutu M, et al. Radical prostatectomy versus watchful waiting in early prostate cancer. N Engl J Med. May 12 2005; 352(19): 1977-84. PMID 15888698
- 10. Thompson IM, Goodman PJ, Tangen CM, et al. Long-term survival of participants in the prostate cancer prevention trial. N Engl J Med. Aug 15 2013; 369(7): 603-10. PMID 23944298
- 11. Albertsen PC, Hanley JA, Fine J. 20-year outcomes following conservative management of clinically localized prostate cancer. JAMA. May 04 2005; 293(17): 2095-101. PMID 15870412
- National Comprehensive Cancer Network (NCCN). NCCN Clincal Practice Guidelines in Oncology: Prostate Cancer. Version 1.2023. https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf. Accessed November 2, 2022.
- American Urological Association (AUA). Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. 2017; http://www.auanet.org/guidelines/clinically-localizedprostate-cancer-new-(aua/astro/suo-guideline-2017). Accessed December 3, 2021.
- 14. Thompson IM, Valicenti RK, Albertsen P, et al. Adjuvant and salvage radiotherapy after prostatectomy: AUA/ASTRO Guideline. J Urol. Aug 2013; 190(2): 441-9. PMID 23707439
- Food and Drug Administration (FDA). The Public Health Evidence for FDA Oversight of Laboratory Developed Tests: 20 Case Studies. 2015; http://www.nilausa.org/images/nila/The%20Public%20Health%20Case%20for%20FDA%20Oversight%20o f%20LDTs%20110915(2)_508ed%20(1).pdf. Accessed November 2, 2022.
- Borley N, Feneley MR. Prostate cancer: diagnosis and staging. Asian J Androl. Jan 2009; 11(1): 74-80. PMID 19050692
- 17. Freedland SJ. Screening, risk assessment, and the approach to therapy in patients with prostate cancer. Cancer. Mar 15 2011; 117(6): 1123-35. PMID 20960523
- 18. Albertsen PC. Treatment of localized prostate cancer: when is active surveillance appropriate?. Nat Rev Clin Oncol. Jul 2010; 7(7): 394-400. PMID 20440282
- Ip S, Dahabreh IJ, Chung M, et al. An evidence review of active surveillance in men with localized prostate cancer. Evid Rep Technol Assess (Full Rep). Dec 2011; (204): 1-341. PMID 23126653
- Nam RK, Cheung P, Herschorn S, et al. Incidence of complications other than urinary incontinence or erectile dysfunction after radical prostatectomy or radiotherapy for prostate cancer: a population-based cohort study. Lancet Oncol. Feb 2014; 15(2): 223-31. PMID 24440474
- Hamdy FC, Donovan JL, Lane JA, et al. 10-Year Outcomes after Monitoring, Surgery, or Radiotherapy for Localized Prostate Cancer. N Engl J Med. Oct 13 2016; 375(15): 1415-1424. PMID 27626136
- 22. Tosoian JJ, Mamawala M, Epstein JI, et al. Intermediate and Longer-Term Outcomes From a Prospective Active-Surveillance Program for Favorable-Risk Prostate Cancer. J Clin Oncol. Oct 20 2015; 33(30): 3379-85. PMID 26324359
- 23. Klotz L, Vesprini D, Sethukavalan P, et al. Long-term follow-up of a large active surveillance cohort of patients with prostate cancer. J Clin Oncol. Jan 20 2015; 33(3): 272-7. PMID 25512465
- 24. Wilt TJ, Brawer MK, Jones KM, et al. Radical prostatectomy versus observation for localized prostate cancer. N Engl J Med. Jul 19 2012; 367(3): 203-13. PMID 22808955
- 25. Wilt TJ, Jones KM, Barry MJ, et al. Follow-up of Prostatectomy versus Observation for Early Prostate Cancer. N Engl J Med. Jul 13 2017; 377(2): 132-142. PMID 28700844
- 26. van den Bergh RC, Korfage IJ, Roobol MJ, et al. Sexual function with localized prostate cancer: active surveillance vs radical therapy. BJU Int. Oct 2012; 110(7): 1032-9. PMID 22260273
- 27. Johansson E, Steineck G, Holmberg L, et al. Long-term quality-of-life outcomes after radical prostatectomy or watchful waiting: the Scandinavian Prostate Cancer Group-4 randomised trial. Lancet Oncol. Sep 2011; 12(9): 891-9. PMID 21821474

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 46 of 54

- Wu CL, Schroeder BE, Ma XJ, et al. Development and validation of a 32-gene prognostic index for prostate cancer progression. Proc Natl Acad Sci U S A. Apr 09 2013; 110(15): 6121-6. PMID 23533275
- 29. Spans L, Clinckemalie L, Helsen C, et al. The genomic landscape of prostate cancer. Int J Mol Sci. May 24 2013; 14(6): 10822-51. PMID 23708091
- Schoenborn JR, Nelson P, Fang M. Genomic profiling defines subtypes of prostate cancer with the potential for therapeutic stratification. Clin Cancer Res. Aug 01 2013; 19(15): 4058-66. PMID 23704282
- Huang J, Wang JK, Sun Y. Molecular pathology of prostate cancer revealed by nextgeneration sequencing: opportunities for genome-based personalized therapy. Curr Opin Urol. May 2013; 23(3): 189-93. PMID 23385974
- 32. Yu YP, Song C, Tseng G, et al. Genome abnormalities precede prostate cancer and predict clinical relapse. Am J Pathol. Jun 2012; 180(6): 2240-8. PMID 22569189
- 33. Agell L, Hernandez S, Nonell L, et al. A 12-gene expression signature is associated with aggressive histological in prostate cancer: SEC14L1 and TCEB1 genes are potential markers of progression. Am J Pathol. Nov 2012; 181(5): 1585-94. PMID 23083832
- 34. Thompson I, Thrasher JB, Aus G, et al. Guideline for the management of clinically localized prostate cancer: 2007 update. J Urol. Jun 2007; 177(6): 2106-31. PMID 17509297
- 35. Kattan MW, Eastham JA, Wheeler TM, et al. Counseling men with prostate cancer: a nomogram for predicting the presence of small, moderately differentiated, confined tumors. J Urol. Nov 2003; 170(5): 1792-7. PMID 14532778
- 36. Cooperberg MR, Freedland SJ, Pasta DJ, et al. Multiinstitutional validation of the UCSF cancer of the prostate risk assessment for prediction of recurrence after radical prostatectomy. Cancer. Nov 15 2006; 107(10): 2384-91. PMID 17039503
- Chen RC, Chang P, Vetter RJ, et al. Recommended patient-reported core set of symptoms to measure in prostate cancer treatment trials. J Natl Cancer Inst. Jul 2014; 106(7). PMID 25006192
- 38. Cuzick J, Berney DM, Fisher G, et al. Prognostic value of a cell cycle progression signature for prostate cancer death in a conservatively managed needle biopsy cohort. Br J Cancer. Mar 13 2012; 106(6): 1095-9. PMID 22361632
- 39. Cuzick J, Stone S, Fisher G, et al. Validation of an RNA cell cycle progression score for predicting death from prostate cancer in a conservatively managed needle biopsy cohort. Br J Cancer. Jul 28 2015; 113(3): 382-9. PMID 26103570
- 40. Lin DW, Crawford ED, Keane T, et al. Identification of men with low-risk biopsy-confirmed prostate cancer as candidates for active surveillance. Urol Oncol. Jun 2018; 36(6): 310.e7-310.e13. PMID 29655620
- Montironi R, Mazzuccheli R, Scarpelli M, et al. Gleason grading of prostate cancer in needle biopsies or radical prostatectomy specimens: contemporary approach, current clinical significance and sources of pathology discrepancies. BJU Int. Jun 2005; 95(8): 1146-52. PMID 15877724
- 42. Tward JD, Schlomm T, Bardot S, et al. Personalizing Localized Prostate Cancer: Validation of a Combined Clinical Cell-cycle Risk (CCR) Score Threshold for Prognosticating Benefit From Multimodality Therapy. Clin Genitourin Cancer. Aug 2021; 19(4): 296-304.e3. PMID 33608228
- Crawford ED, Scholz MC, Kar AJ, et al. Cell cycle progression score and treatment decisions in prostate cancer: results from an ongoing registry. Curr Med Res Opin. Jun 2014; 30(6): 1025-31. PMID 24576172
- Shore N, Concepcion R, Saltzstein D, et al. Clinical utility of a biopsy-based cell cycle gene expression assay in localized prostate cancer. Curr Med Res Opin. Apr 2014; 30(4): 547-53. PMID 24320750
- Shore ND, Kella N, Moran B, et al. Impact of the Cell Cycle Progression Test on Physician and Patient Treatment Selection for Localized Prostate Cancer. J Urol. Mar 2016; 195(3): 612-8.
 PMID 26403586

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 47 of 54

- Schaink A, Li C, Wells D, et al. Prolaris Cell Cycle Progression Test for Localized Prostate Cancer: A Health Technology Assessment. Ont Health Technol Assess Ser. 2017; 17(6): 1-75. PMID 28572867
- 47. Klein EA, Cooperberg MR, Magi-Galluzzi C, et al. A 17-gene assay to predict prostate cancer aggressiveness in the context of Gleason grade heterogeneity, tumor multifocality, and biopsy undersampling. Eur Urol. Sep 2014; 66(3): 550-60. PMID 24836057
- 48. Cullen J, Rosner IL, Brand TC, et al. A Biopsy-based 17-gene Genomic Prostate Score Predicts Recurrence After Radical Prostatectomy and Adverse Surgical Pathology in a Racially Diverse Population of Men with Clinically Low- and Intermediate-risk Prostate Cancer. Eur Urol. Jul 2015; 68(1): 123-31. PMID 25465337
- 49. Whalen MJ, Hackert V, Rothberg MB, et al. Prospective correlation between likelihood of favorable pathology on the 17-Gene Genomic Prostate Score and actual pathological outcomes at radical prostatectomy. Urol Pract. Sep 2016;3(5):379-386.
- 50. Van Den Eeden SK, Lu R, Zhang N, et al. A Biopsy-based 17-gene Genomic Prostate Score as a Predictor of Metastases and Prostate Cancer Death in Surgically Treated Men with Clinically Localized Disease. Eur Urol. Jan 2018; 73(1): 129-138. PMID 28988753
- 51. Salmasi A, Said J, Shindel AW, et al. A 17-Gene Genomic Prostate Score Assay Provides Independent Information on Adverse Pathology in the Setting of Combined Multiparametric Magnetic Resonance Imaging Fusion Targeted and Systematic Prostate Biopsy. J Urol. Sep 2018; 200(3): 564-572. PMID 29524506
- 52. Cooperberg MR, Simko JP, Cowan JE, et al. Validation of a cell-cycle progression gene panel to improve risk stratification in a contemporary prostatectomy cohort. J Clin Oncol. Apr 10 2013; 31(11): 1428-34. PMID 23460710
- 53. McShane LM, Altman DG, Sauerbrei W, et al. Reporting recommendations for tumor marker prognostic studies. J Clin Oncol. Dec 20 2005; 23(36): 9067-72. PMID 16172462
- 54. Epstein JI, Allsbrook WC, Amin MB, et al. The 2005 International Society of Urological Pathology (ISUP) Consensus Conference on Gleason Grading of Prostatic Carcinoma. Am J Surg Pathol. Sep 2005; 29(9): 1228-42. PMID 16096414
- 55. Brand TC, Zhang N, Crager MR, et al. Patient-specific Meta-analysis of 2 Clinical Validation Studies to Predict Pathologic Outcomes in Prostate Cancer Using the 17-Gene Genomic Prostate Score. Urology. Mar 2016; 89: 69-75. PMID 26723180
- Albala D, Kemeter MJ, Febbo PG, et al. Health Economic Impact and Prospective Clinical Utility of Oncotype DX(R) Genomic Prostate Score. Rev Urol. 2016; 18(3): 123-132. PMID 27833462
- 57. Eure G, Germany R, Given R, et al. Use of a 17-Gene Prognostic Assay in Contemporary Urologic Practice: Results of an Interim Analysis in an Observational Cohort. Urology. Sep 2017; 107: 67-75. PMID 28454985
- 58. Badani KK, Kemeter MJ, Febbo PG, et al. The impact of a biopsy based 17-Gene Genomic Prostate Score on treatment recommendations in men with newly diagnosed clinically prostate cancer who are candidates for active surveillance. Urol Pract. 2015;2(4):181-189. PMID not Indexed in Pubmed
- 59. Canfield SK, M.J.; Febbo, P.G.; Hornberger, J. Balancing confounding and generalizability using observational, real-world data: 17-gene genomic prostate score assay effect on active surveillance. Rev Urol. 2018;20(2):69-76.
- 60. Canfield S, Kemeter MJ, Hornberger J, et al. Active Surveillance Use Among a Low-risk Prostate Cancer Population in a Large US Payer System: 17-Gene Genomic Prostate Score Versus Other Risk Stratification Methods. Rev Urol. 2017; 19(4): 203-212. PMID 29472824
- 61. Vickers AJ, Elkin EB. Decision curve analysis: a novel method for evaluating prediction models. Med Decis Making. Nov-Dec 2006; 26(6): 565-74. PMID 17099194
- 62. Cooperberg MR, Broering JM, Carroll PR. Risk assessment for prostate cancer metastasis and mortality at the time of diagnosis. J Natl Cancer Inst. Jun 16 2009; 101(12): 878-87. PMID 19509351

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 48 of 54

- 63. Berlin A, Murgic J, Hosni A, et al. Genomic Classifier for Guiding Treatment of Intermediate-Risk Prostate Cancers to Dose-Escalated Image Guided Radiation Therapy Without Hormone Therapy. Int J Radiat Oncol Biol Phys. Jan 01 2019; 103(1): 84-91. PMID 30170099
- 64. Nguyen PL, Shin H, Yousefi K, et al. Impact of a Genomic Classifier of Metastatic Risk on Postprostatectomy Treatment Recommendations by Radiation Oncologists and Urologists. Urology. Jul 2015; 86(1): 35-40. PMID 26142578
- 65. Tosoian JJ, Birer SR, Jeffrey Karnes R, et al. Performance of clinicopathologic models in men with high risk localized prostate cancer: impact of a 22-gene genomic classifier. Prostate Cancer Prostatic Dis. Dec 2020; 23(4): 646-653. PMID 32231245
- 66. Blume-Jensen P, Berman DM, Rimm DL, et al. Development and clinical validation of an in situ biopsy-based multimarker assay for risk stratification in prostate cancer. Clin Cancer Res. Jun 01 2015; 21(11): 2591-600. PMID 25733599
- 67. Fossati N, Karnes RJ, Boorjian SA, et al. Long-term Impact of Adjuvant Versus Early Salvage Radiation Therapy in pT3N0 Prostate Cancer Patients Treated with Radical Prostatectomy: Results from a Multi-institutional Series. Eur Urol. Jun 2017; 71(6): 886-893. PMID 27484843
- 68. Hwang WL, Tendulkar RD, Niemierko A, et al. Comparison Between Adjuvant and Early-Salvage Postprostatectomy Radiotherapy for Prostate Cancer With Adverse Pathological Features. JAMA Oncol. May 10 2018; 4(5): e175230. PMID 29372236
- 69. Buscariollo DL, Drumm M, Niemierko A, et al. Long-term results of adjuvant versus early salvage postprostatectomy radiation: A large single-institutional experience. Pract Radiat Oncol. Mar 2017; 7(2): e125-e133. PMID 28274403
- Freedland SJ, Rumble RB, Finelli A, et al. Adjuvant and salvage radiotherapy after prostatectomy: American Society of Clinical Oncology clinical practice guideline endorsement. J Clin Oncol. Dec 01 2014; 32(34): 3892-8. PMID 25366677
- Stephenson AJ, Scardino PT, Kattan MW, et al. Predicting the outcome of salvage radiation therapy for recurrent prostate cancer after radical prostatectomy. J Clin Oncol. May 20 2007; 25(15): 2035-41. PMID 17513807
- 72. Stephenson AJ, Scardino PT, Eastham JA, et al. Postoperative nomogram predicting the 10year probability of prostate cancer recurrence after radical prostatectomy. J Clin Oncol. Oct 01 2005; 23(28): 7005-12. PMID 16192588
- Cooperberg MR, Hilton JF, Carroll PR. The CAPRA-S score: A straightforward tool for improved prediction of outcomes after radical prostatectomy. Cancer. Nov 15 2011; 117(22): 5039-46. PMID 21647869
- 74. Cuzick J, Swanson GP, Fisher G, et al. Prognostic value of an RNA expression signature derived from cell cycle proliferation genes in patients with prostate cancer: a retrospective study. Lancet Oncol. Mar 2011; 12(3): 245-55. PMID 21310658
- 75. Bishoff JT, Freedland SJ, Gerber L, et al. Prognostic utility of the cell cycle progression score generated from biopsy in men treated with prostatectomy. J Urol. Aug 2014; 192(2): 409-14. PMID 24508632
- 76. Swanson GP, Lenz L, Stone S, et al. Cell-cycle risk score more accurately determines the risk for metastases and death in prostatectomy patients compared with clinical features alone. Prostate. Mar 2021; 81(4): 261-267. PMID 33475174
- 77. Koch MO, Cho JS, Kaimakliotis HZ, et al. Use of the cell cycle progression (CCP) score for predicting systemic disease and response to radiation of biochemical recurrence. Cancer Biomark. Jun 07 2016; 17(1): 83-8. PMID 27314296
- 78. Freedland SJ, Gerber L, Reid J, et al. Prognostic utility of cell cycle progression score in men with prostate cancer after primary external beam radiation therapy. Int J Radiat Oncol Biol Phys. Aug 01 2013; 86(5): 848-53. PMID 23755923
- Klein EA, Yousefi K, Haddad Z, et al. A genomic classifier improves prediction of metastatic disease within 5 years after surgery in node-negative high-risk prostate cancer patients managed by radical prostatectomy without adjuvant therapy. Eur Urol. Apr 2015; 67(4): 778-86. PMID 25466945

- 80. Sommariva S, Tarricone R, Lazzeri M, et al. Prognostic Value of the Cell Cycle Progression Score in Patients with Prostate Cancer: A Systematic Review and Meta-analysis. Eur Urol. Jan 2016; 69(1): 107-15. PMID 25481455
- Feng FY, Huang HC, Spratt DE, et al. Validation of a 22-Gene Genomic Classifier in Patients With Recurrent Prostate Cancer: An Ancillary Study of the NRG/RTOG 9601 Randomized Clinical Trial. JAMA Oncol. Apr 01 2021; 7(4): 544-552. PMID 33570548
- 82. Den RB, Yousefi K, Trabulsi EJ, et al. Genomic classifier identifies men with adverse pathology after radical prostatectomy who benefit from adjuvant radiation therapy. J Clin Oncol. Mar 10 2015; 33(8): 944-51. PMID 25667284
- 83. Den RB, Feng FY, Showalter TN, et al. Genomic prostate cancer classifier predicts biochemical failure and metastases in patients after postoperative radiation therapy. Int J Radiat Oncol Biol Phys. Aug 01 2014; 89(5): 1038-1046. PMID 25035207
- 84. Cooperberg MR, Davicioni E, Crisan A, et al. Combined value of validated clinical and genomic risk stratification tools for predicting prostate cancer mortality in a high-risk prostatectomy cohort. Eur Urol. Feb 2015; 67(2): 326-33. PMID 24998118
- 85. Ross AE, Feng FY, Ghadessi M, et al. A genomic classifier predicting metastatic disease progression in men with biochemical recurrence after prostatectomy. Prostate Cancer Prostatic Dis. Mar 2014; 17(1): 64-9. PMID 24145624
- Karnes RJ, Bergstralh EJ, Davicioni E, et al. Validation of a genomic classifier that predicts metastasis following radical prostatectomy in an at risk patient population. J Urol. Dec 2013; 190(6): 2047-53. PMID 23770138
- Erho N, Crisan A, Vergara IA, et al. Discovery and validation of a prostate cancer genomic classifier that predicts early metastasis following radical prostatectomy. PLoS One. 2013; 8(6): e66855. PMID 23826159
- 88. Ross AE, Johnson MH, Yousefi K, et al. Tissue-based Genomics Augments Post-prostatectomy Risk Stratification in a Natural History Cohort of Intermediate- and High-Risk Men. Eur Urol. Jan 2016; 69(1): 157-65. PMID 26058959
- Freedland SJ, Choeurng V, Howard L, et al. Utilization of a Genomic Classifier for Prediction of Metastasis Following Salvage Radiation Therapy after Radical Prostatectomy. Eur Urol. Oct 2016; 70(4): 588-596. PMID 26806658
- Glass AG, Leo MC, Haddad Z, et al. Validation of a Genomic Classifier for Predicting Post-Prostatectomy Recurrence in a Community Based Health Care Setting. J Urol. Jun 2016; 195(6): 1748-53. PMID 26626216
- 91. Klein EA, Haddad Z, Yousefi K, et al. Decipher Genomic Classifier Measured on Prostate Biopsy Predicts Metastasis Risk. Urology. Apr 2016; 90: 148-52. PMID 26809071
- 92. Spratt DE, Dai DLY, Den RB, et al. Performance of a Prostate Cancer Genomic Classifier in Predicting Metastasis in Men with Prostate-specific Antigen Persistence Postprostatectomy. Eur Urol. Jul 2018; 74(1): 107-114. PMID 29233664
- 93. Karnes RJ, Choeurng V, Ross AE, et al. Validation of a Genomic Risk Classifier to Predict Prostate Cancer-specific Mortality in Men with Adverse Pathologic Features. Eur Urol. Feb 2018; 73(2): 168-175. PMID 28400167
- Ross AE, Den RB, Yousefi K, et al. Efficacy of post-operative radiation in a prostatectomy cohort adjusted for clinical and genomic risk. Prostate Cancer Prostatic Dis. Sep 2016; 19(3): 277-82. PMID 27136742
- 95. Lobo JM, Dicker AP, Buerki C, et al. Evaluating the clinical impact of a genomic classifier in prostate cancer using individualized decision analysis. PLoS One. 2015; 10(3): e0116866. PMID 25837660
- 96. West TA, Kiely BE, Stockler MR. Estimating scenarios for survival time in men starting systemic therapies for castration-resistant prostate cancer: a systematic review of randomised trials. Eur J Cancer. Jul 2014; 50(11): 1916-24. PMID 24825113
- 97. Scher HI, Graf RP, Schreiber NA, et al. Nuclear-specific AR-V7 Protein Localization is Necessary to Guide Treatment Selection in Metastatic Castration-resistant Prostate Cancer. Eur Urol. Jun 2017; 71(6): 874-882. PMID 27979426

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 50 of 54

- 98. Armstrong AJ, Halabi S, Luo J, et al. Prospective Multicenter Validation of Androgen Receptor Splice Variant 7 and Hormone Therapy Resistance in High-Risk Castration-Resistant Prostate Cancer: The PROPHECY Study. J Clin Oncol. May 01 2019; 37(13): 1120-1129. PMID 30865549
- Sanda MG, Cadeddu JA, Kirkby E, et al. Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part I: Risk Stratification, Shared Decision Making, and Care Options. J Urol. Mar 2018; 199(3): 683-690. PMID 29203269
- 100. Sanda MG, Cadeddu JA, Kirkby E, et al. Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part II: Recommended Approaches and Details of Specific Care Options. J Urol. Apr 2018; 199(4): 990-997. PMID 29331546
- 101. Antonarakis ES, Lu C, Wang H, et al. AR-V7 and resistance to enzalutamide and abiraterone in prostate cancer. N Engl J Med. Sep 11 2014; 371(11): 1028-38. PMID 25184630
- 102. Scher HI, Lu D, Schreiber NA, et al. Association of AR-V7 on Circulating Tumor Cells as a Treatment-Specific Biomarker With Outcomes and Survival in Castration-Resistant Prostate Cancer. JAMA Oncol. Nov 01 2016; 2(11): 1441-1449. PMID 27262168
- 103. Scher HI, Graf RP, Schreiber NA, et al. Assessment of the Validity of Nuclear-Localized Androgen Receptor Splice Variant 7 in Circulating Tumor Cells as a Predictive Biomarker for Castration-Resistant Prostate Cancer. JAMA Oncol. Sep 01 2018; 4(9): 1179-1186. PMID 29955787
- 104. Eggener SE, Rumble RB, Armstrong AJ, et al. Molecular Biomarkers in Localized Prostate Cancer: ASCO Guideline. J Clin Oncol. May 01 2020; 38(13): 1474-1494. PMID 31829902
- 105. Eastham JA, Auffenberg GB, Barocas DA, et al. Clinically Localized Prostate Cancer: AUA/ASTRO Guideline, Part I: Introduction, Risk Assessment, Staging, and Risk-Based Management. J Urol. Jul 2022; 208(1): 10-18. PMID 35536144
- 106. Lowrance WT, Murad MH, Oh WK, et al. Castration-Resistant Prostate Cancer: AUA Guideline Amendment 2018. J Urol. Dec 2018; 200(6): 1264-1272. PMID 30086276
- 107. National Institute for Health and Care Excellence (NICE). Prostate cancer: diagnosis and management [NG131]. 2019 (Updated 2021); https://www.nice.org.uk/guidance/ng131. Accessed November 2, 2022.

Documentation for Clinical Review

Please provide the following documentation:

- History and physical and/or consultation notes including:
 - o Clinical stage (TNM)
 - o NCCN risk category
 - o Biopsy Gleason score
 - o PSA level/density
 - o Number of biopsy cores with presence of disease including cancer involvement
 - o Pertinet comorbidities
- Test requested and reason for test
- Projected life expectancy
- Documentation that individual is a candidate for active surveillance or definitive therapy
- Prior treatment (if applicable) including prostatectomy if applicable

Post Service (in addition to the above, please include the following):

• Laboratory report/results

Coding

This Policy relates only to the services or supplies described herein. Benefits may vary according to product design; therefore, contract language should be reviewed before applying the terms of the Policy.

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 51 of 54

The following codes are included below for informational purposes. Inclusion or exclusion of a code(s) does not constitute or imply member coverage or provider reimbursement policy. Policy Statements are intended to provide member coverage information and may include the use of some codes for clarity. The Policy Guidelines section may also provide additional information for how to interpret the Policy Statements and to provide coding guidance in some cases.

Туре	Code	Description
CPT*	0005U	Oncology (prostate) gene expression profile by real-time RT-PCR of 3
		genes (ERG, PCA3, and SPDEF), urine, algorithm reported as risk score
	0011M	Oncology, prostate cancer, mRNA expression assay of 12 genes (10
		content and 2 housekeeping), RT-PCR test utilizing blood plasma
		and/or urine, algorithms to predict high-grade prostate cancer risk
	0047U	Oncology (prostate), mRNA, gene expression profiling by real-time RT-
		PCR of 17 genes (12 content and 5 housekeeping), utilizing formalin-fixed
		paraffin-embedded tissue, algorithm reported as a risk score
	81479	Unlisted molecular pathology procedure
	81541	Oncology (prostate), mRNA gene expression profiling by real-time RT-
		PCR of 46 genes (31 content and 15 housekeeping), utilizing formalin-
		fixed paraffin-embedded tissue, algorithm reported as a disease-
		specific mortality risk score
	81542	Oncology (prostate), mRNA, microarray gene expression profiling of 22
		content genes, utilizing formalin-fixed paraffin-embedded tissue,
		algorithm reported as metastasis risk score
	81599	Unlisted multianalyte assay with algorithmic analysis
HCPCS	None	

Policy History

This section provides a chronological history of the activities, updates and changes that have occurred with this Medical Policy.

Effective Date	Action
03/01/2016	BCBSA Medical Policy adoption
02/01/2017	Policy revision without position change
01/01/2018	Policy revision without position change
	Coding update
05/01/2018	Coding update
11/01/2018	Policy revision with position change
01/01/2019	Policy revision without position change
03/01/2020	Coding update
04/01/2020	Annual review. No change to policy statement.
01/01/2021	Annual review. No change to policy statement. Literature review updated.
12/01/2021	Administrative update. Policy statement updated.
02/01/2021	Annual review. Literature review updated. Policy statement and literature
02/01/2021	updated.
01/01/2023	Annual review. No change to policy statement. Literature review updated.

Definitions of Decision Determinations

Medically Necessary: Services that are Medically Necessary include only those which have been established as safe and effective, are furnished under generally accepted professional standards to treat illness, injury or medical condition, and which, as determined by Blue Shield, are: (a) consistent

2.04.111 Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management Page 52 of 54

with Blue Shield medical policy; (b) consistent with the symptoms or diagnosis; (c) not furnished primarily for the convenience of the patient, the attending Physician or other provider; (d) furnished at the most appropriate level which can be provided safely and effectively to the patient; and (e) not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of the Member's illness, injury, or disease.

Investigational/Experimental: A treatment, procedure, or drug is investigational when it has not been recognized as safe and effective for use in treating the particular condition in accordance with generally accepted professional medical standards. This includes services where approval by the federal or state governmental is required prior to use, but has not yet been granted.

Split Evaluation: Blue Shield of California/Blue Shield of California Life & Health Insurance Company (Blue Shield) policy review can result in a split evaluation, where a treatment, procedure, or drug will be considered to be investigational for certain indications or conditions, but will be deemed safe and effective for other indications or conditions, and therefore potentially medically necessary in those instances.

Prior Authorization Requirements and Feedback (as applicable to your plan)

Within five days before the actual date of service, the provider must confirm with Blue Shield that the member's health plan coverage is still in effect. Blue Shield reserves the right to revoke an authorization prior to services being rendered based on cancellation of the member's eligibility. Final determination of benefits will be made after review of the claim for limitations or exclusions.

Questions regarding the applicability of this policy should be directed to the Prior Authorization Department at (800) 541-6652, or the Transplant Case Management Department at (800) 637-2066 ext. 3507708 or visit the provider portal at <u>www.blueshieldca.com/provider</u>.

We are interested in receiving feedback relative to developing, adopting, and reviewing criteria for medical policy. Any licensed practitioner who is contracted with Blue Shield of California or Blue Shield of California Promise Health Plan is welcome to provide comments, suggestions, or concerns. Our internal policy committees will receive and take your comments into consideration.

For utilization and medical policy feedback, please send comments to: <u>MedPolicy@blueshieldca.com</u>

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. Blue Shield of California may consider published peer-reviewed scientific literature, national guidelines, and local standards of practice in developing its medical policy. Federal and state law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over medical policy and must be considered first in determining covered services. Member contracts may differ in their benefits. Blue Shield reserves the right to review and update policies as appropriate.

Appendix A

POLICY STATEMENT					
(No changes)					
BEFORE	AFTER				
Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management 2.04.111	Gene Expression Profiling and Protein Biomarkers for Prostate Cancer Management 2.04.111				
 Policy Statement: The one-time use of Oncotype DX® Prostate, Decipher® (prostate biopsy) or Prolaris® gene expression profiling tests may be considered medically necessary to guide management of prostate cancer when all of the following have been met: Needle biopsy discloses localized (i.e., less than T3) adenocarcinoma of prostate (with no clinical evidence of metastasis or lymph node involvement, i.e., NOMO) Individual falls into one of the following Risk categories as outlined by the National Comprehensive Cancer Network (NCCN) guidelines (see Policy Guidelines)*: Very Low risk Low risk No documentation of significant co-morbidities, advanced age, or other factors that suggest the individual might have a life expectancy of less than 10 years IV. No documentation the individual is not a candidate for either active surveillance or definitive therapy (e.g., external beam radiotherapy, brachytherapy, or radical prostatectomy) V. No documentation the individual has received prior treatment for prostate cancer 	 Policy Statement: The one-time use of Oncotype DX® Prostate, Decipher® (prostate biopsy) or Prolaris® gene expression profiling tests may be considered medically necessary to guide management of prostate cancer when all of the following have been met: Needle biopsy discloses localized (i.e., less than T3) adenocarcinoma of prostate (with no clinical evidence of metastasis or lymph node involvement, i.e., NOMO) Individual falls into one of the following Risk categories as outlined by the National Comprehensive Cancer Network (NCCN) guidelines (see Policy Guidelines)*: Very Low risk Forvarable Intermediate risk No documentation of significant co-morbidities, advanced age, or other factors that suggest the individual might have a life expectancy of less than 10 years No documentation the individual is not a candidate for either active surveillance or definitive therapy (e.g., external beam radiotherapy, brachytherapy, or radical prostatectomy) No documentation the individual has received prior treatment for prostate cancer 				
The one-time use of the Decipher or Prolaris gene expression profiling tests may be considered medically necessary in patients with unfavorable intermediate and high risk disease and no documentation of significant co- morbidities, advanced age, or other factors that suggest the individual might have a life expectancy of less than 10 years The one-time use of Decipher Prostate RP (genomic classifier) may be considered medically necessary to inform adjuvant treatment if adverse	II. The one-time use of the Decipher or Prolaris gene expression profiling tests may be considered medically necessary in patients with unfavorable intermediate and high risk disease and no documentation of significant co-morbidities, advanced age, or other factors that suggest the individual might have a life expectancy of less than 10 years				

POLICY STATEMENT					
(No changes)					
BEFORE	AFTER				
features are found after radical prostatectomy or during workup for radical	III. The one-time use of Decipher Prostate RP (genomic classifier) may				
prostatectomy with PSA persistence or recurrence (category 2B for	be considered medically necessary to inform adjuvant treatment if				
recurrence).	adverse features are found after radical prostatectomy or during				
	workup for radical prostatectomy with PSA persistence or				
The use of gene expression analysis other than listed above, and protein	recurrence (category 2B for recurrence).				
biomarkers to guide management of prostate cancer is considered					
investigational in all situations.	IV. The use of gene expression analysis other than listed above, and				
	protein biomarkers to guide management of prostate cancer is				
	considered investigational in all situations.				