

2.04.43	Genetic Testing for Cardiac Ion Channelopathies		
Original Policy Date:	July 6, 2012	Effective Date:	March 1, 2022
Section:	2.0 Medicine	Page:	Page 1 of 39

Policy Statement

Long QT Syndrome

Genetic testing to confirm a diagnosis of congenital long QT syndrome (LQTS) may be considered **medically necessary** when signs and/or symptoms of LQTS are present, but a definitive diagnosis cannot be made without genetic testing. This includes:

- I. Individuals who do not meet the clinical criteria for LQTS (i.e., those with a Schwartz score less than 4) but have a moderate-to-high pretest probability (see Policy Guidelines section) based on the Schwartz score and/or other clinical criteria.

Genetic testing of asymptomatic individuals to determine future risk of LQTS may be considered **medically necessary** when at least **one or more** of the following criteria is met:

- I. A [close relative](#) (i.e., first-, second-, or third-degree relative) with a known LQTS variant
- II. A [close relative](#) diagnosed with LQTS by clinical means whose genetic status is unavailable.

Genetic testing for LQTS for all other situations not meeting the criteria outlined above, including but not limited to determining prognosis and/or directing therapy in patients with known LQTS, is considered **investigational**.

Brugada Syndrome

Genetic testing to confirm a diagnosis of Brugada syndrome (BrS) may be considered **medically necessary** when [signs and/or symptoms](#) consistent with BrS (see Policy Guidelines section) are present, but a definitive diagnosis cannot be made without genetic testing.

Genetic testing of asymptomatic individuals to determine future risk of BrS may be considered **medically necessary** when patients have a [close relative](#) (i.e., first-, second-, or third-degree relative) with a known BrS variant.

Genetic testing for BrS for all other situations not meeting the criteria outlined above is considered **investigational**.

Catecholaminergic Polymorphic Ventricular Tachycardia

Genetic testing to confirm a diagnosis of catecholaminergic polymorphic ventricular tachycardia (CPVT) may be considered **medically necessary** when signs and/or symptoms of CPVT are present, but a definitive diagnosis cannot be made without genetic testing.

Genetic testing of asymptomatic individuals to determine future risk of CPVT may be considered **medically necessary** when at least **one or more** of the following criteria is met:

- I. A [close relative](#) (i.e., first-, second-, or third-degree relative) with a known CPVT variant
- II. A [close relative](#) diagnosed with CPVT by clinical means whose genetic status is unavailable.

Genetic testing for CPVT for all other situations not meeting the criteria outlined above is considered **investigational**.

Short QT Syndrome

Genetic testing of asymptomatic individuals to determine future risk of short QT syndrome (SQTS) may be considered **medically necessary** when patients have a [close relative](#) (i.e., first-, second-, or third-degree relative) with a known SQTS variant.

Genetic testing for SQTS for all other situations not meeting the criteria outlined above is considered **investigational**.

NOTE: Refer to [Appendix A](#) to see the policy statement changes (if any) from the previous version.

Policy Guidelines

First-, Second-, and Third-Degree Relatives

Note: Both maternal and paternal family histories are important and each lineage must be considered separately:

- First degree relatives are defined as a blood relative with whom the individual shares approximately 50% of his/her genes, including parents, full-siblings, and children on both maternal and paternal sides
- Second degree relatives are defined as a blood relative with whom an individual shares approximately 25% of his/her genes, including grandparents, grandchildren, uncles, aunts, nieces, nephews, and half-siblings
- Third degree relatives are defined as a blood relative with whom an individual shares approximately 12.5% of his/her genes, including first-cousins, great-grandparents or great-grandchildren

Genetic testing should be performed by an expert in genetic testing and/or cardiac ion channelopathies.

Determining the pretest probability of long QT syndrome (LQTS) is not standardized. An example of a patient with a moderate-to-high pretest probability of LQTS is a patient with a Schwartz score of 2 or 3.

Signs and symptoms suggestive of Brugada syndrome (BrS) include the presence of a characteristic electrocardiographic pattern, documented ventricular arrhythmia, sudden cardiac death (SCD) in a family member younger than 45 years old, a characteristic electrocardiographic pattern in a family member, inducible ventricular arrhythmias on electrophysiologic studies, syncope, or nocturnal agonal respirations. An index patient with suspected short QT syndrome (SQTS) would be expected to have a shortened (less than 2 standard deviation [SD] below from the mean) rate-corrected shortened QT interval (QTc). Cutoffs below 350 ms for men and 360 ms for women have been derived from population normal values (Tristani-Firouzi, 2014). The presence of a short QTc interval alone does not make the diagnosis of SQTS. Clinical history, family history, other electrocardiographic findings, and genetic testing may be used to confirm the diagnosis.

Testing Strategy

In general, testing for patients with suspected congenital LQTS, catecholaminergic polymorphic ventricular tachycardia (CPVT), or BrS should begin with a known familial variant, if one has been identified.

In cases where the family member's genetic diagnosis is unavailable, testing is available through either single-gene testing or panel testing. The evaluation of the clinical utility of panel testing is outlined in Blue Shield of California Medical Policy: General Approach to Evaluating the Utility of Genetic Panels. Panels for cardiac ion channelopathies are diagnostic test panels that may fall into one of several categories: panels that include variants for a single condition; panels that include variants for multiple conditions (indicated plus nonindicated conditions); and panels that include variants for multiple conditions (clinical syndrome for which clinical diagnosis not possible).

For situations in which a relative of a proband with unexplained cardiac death or unexplained sudden cardiac arrest (SCA) or an individual with unexplained SCA is being evaluated, genetic testing may be part of a diagnostic strategy that includes a comprehensive history and physical

exam and 12-lead electrocardiogram (ECG), along with exercise stress test, transthoracic echocardiography, and additional evaluation as guided by the initial studies. Studies have suggested that, in such cases, a probable diagnosis of an inherited cardiac condition can be made following a nongenetic evaluation in 50% to 80% of cases (Behr et al, 2008; Krahn et al, 2009; Kumar et al, 2013; Wong et al, 2014). If, after a comprehensive evaluation, a diagnosis of CPVT, LQTS, or BrS is suspected but not definitive (i.e., if there is a moderate-to-high pretest probability of either condition), genetic testing could be considered.

Genetic Counseling

Genetic counseling is primarily aimed at patients who are at risk for inherited disorders, and experts recommend formal genetic counseling in most cases when genetic testing for an inherited condition is considered. The interpretation of the results of genetic tests and the understanding of risk factors can be very difficult and complex. Therefore, genetic counseling will assist individuals in understanding the possible benefits and harms of genetic testing, including the possible impact of the information on the individual's family. Genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing. Genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

Coding

The following CPT genomic sequencing procedure codes may be used for this testing:

- **81413:** Cardiac ion channelopathies (e.g., Brugada syndrome, long QT syndrome, short QT syndrome, catecholaminergic polymorphic ventricular tachycardia); genomic sequence analysis panel, must include sequencing of at least 10 genes, including *ANK2*, *CASQ2*, *CAV3*, *KCNE1*, *KCNE2*, *KCNH2*, *KCNJ2*, *KCNQ1*, *RYR2*, and *SCN5A*
- **81414:** Cardiac ion channelopathies (e.g., Brugada syndrome, long QT syndrome, short QT syndrome, catecholaminergic polymorphic ventricular tachycardia); duplication/deletion gene analysis panel, must include analysis of at least 2 genes, including *KCNH2* and *KCNQ1*

The following CPT genomic sequencing procedure code is for this testing:

- **0237U:** Cardiac ion channelopathies (e.g., Brugada syndrome, long QT syndrome, short QT syndrome, catecholaminergic polymorphic ventricular tachycardia), genomic sequence analysis panel including *ANK2*, *CASQ2*, *CAV3*, *KCNE1*, *KCNE2*, *KCNH2*, *KCNJ2*, *KCNQ1*, *RYR2*, and *SCN5A*, including small sequence changes in exonic and intronic regions, deletions, duplications, mobile element insertions, and variants in non-uniquely mappable regions

Other analyses related to this testing are listed under the following CPT tier 2 molecular pathology codes:

- Under code **81403:** Molecular Pathology Procedure Level 4
 - *KCNJ2* (*potassium inwardly-rectifying channel, subfamily J, member 2*) (e.g., Andersen-Tawil syndrome), full gene sequence
- Under code **81404:** Molecular pathology procedure level 5
 - *SCN1B* (*sodium channel, voltage-gated, type I, beta*) (e.g., Brugada syndrome), full gene sequence
- Under code **81405:** Molecular Pathology Procedure Level 6
 - *CASQ2* (*calsequestrin 2 [cardiac muscle]*) (e.g., catecholaminergic polymorphic ventricular tachycardia), full gene sequence
- Under code **81406:** Molecular Pathology Procedure Level 7
 - *KCNH2* (*potassium voltage-gated channel, subfamily H[lead-related], member 2*) (e.g., short QT syndrome, long QT syndrome), full gene sequence
 - *KCNQ1* (*potassium voltage-gated channel, KQT-like subfamily, member 1*) (e.g., short QT syndrome, long QT syndrome), full gene sequence
- Under code **81407:** Molecular Pathology Procedure Level 8

- *SCN5A* (sodium channel, voltage-gated, type V, alpha subunit) (e.g., familial dilated cardiomyopathy), full gene sequence
- Under code **81408**: Molecular Pathology Procedure Level 9
 - *RYR2* (ryanodine receptor 2 [cardiac]) (e.g., catecholaminergic polymorphic ventricular tachycardia, arrhythmogenic right ventricular dysplasia), full gene sequence or targeted sequence analysis of > 50 exons

There is a HCPCS S code for testing for suspected Brugada syndrome:

- **S3861**: Genetic testing, sodium channel, voltage-gated, type V, alpha subunit (SCN5A) and variants for suspected Brugada syndrome

Description

Genetic testing is available for patients suspected of having cardiac ion channelopathies, including long QT syndrome (LQTS), catecholaminergic polymorphic ventricular tachycardia (CPVT), Brugada syndrome (BrS), and short QT syndrome (SQTS). These disorders are clinically heterogeneous and may range from asymptomatic to presenting with sudden cardiac death (SCD). Testing for variants associated with these channelopathies may assist in diagnosis, risk-stratify prognosis, and/or identify susceptibility for the disorders in asymptomatic family members.

Related Policies

- General Approach to Evaluating the Utility of Genetic Panels

Benefit Application

Benefit determinations should be based in all cases on the applicable contract language. To the extent there are any conflicts between these guidelines and the contract language, the contract language will control. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

Some state or federal mandates (e.g., Federal Employee Program [FEP]) prohibits plans from denying Food and Drug Administration (FDA)-approved technologies as investigational. In these instances, plans may have to consider the coverage eligibility of FDA-approved technologies on the basis of medical necessity alone.

Regulatory Status

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Amendments (CLIA). Laboratories that offer laboratory-developed tests must be licensed by the CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

Rationale

Background

Cardiac Ion Channelopathies

Cardiac ion channelopathies result from variants in genes that code for protein subunits of the cardiac ion channels. These channels are essential to cell membrane components that open or close to allow ions to flow into or out of the cell. Regulation of these ions is essential for the maintenance of a normal cardiac action potential. This group of disorders is associated with ventricular arrhythmias and an increased risk of sudden cardiac death (SCD). These congenital

cardiac channelopathies can be difficult to diagnose, and the implications of an incorrect diagnosis could be catastrophic.

The prevalence of any cardiac channelopathy is still ill-defined but is thought to be between 1 in 2000 and 1 in 3000 persons in the general population.¹ Data about the individual prevalences of long QT syndrome (LQTS), Brugada syndrome (BrS), catecholaminergic polymorphic ventricular tachycardia (CPVT), and short QT syndrome (SQTS) are presented in Table 1.

Table 1. Epidemiology of Cardiac Ion Channelopathies

Variables	LQTS	BrS	CPVT	SQTS
Prevalence	1:2000-5000	1:6000	1:7000-10,000	Unidentified
Annual mortality rate	0.3% (LQT1) 0.6% (LQT2) 0.56% (LQT3)	4% ^a	3.1%	Unidentified
Mean age at first event, y	14	42 ^a	15	40

Adapted from Modell et al (2012).²

BrS: Brugada syndrome; CPVT: catecholaminergic polymorphic ventricular tachycardia; LQTS: long QT syndrome; SQTS: short QT syndrome.

^a Type 1 electrocardiographic pattern.

Long QT Syndrome

Congenital LQTS is an inherited disorder characterized by the lengthening of the repolarization phase of the ventricular action potential, increasing the risk for arrhythmic events, such as torsades de pointes, which may, in turn, result in syncope and SCD.

Congenital LQTS usually manifests before the age of 40 years. It is estimated that more than half of the 8000 sudden unexpected deaths in children may be related to LQTS. The mortality rate of untreated patients with LQTS is estimated at 1% to 2% per year, although this figure will vary with the genotype.

Brugada Syndrome

Brugada syndrome is characterized by cardiac conduction abnormalities that increase the risk of syncope, ventricular arrhythmia, and SCD. The disorder primarily manifests during adulthood, although ages between 2 days and 85 years have been reported.³ Brugada syndrome is an autosomal dominant disorder with an unexplained male predominance. Males are more likely to be affected than females (approximate ratio, 8:1). Brugada syndrome is estimated to be responsible for 12% of SCD cases.¹ For both sexes, there is an equally high risk of ventricular arrhythmias or sudden death.⁴ Penetrance is highly variable, with phenotypes ranging from asymptomatic expression to death within the first year of life.⁵

Catecholaminergic Polymorphic Ventricular Tachycardia

Catecholaminergic polymorphic ventricular tachycardia is a rare, inherited channelopathy that may present with autosomal dominant or autosomal recessive inheritance. The disorder manifests as a bidirectional or polymorphic ventricular tachycardia precipitated by exercise or emotional stress.⁶ The prevalence of CPVT is estimated between 1 in 7000 and 1 in 10,000 persons. Catecholaminergic polymorphic ventricular tachycardia has a mortality rate of 30% to 50% by age 35 and is responsible for 13% of cardiac arrests in structurally normal hearts.⁶ Catecholaminergic polymorphic ventricular tachycardia was previously believed to manifest only during childhood, but studies have now identified presentation between infancy and 40 years of age.⁷

Short QT Syndrome

Short QT syndrome is characterized by a shortened QT interval on the electrocardiogram (ECG) and, at the cellular level, a shortening of the action potential.⁸ The clinical manifestations are an increased risk of atrial and/or ventricular arrhythmias. Because of the disease's rarity, the prevalence and risk of sudden death are currently unknown.⁶

Sudden Cardiac Arrest or Sudden Cardiac Death

Sudden cardiac arrest (SCA) and SCD refer to the sudden interruption of cardiac activity with circulatory collapse. The most common cause is coronary artery disease. Approximately 5% to 10% of SCA and SCD is due to arrhythmias without structural cardiac disease and are related to the primary electrical disease syndromes. The previously described cardiac ion channelopathies are among the primary electrical disease syndromes.

The evaluation and management of a survivor of SCA include an assessment of the circumstances of the event as well as a comprehensive physical examination emphasizing cardiovascular and neurologic systems, laboratory testing, ECG, and more advanced cardiac imaging or electrophysiologic testing as may be warranted. Genetic testing might be considered when, after completion of a comprehensive evaluation, there are findings consistent with a moderate-to-high likelihood of a primary electrical disease. Postmortem protocols for evaluation of a fatal SCA should be implemented when possible.

Genetics of Cardiac Ion Channelopathies

Long QT Syndrome

There are more than 1200 unique variants on at least 13 genes encoding potassium-channel proteins, sodium-channel proteins, calcium channel-related factors, and membrane adaptor proteins that have been associated with LQTS. In addition to single variants, some cases of LQTS are associated with deletions or duplications of genes.⁹

The absence of a variant does not imply the absence of LQTS; it is estimated that variants are only identified in 70% to 75% of patients with a clinical diagnosis of LQTS.¹⁰ A negative test is only definitive when there is a known variant identified in a family member and targeted testing for this variant is negative.

Another factor complicating interpretation of the genetic analysis is the penetrance of a given variant or the presence of multiple phenotypic expressions. For example, approximately 50% of variant carriers never have any symptoms. There is variable penetrance for LQTS, and penetrance may differ for the various subtypes. While linkage studies in the past have indicated that penetrance was 90% or greater, a 1999 analysis using molecular genetics challenged this estimate and suggested that penetrance may be as low as 25% for some families.¹¹

Variants involving *KCNQ1*, *KCNH2*, and *SCN5A* are the most commonly detected in patients with genetically confirmed LQTS. Some variants are associated with extra-cardiac abnormalities in addition to the cardiac ion channel abnormalities. A summary of clinical syndromes associated with hereditary LQTS is shown in Table 2. A 2021 analysis of 49 patients with channelopathies identified 3 rare variants that were pathogenic for LQTS and 8 rare variants that were likely pathogenic for LQTS, all involving *KCNQ1* or *KCNH2*.¹²

Table 2. Genetics of LQTS

Type	Other Names	Chromosome Locus	Mutated Gene	Ion Current(s) Affected	Associated Findings
LQT1	RWS	11p15.5-p.15.4	<i>KCNQ1</i>	Potassium	
LQT2	RWS	7q36.1	<i>KCNH2</i>	Potassium	
LQT3	RWS	3p22.2	<i>SCN5A</i>	Sodium	
LQT4	Ankyrin B syndrome	4q25-26	<i>ANK2</i>	Sodium, potassium, calcium	Catecholaminergic polymorphic ventricular arrhythmias, sinus node dysfunction, AF
LQT5	RWS	21q22.12	<i>KCNE1</i>	Potassium	
LQT6	RWS	21q22.11	<i>KNCE2</i>	Potassium	
LQT7	Andersen-Tawil syndrome	17.qq2432	<i>KCNJ2</i>	Potassium	Episodic muscle weakness, congenital anomalies

Type	Other Names	Chromosome Locus	Mutated Gene	Ion Current(s) Affected	Associated Findings
LQT8	Timothy syndrome	12q13.33	<i>CACNA1C</i>	Calcium	Congenital heart defects, hand/foot syndactyly, ASD
LQT9	RWS	3p25.3	<i>CAV3</i>	Sodium	
LQT10	RWS	11q23.3	<i>SCN4B</i>	Sodium	
LQT11	RWS	7q21.2	<i>AKAP9</i>	Potassium	
LQT12	RWS	20q11.21	<i>SNTA1</i>	Sodium	
LQT13	RWS	11q24.3	<i>KCNJ5</i>	Potassium	
LQT14		14q32.11	<i>CALM1</i>	Calmodulin	
LQT15		2p21	<i>CALM2</i>	Calmodulin	
LQT16		19q13.32	<i>CALM3</i>	Calmodulin	
JLNS1	JLNS	11p15.5-11p15.4	<i>KCNQ1</i> (homozygotes or compound heterozygotes)	Potassium	Congenital sensorineural hearing loss
JLNS2	JLNS	21q22.12	<i>KCNE1</i> (homozygotes or compound heterozygotes)	Potassium	Congenital sensorineural hearing loss

Adapted from Beckmann et al (2021),¹³ Arking et al (2014),¹⁴ and Alders (2015).¹⁵

AF: atrial fibrillation; ASD: autism spectrum disorder; LQT: long QT; LQTS: long QT syndrome; JLNS: Jervell and Lange-Nielsen syndrome; RWS: Romano-Ward syndrome.

Brugada Syndrome

Brugada syndrome is typically inherited in an autosomal dominant manner with incomplete penetrance. The proportion of cases that are inherited, versus de novo variants, is uncertain. Although some have reported up to 50% of cases are sporadic, others have reported that the instance of de novo variants is very low and is estimated to be only 1% of cases.⁴

Variants in 16 genes have been identified as causative of BrS, all of which lead to a decrease in the inward sodium or calcium current or an increase in one of the outward potassium currents. Of these, *SCN5A* is the most important, accounting for more than an estimated 20% of cases⁷; *SCN10A* has also been implicated. The other genes are of minor significance and account together for approximately 5% of cases.⁶ The absence of a positive test does not indicate the absence of BrS, with more than 65% of cases not having an identified genetic cause. Penetrance of BrS among persons with an *SCN5A* variant is 80% when undergoing ECG with sodium-channel blocker challenge and 25% when not using the ECG challenge.⁴ A 2021 analysis of 49 patients with channelopathies identified 1 rare variant that was pathogenic for BrS and 3 rare variants that were likely pathogenic for BrS, all involving the *SCN5A* gene.¹²

Catecholaminergic Polymorphic Ventricular Tachycardia

Variants in 4 genes are known to cause CPVT, and investigators believe other unidentified loci are involved as well. Currently, only 55% to 65% of patients with CPVT have an identified causative variant. Variants of the gene encoding the cardiac ryanodine receptor (*RYR2*) or to *KCNJ2* result in an autosomal dominant form of CPVT. *CASQ2* (cardiac calsequestrin) and *TRDN*-related CPVT exhibit autosomal recessive inheritance. A channelopathy expert panel review has also found moderate to definitive evidence for an autosomal dominant inheritance of *CALM1*, *CALM2*, and *CALM3* and an autosomal recessive inheritance of *TECRL*.¹⁶ Some have reported heterozygotes for *CASQ2* and *TRDN* variants for rare, benign arrhythmias.¹⁷ *RYR2* variants represent most CPVT cases (50% to 55%), with *CASQ2* accounting for 1% to 2% and *TRDN* accounting for an unknown proportion of cases. The penetrance of *RYR2* variants is approximated at 83%.¹⁷

An estimated 50% to 70% of patients will have the dominant form of CPVT with a disease-causing variant. Most variants (90%) to *RYR2* are missense variants, but in a small proportion of unrelated CPVT patients, large gene rearrangements or exon deletions have been reported.⁷ Additionally, nearly a third of patients diagnosed as LQTS with normal QT intervals have CPVT due to identified *RYR2* variants. Another misclassification, CPVT diagnosed as Anderson-Tawil syndrome

may result in more aggressive prophylaxis for CPVT whereas a correct diagnosis can spare this treatment because Anderson-Tawil syndrome is rarely fatal.

Short QT Syndrome

Short QT syndrome has been linked predominantly to variants in 3 genes (KCNH2, KCNJ2, KCNQ1).¹⁴ Variants in genes encoding alpha- and beta-subunits of the L-type cardiac calcium channel (CACNA1C, CACNB2) have also been associated with SQTs. Some individuals with SQTs do not have a variant in these genes, suggesting changes in other genes may also cause this disorder. A channelopathy expert panel concluded that only *KCNH2* had a definitive relationship with SQTs and *KCNQ1*, *KCNJ2*, and *SLC4A3* had strong to moderate causative evidence.¹⁶ Short QT syndrome is believed to be inherited in an autosomal dominant pattern. Although sporadic cases have been reported, patients frequently have a family history of the syndrome or SCD.

Literature Review

(see Appendix Table 1 for genetic testing categories).

Evidence reviews assess whether a medical test is clinically useful. A useful test provides information to make a clinical management decision that improves the net health outcome. That is, the balance of benefits and harms is better when the test is used to manage the condition than when another test or no test is used to manage the condition.

The first step in assessing a medical test is to formulate the clinical context and purpose of the test. The test must be technically reliable, clinically valid, and clinically useful for that purpose. Evidence reviews assess the evidence on whether a test is clinically valid and clinically useful. Technical reliability is outside the scope of these reviews, and credible information on technical reliability is available from other sources.

Genetic Testing for Variants Associated with Cardiac Ion Channelopathies

Clinical Context and Test Purpose

The purpose of genetic testing in patients with unexplained cardiac arrhythmias and/or other conduction abnormalities is to confirm the presence or absence of a cardiac ion channelopathy and inform clinical management.

The question addressed in this evidence review is: Does genetic testing for cardiac ion channelopathies (e.g., long QT syndrome [LQTS], Brugada syndrome [BrS], catecholaminergic polymorphic ventricular tachycardia [CPVT], short QT syndrome [SQTs]) improve health outcomes in individuals with suspected channelopathies or in individuals with a close relative with known or suspected channelopathies?

The following PICO was used to select literature to inform this review.

Populations

The population of interest are individuals with suspected cardiac ion channelopathies or asymptomatic individuals with a close relative(s) with a known cardiac ion channelopathy variants.

The channelopathies discussed herein are genetically heterogeneous with hundreds of identified variants, but the group of disorders shares basic clinical expression. The most common presentation is spontaneous or exercise-triggered syncope due to ventricular dysrhythmia. These can be self-limiting or potentially lethal cardiac events. The electrocardiographic features of each channelopathy are characteristic, but the electrocardiogram (ECG) is not diagnostic in all cases, and some secondary events (e.g., electrolyte disturbance, cardiomyopathies, or subarachnoid hemorrhage) may result in an ECG similar to those observed in a cardiac channelopathy.

Interventions

The intervention being considered is genetic testing for variants associated with cardiac ion channelopathies. Genetic tests are conducted in clinical laboratories. Genetic testing should be accompanied by genetic counseling including discussions with the patient or guardians about the importance and interpretation of genetic information and sharing of information with potentially affected family members as appropriate.

Genetic testing can be comprehensive (testing for all possible variants in multiple genes) or targeted (testing for a single variant identified in a family member). For comprehensive testing, the probability that a specific variant is pathophysiologically significant is greatly increased if the same variant has been reported in other cases. A variant may also be found that has not been associated with a disorder and therefore may or may not be pathogenic. Variants are classified by their pathogenic potential; an example of such a classification system used in the Familion assay is as follows in Table 3.

Table 3. Familion Assay Classification System

Class	Description
I	Deleterious and probable deleterious mutations. They are mutations that have either previously been identified as pathogenic (deleterious mutations), represent a major change in the protein, or cause an amino acid substitution in a critical region of the protein(s) (probable deleterious mutations).
II	Possible deleterious mutations. These variants encode changes to protein(s) but occur in regions that are not considered critical. Approximately 5% of unselected patients without LQTS will exhibit mutations in this category.
III	Variants not generally expected to be deleterious. These variants encode modified protein(s); however, they are considered more likely to represent benign polymorphisms. Approximately 90% of unselected patients without LQTS will have one or more of these variants; therefore patients with only class III variants are considered "negative."
IV	Non-protein-altering variants. These variants are not considered to have clinical significance and are not reported in the results of the Familion test.

LQTS: long QT syndrome.

Genetic testing for specific disorders, which may include 1 or more specific genes, is available from multiple academic and commercial laboratories, generally by next-generation sequencing or Sanger sequencing. Also, panel testing for 1 or more cardiac ion channelopathies is available from a number of genetic diagnostics laboratories, but there is some variation among manufacturers on the included genes.

There are also commercially available panels that include genetic testing for cardiac ion channelopathies along with other hereditary cardiac disorders, such as hypertrophic cardiomyopathy, dilated cardiomyopathy, and arrhythmogenic right ventricular cardiomyopathy.

Comparators

The following practice is currently being used: standard management without genetic testing. Diagnosis and management are described in the following sections by the condition.

Long QT Syndrome Diagnosis

The Schwartz criteria are commonly used as a diagnostic scoring system for LQTS.¹⁸ The most recent version is shown in Table 4. A score of 3.5 or higher indicates a high probability that LQTS is present; a score of 1.5 to 3, an intermediate probability; and a score of 1 or less indicates a low probability of the disorder. Before the availability of genetic testing, it was not possible to test the sensitivity and specificity of this scoring system, and because there is still no perfect criterion standard for diagnosing LQTS, the accuracy of this scoring system remains ill-defined.

Table 4. Diagnostic Scoring System for LQTS

Schwartz Criteria	Points
Electrocardiographic findings	
QT corrected >480 ms	3
QT corrected 460-470 ms	2
QT corrected <450 ms	1
History of torsades de pointes	2
T-wave alternans	1
Notched T waves in 3 leads	1
Low heart rate for age	0.5
Clinical history	
Syncope brought on by stress	2
Syncope without stress	1
Congenital deafness	0.5
Family history	
Family members with definite LQTS	1
Unexplained sudden death in immediate family members <30 y of age	0.5

Adapted from Perrin and Gollob (2012).¹⁹.

LQTS: long QT syndrome.

Long QT Standard Management

Primary management of asymptomatic or symptomatic long QT is β -blocker treatment with an intensification of therapy, if necessary due to recurrent arrhythmic events or intolerable side effects, including additional medication, left cardiac sympathetic denervation or placement of an implantable cardioverter-defibrillator (ICD). Avoidance of medications known to prolong the QT interval and the aggressive treatment of electrolyte imbalances are also advised.

Brugada Diagnosis

The diagnosis of BrS is made by the presence of a type 1 Brugada pattern on the ECG in addition to other clinical features.²⁰ This ECG pattern includes a coved ST-segment and a J-point elevation of 0.2 mV or higher followed by a negative T wave. This pattern should be observed in 2 or more of the right precordial ECG leads (V₁-V₃). This pattern may be concealed and can be revealed by administering a sodium-channel-blocking agent (e.g., flecainide).²¹ Two additional ECG patterns have been described (type 2, type 3) but are less specific for the disorder.²² The diagnosis of BrS is considered definitive when the characteristic ECG pattern is present with at least 1 of the following clinical features: documented ventricular arrhythmia, sudden cardiac death (SCD) in a family member younger than 45 years old, characteristic ECG pattern in a family member, inducible ventricular arrhythmias on electrophysiology studies, syncope, or nocturnal apneal respirations.

Brugada Standard Management

Management has focused on the use of ICDs in patients with syncope or cardiac arrest and isoproterenol for electrical storms. Patients who are asymptomatic can be closely followed to determine if ICD implantation is necessary.

Catecholaminergic Polymorphic Ventricular Tachycardia Diagnosis

Patients generally present with syncope or cardiac arrest during the first or second decade of life. The symptoms are nearly always triggered by exercise or emotional stress. The resting ECG of patients with CPVT is typically normal, but exercise stress testing can induce a ventricular arrhythmia in most cases (75% to 100%).¹⁹ Premature ventricular contractions, couplets, bigeminy, or polymorphic ventricular tachycardia (VT) are possible outcomes to the ECG stress test. For patients who are unable to exercise, an infusion of epinephrine may induce ventricular arrhythmia, but this is less effective than exercise testing.²³

Catecholaminergic Polymorphic Ventricular Tachycardia Standard Management

Management of CPVT is primarily with the β -blockers nadolol (1 to 2.5 mg/kg/d) or propranolol (2 to 4 mg/kg/d). If protection is incomplete (i.e., recurrence of syncope or arrhythmia), then flecainide (100 to 300 mg/d) may be added. If recurrence continues, an ICD may be necessary

with optimized pharmacologic management continued post-implantation.¹⁷ Lifestyle modification with the avoidance of strenuous exercise is recommended for all CPVT patients.

Short QT Diagnosis

Patients generally present with syncope, pre-syncope, or cardiac arrest. An ECG with a corrected QT interval less than 330 ms, sharp T wave at the end of the QRS complex, and a brief or absent ST-segment are characteristic of the syndrome.²⁴ However, higher QT intervals on ECG might also indicate SQTS, and the clinician has to determine if this is within the normative range of QT values. An index patient with suspected SQTS would be expected to have a shortened (<2 standard deviations below from the mean) rate-corrected shortened QT interval (QTc). Cutoffs below 350 ms for men and 360 ms for women have been derived from population normal values.²⁵ The length of the QT interval was not associated with severity of symptoms in a 2006 series of 29 patients with SQTS.²⁶ Electrophysiologic studies may be used to diagnose SQTS if the diagnosis is uncertain to evaluate for short refractory periods and inducible VT. However, in the series of 29 patients with SQTS described above, VT was inducible in only 3 of 6 subjects who underwent an electrophysiologic study.²⁶ A diagnostic scoring system was proposed by Gollob et al (2011) to help decision making after a review of 61 SQTS cases (see Table 5).²⁷

Table 5. Diagnostic Scoring System for SQTS

Gollob Criteria	Points
Electrocardiographic findings	
QT corrected <370 ms	1
QT corrected <350 ms	2
QT corrected <330 ms	3
J point-T peak interval <120 ms	1
Clinical history	
History of SCD	2
Documented polymorphic ventricular fibrillation or VT	2
Unexplained syncope	1
AF	1
Family history	
First- or second-degree relative with high probability SQTS	2
First- or second-degree relative with autopsy-negative SCD	1
Sudden infant death syndrome	1
Genotype	
Genotype positive	2
Mutation of undetermined significance in a culprit gene	1

Adapted from Perrin and Gollob (2012).¹⁹

AF: atrial fibrillation; SCD: sudden cardiac death; SQTS: short QT syndrome; VT: ventricular tachycardia.

Short QT Standard Management

The primary management of SQTS is with ICD therapy. Decisions about ICD therapy are based on the degree to which SQTS is considered likely, which depends on ECG features, family history, personal history of cardiac arrest or ventricular arrhythmias, and the ability to induce VT on electrophysiologic studies.

Anti-arrhythmic drug management of the disease is complicated because the binding target for QT-prolonging drugs (e.g., sotalol) is Kv11.1, which is coded for by *KCNH2*, the most common site for variants in SQTS (subtype 1). Treatment with quinidine (which is able to bind to both open and inactivated states of Kv11.1) is an appropriate QT-prolonging treatment. This treatment has been reported to reduce the rate of arrhythmias from 4.9% to 0% per year. For those with recurrence while on quinidine, an ICD is recommended.¹⁹

Outcomes

The general outcomes of interest are overall survival (OS), test validity, changes in reproductive decision making, and morbid events (e.g., cardiac events).

A positive diagnosis of LQTS or CPVT in symptomatic patients may lead to treatment with β -blockers or with ICDs, which can reduce the risk for ventricular arrhythmias and SCD. A positive test for BrS in symptomatic patients may influence the decision for treatment with an ICD.

It is unknown how a positive SQTs test in symptomatic patients would influence treatment decisions.

Positive tests in asymptomatic family members can inform lifestyle changes and prevention treatment decisions.

The genetic assays may be recommended as part of a diagnostic strategy for patients who exhibit clinical symptoms that are not considered definitive.

The tests may also be recommended for asymptomatic family members of patients with known cardiac ion channel variants.

The evidence related to the clinical validity and utility of genetic testing for the cardiac channelopathies consists primarily of studies that evaluate the yield of genetic testing and the impact of genetic testing on the diagnosis and subsequent management of a specific cardiac channelopathy. Many cardiac channelopathies lead to a common clinical outcome - increased risk of ventricular arrhythmias leading to an increased risk of SCD. Studies that evaluate the role of genetic testing for cardiac channelopathies as part of a diagnostic strategy in the evaluation of ventricular fibrillation or SCD from an unknown cause are discussed separately.

The evidence is presented as follows. First, for patients who are candidates for testing of specific channelopathies (LQTS, BrS, CPVT, SQTs) and asymptomatic family members of variant-positive probands. Finally, the evidence is presented for the genetic testing of family members in cases of SCD when a specific clinical diagnosis has not been made.

Study Selection Criteria

For the evaluation of clinical validity of genetic testing for specific cardiac ion channelopathies, studies that meet the following eligibility criteria were considered:

- Reported on the yield of genetic testing in patients with suspected or confirmed channelopathy;
- Included clinical diagnosis;
- Patient/sample clinical characteristics were described;
- Patient/sample selection criteria were described.

In addition, studies reporting on the clinical specificity will be discussed briefly when available.

Genetic Testing for the Diagnosis of Specific Cardiac Ion Channelopathies Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

The true clinical sensitivity and specificity of genetic testing for specific cardiac ion channelopathies cannot be determined with certainty because there is no independent criterion standard for the diagnosis. The clinical diagnosis can be compared with the genetic diagnosis, and vice versa, but neither the clinical diagnosis nor the results of genetic testing can be considered an adequate criterion standard.

Survivors of Sudden Cardiac Arrest

Asatryan et al (2019) evaluated the diagnostic validity and clinical utility of genetic testing in sudden cardiac arrest (SCA) survivors (n=60) with or without previous clinical evidence of heart

disease.²⁸ Patients without coronary artery disease were included; 24 (40%) with clear detectable cardiac phenotype [Ph(+)]SCA and 36 (60%) with no clear cardiac phenotype [Ph(-)]SCA. Targeted exome sequencing was performed using the TruSight-One Sequencing Panel (Illumina). A total of 32 pathogenic or likely pathogenic gene variants were found in 27 (45%) patients: 17 (71%) in the Ph(+)]SCA group and 10 (28%) in the Ph(-)]SCA group. Mutations in 16 (67%) Ph(+)]SCA patients were congruent with the suspected phenotype, consisting of 12 (50%) cardiomyopathies and 4 (17%) channelopathies. Mutations in 6 (17%) Ph(-)]SCA patients revealed a cardiac ion channelopathy explaining their SCA event. An additional 4 (11%) mutations in this group could not explain the phenotype and require additional studies. Overall, cardiac genetic testing was positive in 2/3 of the Ph(+)]SCA group and 1/6 of the Ph(-)]SCA group. The study was limited in its description of clinical criteria for establishing a diagnostic clinical phenotype. While the authors suggest the testing was useful to identify or confirm an inherited heart disease, with important impact on patient care and first-degree relatives at risk, health outcomes pertaining to clinical management of patients or asymptomatic familial probands were not reported.

Chiu et al (2021) performed genetic tests on 36 survivors of pediatric cardiac arrest (median age, 13.3 years).²⁹ The yield rate of genetic testing in the study cohort was 84.6%, including 14 pathogenic and 8 likely pathogenic variants. Long QT syndrome, CPVT, and BrS were diagnosed in 25%, 16.7%, and 6% of patients, respectively; genetic testing led to a change in diagnosis from CPVT to LQTS in 1 patient. Assessment of long-term outcomes showed that 10-year transplant-free survival was higher among patients who received genetic testing soon after the cardiac arrest event. Subsequent testing of family members of 15 probands identified 8 family members with positive genetic tests, but information on subsequent management of these patients was lacking.

Long QT Syndrome

Tester et al (2006) completed the largest study to evaluate the percentage of individuals with a clinical diagnosis of LQTS found to have a genetic variant.³⁰ The sample was 541 consecutive patients referred for evaluation of LQTS. Clinical assessments of the patients were made while blinded to the genetic testing results. Among the 123 patients with a high probability of LQTS based on clinical assessments, defined as a Schwartz score of 4 or more, 72% (89/123) had a genetic variant. Among patients with a QTc greater than 480 ms, 62% had a genetic variant. Characteristics and results of selected studies are shown in Tables 6 and 7 below.

Table 6. Characteristics of Clinical Validity Studies of Genetic Testing for LQTS

Study	Study Population	Design	Clinical Diagnosis	Genes Included	Blinding of Assessors
Tester et al (2006) ³⁰	Unrelated patients referred to Mayo Clinic's Sudden Death Genomics Laboratory for LQTS genetic testing from 1997 to 2004	Consecutive; prospective	Schwartz and Moss score (≥ 4) suggests strong probability for LQTS)	Unclear but described as 'comprehensive mutational analysis'	Yes
Bai et al (2009) ³¹	Patients from a sample of 1394 consecutive probands with either a clinically confirmed or suspected diagnosis of LQTS, BrS, or CPVT or a personal or family history of idiopathic ventricular fibrillation/cardiac arrest/SCD referred for molecular diagnosis	Consecutive; prospective	Diagnosed clinically as conclusive or possible; criteria not specified	KCNQ1, KCNH2, SCN5A, KCNE1, and KCNE2	NR

BrS: Brugada syndrome; CPVT: catecholaminergic polymorphic ventricular tachycardia; LQTS: long QT syndrome; NR: not reported; SCD: sudden cardiac death.

Table 7. Yield of Genetic Testing for LQTS

Study	N	Excluded Samples	Yield of Genetic Testing
Tester et al (2006) ³⁰ .	541	None	NR
Schwartz and Moss ≥ 4	123	Unknown Schwartz/Moss (n=124)	72%
Schwartz and Moss < 4	294	Unknown Schwartz/Moss (n=124)	44%
Bai et al (2009) ³¹ .	546	NR	40%
Conclusive Dx	304	NR	64%
Possible Dx	160	NR	14%

Dx: diagnosis; LQTS: long QT syndrome; NR: not reported.

The purpose of limitations tables (see Tables 8 and 9) is to display notable limitations identified in each study. This information is synthesized as a summary of the body of evidence and provides the conclusions on the sufficiency of the evidence supporting the position statement.

Table 8. Study Relevance Limitations of Clinical Validity Studies of Genetic Testing for LQTS

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Tester et al (2006) ³⁰ .		1: Not clear which genes were tested			
Bai et al (2009) ³¹ .	3: Criteria for clinical diagnosis unclear				

LQTS: long QT syndrome.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

Table 9. Study Design and Conduct Limitations of Clinical Validity Studies of Genetic Testing for LQTS

Study	Selection ^a	Blinding ^b	Test Delivery ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
Tester et al (2006) ³⁰ .					2: Insufficient data for clinical score in 23% of samples that had genetic testing	
Bai et al (2009) ³¹ .		1: Blinding not described			1: No description of exclusions or indeterminate results	

LQTS: long QT syndrome.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

^b Blinding key: 1. Not blinded to results of reference or other comparator tests.

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

The evidence on clinical specificity focuses on the frequency and interpretation of variants identified but not known to be pathogenic. If a variant identified is known to be pathogenic, then the specificity of this finding is high. However, many variants are not known to be pathogenic, and the specificity for these variants is lower. The rate of identification of variants is estimated at 5% for patients who do not have LQTS.³²

A 2012 publication from the National Heart, Lung, and Blood Institute GO Exome Sequencing Project (ESP) reported on the rate of sequence variants in a large number of patients without LQTS.³³ The ESP sequenced all genome regions of protein-coding in a sample of 5400 persons drawn from various populations, none of whom specifically had heart disease and/or channelopathies. Exome data were systematically searched to identify sequence variants previously associated with LQTS, including both nonsense variants, which are generally pathogenic, and missense variants, which are less likely to be pathogenic. Thirty-three such sequence variants were identified in the total population—all missense variations. The percentage of the population that had at least 1 of these missense variants was 5.2%. No nonsense variants were associated with LQTS found among the entire population.

Brugada Syndrome

Priori et al (2000) reported an early paper to describe the yield of genetic testing for BrS.³⁴ In 58 probands with a clinical diagnosis of BrS, the yield of *SCN5A* testing was 15%.

Kapplinger et al (2010) reported results from an international compendium of *SCN5A* variants of more than 2000 patients referred for BrS genetic testing which yielded almost 300 distinct mutations in 438 of 2111 (21%) patients, ranging from 11% to 28% across the 9 testing centers.³⁵ Hu et al (2014) evaluated the prevalence of *SCN10A* variants in 120 probands with BrS.³⁶ Seventeen *SCN10A* variants were identified in 25 probands, with a variant detection rate of 16.7% in BrS probands.

Behr et al (2015) evaluated 7 candidate genes (*SCN10A*, *HAND1*, *PLN*, *CASQ2*, *TKT*, *TBX3*, *TBX5*) among 156 patients negative for *SCN5A* variants with symptoms indicative of BrS (64%) and/or a family history of sudden death (47%) or BrS (18%).³⁷ Eighteen (11.5%) patients were found to have variants, most often in *SCN10A* (12/18 [67%]).

Andorin et al (2016) described the yield of *SCN5A* genetic testing in 75 patients younger than 19 years of age from 62 families who had a Brugada type I ECG pattern; only 20% were symptomatic.³⁸ The ECG pattern was spontaneous in 34% and drug-induced in 66%. The yield was very high compared to previous studies at 77%. The authors hypothesized that the high yield might have been due to the inclusion of only a pediatric population.

Chen et al (2019) conducted a meta-analysis of 17 studies involving 1780 unrelated and consecutive patients with BrS to assess the relationship between *SCN5A* mutation status and phenotypic features.³⁹ A history of syncope and spontaneous type 1 ECG pattern were observed in 31% and 59% of BrS patients, respectively. A total of 52% of patients had ICD implantation. The average frequency of *SCN5A* mutations was 20%, which ranged from 11% to 43% across studies. The onset of symptoms was found to occur at a younger age in the *SCN5A*(+) group (34 ± 17 vs. 42 ± 16 years; p=.0003). The presence of a spontaneous type 1 ECG pattern was associated with an increased risk of cardiac events in BrS patients based on a pooled analysis of 12 studies (71% vs. 57%; p=.0002). *SCN5A*(+) patients had a higher proportion of sick sinus syndrome (43% vs. 5%; p<.001) and atrial ventricular block (71% vs. 30%; p=.01). However, there was a lower rate of VT/ventricular fibrillation inducibility during electrophysiology

study(41% vs. 51%; $p=.01$), which may partially be explained by heterogeneity in electrophysiology study protocols. The *SCN5A* mutation was associated with an increased risk of major adverse events in the overall BrS (odds ratio, 1.78; 95% confidence interval [CI], 1.19 to 2.26; $p=.005$), Asian (odds ratio, 1.82; 95% CI, 1.07 to 3.11; $p=.03$), and Caucasian (odds ratio, 2.24; 95% CI, 1.02 to 4.90; $p=.04$) patient population.

Monasky et al (2019) evaluated 15 BrS-associated genes (*CACNA1C*, *CACNA2D1*, *CACNB2*, *GPD1L*, *HCN4*, *KCND2*, *KCND3*, *PKP2*, *RANGRF*, *SCN10A*, *SCN1B*, *SCN2B*, *SCN3B*, *SCN5A*, and *TRPM4*) with the TruSight One sequencing kit and NextSeq platform in 297 BrS patients screened for study enrollment.⁴⁰ The 2 most common mutations were *SCN5A* (84 [28.3%]) followed by *SCN10A* (8 [2.7%]). Clinical characteristics of BrS patients harboring *SCN5A* or *SCN10A* mutations were not found to be significantly different between probands, although patients with a variety of type I-III ECG patterns were represented in both cohorts.

Sacilotto et al (2020) reported data from the Genetics of Brazilian Arrhythmias (GenBra) registry.⁴¹ From 1999 to 2020, 138 (22 symptomatic) consecutive patients with type-1 BrS were assessed for invasive and noninvasive parameters and *SCN5A* mutation status. No difference in the rate of *SCN5A*-positive patients was found between asymptomatic and symptomatic groups (20/76 [26.3%] vs. 5/17 [29.4%]; $p=.770$). *SCN5A* carriers had a significantly higher frequency of aVR sign, S wave, and QRS-f.

Milman et al (2021) published an observational study of 678 patients from 14 countries with a first arrhythmic event due to BrS.⁴² Of the 392 probands, 23.5% were *SCN5A*(+) with 44 pathogenic/likely pathogenic variants and 48 variants of unknown significance. The remaining probands were *SCN5A*(-). Patients with pathogenic/likely pathogenic variants were more likely to be aged <16 years ($p=.023$), female ($p=.013$), and have a family history of SCD ($p<.001$) compared to patients who were *SCN5A*(-). Logistic regression found that White ethnicity (odds ratio, 5.41; 95% CI, 2.8 to 11.19; $p<.001$) and family history of SCD (odds ratio, 2.73; 95% CI, 1.28 to 5.82; $p=.009$) were associated with having a pathogenic/likely pathogenic genotype.

The description of the studies are below in Table 10 and results are shown in Table 11.

Table 10. Characteristics of Clinical Validity Studies of Genetic Testing for BrS

Study	Study Population	Design	Clinical Diagnosis	Genes Included	Blinding of Assessors
Priori et al (2000) ³⁴ .	Patients with the typical Brugada ECG pattern, without structural heart disease	Retrospective	Clinical and ECG diagnosis, criteria not specified	<i>SCN5A</i>	Unclear
Bai et al (2009) ³¹ .	Patients from a sample of 1394 consecutive probands with either a clinically confirmed or suspected diagnosis of LQTS, BrS, or CPVT or a personal or family history of idiopathic ventricular fibrillation/cardiac arrest/SCD referred for molecular diagnosis	Consecutive; prospective	Diagnosed clinically as conclusive or possible; criteria not specified	<i>SCN5A</i>	NR
Kapplinger et al (2010) ³⁵ .	Unrelated cases of clinically suspected BrS from international BrS databases (5 Europe, 3 United States, 1 Japan)	Retrospective; unclear whether the samples were consecutive	Referring physician made a clinical diagnosis of either possible or definite BrS, criteria not specified	27 translated exons in <i>SCN5A</i>	Unclear

Study	Study Population	Design	Clinical Diagnosis	Genes Included	Blinding of Assessors
Hu et al (2014) ³⁶ .	Unrelated patients with BrS referred to a single-center for genetic testing	Retrospective; not clear if selection was consecutive	2005 Consensus Conference diagnostic criteria (Heart Rhythm Society and the European Heart Rhythm Association)	SCN10A	Unclear
Behr et al (2015) ³⁷ .	Unrelated BrS Caucasian patients negative for SCN5A variants with symptoms and/or a family history of sudden death or BrS from 8 centers in Europe and U.S.	Retrospective; not clear if selection was consecutive	Locally diagnosed, criteria not specified	SCN10A, HAND1, CASQ2, TKT, PLN, TBX5, TBX3	Unclear
Andorin et al (2016) ³⁸ .	Patients (some from same family) <19 years of age at "diagnosis" of BrS (based on ECG pattern alone) in 16 European hospitals; 20% were symptomatic	Retrospective; not clear if selection was consecutive	Brugada type 1 ECG pattern either spontaneously or after challenge with a sodium channel blocker	SCN5A	Unclear
Chen et al (2019) ³⁹ .	Unrelated BrS patients >16 years of age with spontaneous or drug-induced type 1 ECG pattern from 17 studies in Japan, Europe, China, and others; 59% were spontaneously symptomatic	Meta-analysis; consecutive	Spontaneous or induced Brugada type 1 ECG pattern	SCN5A	NR
Monasky et al (2019) ⁴⁰ .	BrS patients (some from same family) with spontaneous or inducible arrhythmia	Prospective; not clear if selection was consecutive	Clinical diagnosis with EPS study and substrate ablation; unclear requirements for ECG pattern type	SCN5A, SCN10A	NR
Sacilotto et al (2020) ⁴¹ .	BrS patients in Brazilian registry with type-1 ECG pattern	Prospective; consecutive	Spontaneous or induced Brugada type 1 ECG pattern	SCN5A, GPD1L, SCN10A, SCN18, SCN28, SCN38, CACNA1C, CACNB2, KCND3, CACNAD2, KCNJ8, KCNE3, SLMAP, RANGRF	Unclear
Milman et al (2021) ⁴² .	BrS patients from 14 countries with a first arrhythmic event	Observational cohort (Survey on Arrhythmic Events in Brugada Syndrome); selection not reported	NR	SCN5A	None

BrS: Brugada syndrome; CPVT: catecholaminergic polymorphic ventricular tachycardia; ECG: electrocardiogram; EPS: electrophysiological study; LQTS: long QT syndrome; NR: not reported; SCD: sudden cardiac death.

Table 11. Yield of Genetic Testing for BrS

Study	N	Excluded Samples	Yield of Genetic Testing (Range)
Priori et al (2000) ³⁴ .	52	NR	15%
Bai et al (2009) ³¹ .	798		8%
Conclusive Dx	405		13%
Possible Dx	248		4%
Kaplinger et al (2010) ³⁵ .	2111	NR	21% (11% to 28%)
Hu et al (2014) ³⁶ .	150	NR	17%
Behr et al (2015) ³⁷ .	156	SCN5A re-sequencing (n=2) revision of the diagnosis (n=4), non-European ancestry (n=3)	11.5%
Andorin et al (2016) ³⁸ .	75 (from 62 families)	Only 75/106 have genetic analysis; reasons for lack of genetic analysis unclear	77%
Chen et al (2019) ³⁹ .	1780	NR	20% (11% to 43%)
Monasky et al (2019) ⁴⁰ .	294	NR	28.3% (SCN5A) 2.7% (SCN10A)
Sacilotto et al (2020) ⁴¹ .	138 (109 probands; 22/138 symptomatic)	Genetic analysis was only performed in 93/138 patients (76 asymptomatic, 17 symptomatic)	26.3% (SCN5A, asymptomatic) 29.4% (SCN5A, symptomatic)
Milman et al (2021) ⁴² .	678 (392 probands)	NR	23.5%

Dx: diagnosis; NR: not reported.

The purpose of limitations tables (see Tables 12 and 13) is to display notable limitations identified in selected primary studies. This information is synthesized as a summary of the body of evidence and provides the conclusions on the sufficiency of the evidence supporting the position statement.

Table 12. Study Relevance Limitations of Clinical Validity Studies of Genetic Testing for BrS

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Priori et al (2000) ³⁴ .	3: Criteria for clinical diagnosis unclear				
Bai et al (2009) ³¹ .	3: Criteria for clinical diagnosis unclear				
Kaplinger et al (2010) ³⁵ .	3: Criteria for clinical diagnosis unclear				
Hu et al (2014) ³⁶ .					
Behr et al (2015) ³⁷ .	3: Criteria for clinical diagnosis unclear				
Andorin et al (2016) ³⁸ .	4: Majority of probands had only Brugada pattern ECG without symptoms				
Monasky et al (2019) ⁴⁰ .	3: Criteria for clinical diagnosis unclear; patients had variety of type I-III ECG patterns			1: Study does not directly address a key health outcome	

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Sacilotto et al (2020) ⁴¹	4: Majority of probands had only Brugada type 1 ECG pattern without symptoms				
Milman et al (2021) ⁴²	3: Criteria for clinical diagnosis unclear			1: Study does not directly address a key health outcome	

BrS: Brugada syndrome; ECG: electrocardiogram.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

Table 13. Study Design and Conduct Limitations of Clinical Validity Studies of Genetic Testing for BrS

Study	Selection ^a	Blinding ^b	Test Delivery ^c	Selective Reporting ^d	Data Completeness ^e	Statistical
Priori et al (2000) ³⁴	1: Not clear if all eligible patients were included	1: Blinding not described			1: No description of exclusions or indeterminate results	
Bai et al (2009) ³¹		1: Blinding not described			1: No description of exclusions or indeterminate results	
Kapplinger et al (2010) ³⁵	1: Not clear if all eligible patients were included	1: Blinding not described			1: No description of exclusions or indeterminate results	
Hu et al (2014) ³⁶	1, 2: Not clear if all eligible patients were included; not clear how samples were selected	1: Blinding not described			1: No description of exclusions or indeterminate results	
Behr et al (2015) ³⁷	1, 2: Not clear if all eligible patients were included; not clear how samples were selected	1: Blinding not described				
Andorin et al (2016) ³⁸	1, 2: Not clear if all eligible patients were included; not clear how samples were selected	1: Blinding not described			1: Unclear why ~30% of patients did not have genetic analysis	

Study	Selection ^a	Blinding ^b	Test Delivery ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
Monasky et al (2019) ⁴⁰	1, 2: Not clear if all eligible patients were included; not clear how samples were selected	1: Blinding not described		1: Not registered; 2: Evidence of selective reporting; detailed outcomes for SCN5A cohort not reported		
Sacilotto et al (2020) ⁴¹		1: Blinding not described			1: Unclear why ~33% of patients did not have genetic analysis	
Milman et al (2021) ⁴²	1: Selection not described	1: Blinding not described				

BrS: Brugada syndrome.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

^b Blinding key: 1. Not blinded to results of reference or other comparator tests.

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

Catecholaminergic Polymorphic Ventricular Tachycardia

Studies reporting the yield of RyR2 testing in CPVT have been conducted in patients with clinically diagnosed CPVT.^{31,43,44,45} Characteristics are shown in Table 14 and results are shown in Table 15. The yield in cases with a 'strong' diagnosis of CPVT is around 60%.

Table 14. Characteristics of Clinical Validity Studies of Genetic Testing for CPVT

Study	Study Population	Design	Clinical Diagnosis	Genes included	Blinding of Assessors
Priori et al (2002) ⁴³	Patients with documented polymorphic ventricular arrhythmias occurring during physical or emotional stress with a normal heart	Retrospective; unclear whether samples were consecutive	Ventricular fibrillation elicited by physical or emotional stress in the absence of identifiable precipitating factors and in the absence of VT documented at Holter and/or exercise stress testing	RyR2	NR
Medeiros-Domingo et al (2009) ⁴⁴	Patients referred for genetic testing with "strong" diagnosis of CPVT	Retrospective; unclear whether samples were consecutive	Exertional syncope plus documentation of bidirectional or polymorphic VT	RyR2	NR
Bai et al (2009) ³¹	Patients from a sample of 1394 consecutive probands with either a clinically confirmed or	Consecutive; prospective	Diagnosed clinically as conclusive or possible; criteria not specified	RyR2	NR

Study	Study Population	Design	Clinical Diagnosis	Genes included	Blinding of Assessors
	suspected diagnosis of LQTS, BrS, or CPVT or a personal or family history of idiopathic ventricular fibrillation/cardiac arrest/SCD referred for molecular diagnosis				
Kapplinger et al (2018) ⁴⁵ .	Patients referred for commercial genetic testing with well-phenotyped cases and "strong" diagnosis of CPVT	Retrospective; unclear whether samples were consecutive	History of exertional syncope with documentation of exercise-related bidirectional or polymorphic VT	RyR2	NR

BrS: Brugada syndrome; CPVT: catecholaminergic polymorphic ventricular tachycardia; LQTS: long QT syndrome; NR: not reported; SCD: sudden cardiac death; VT: ventricular tachycardia.

Table 15. Yield of Genetic Testing for CPVT

Study	N	Excluded Samples	Yield of Genetic Testing
Priori et al (2002) ⁴³ .	30	NR	47%
Medeiros-Domingo et al (2009) ⁴⁴ .	78	NR	60%
Bai et al (2009) ³¹ .	175	NR	35%
Conclusive Dx	81	NR	62%
Possible Dx	21	NR	5%
Kapplinger et al (2018) ⁴⁵ .	78	NR	59%

CPVT: catecholaminergic polymorphic ventricular tachycardia; Dx: diagnosis; NR: not reported.

The purpose of the limitations tables (see Tables 16 and 17) is to display notable limitations identified in each study. This information is synthesized as a summary of the body of evidence and provides the conclusions on the sufficiency of the evidence supporting the position statement.

Table 16. Study Relevance Limitations of Clinical Validity Studies of Genetic Testing for CPVT

Study	Population ^a	Intervention ^b	Comparator ^c	Outcomes ^d	Duration of Follow-Up ^e
Priori et al (2002) ⁴³ .					
Medeiros-Domingo et al (2009) ⁴⁴ .					
Bai et al (2009) ³¹ .	3: Criteria for clinical diagnosis unclear				
Kapplinger et al (2018) ⁴⁵ .					

CPVT: catecholaminergic polymorphic ventricular tachycardia.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

^b Intervention key: 1. Classification thresholds not defined; 2. Version used unclear; 3. Not intervention of interest.

^c Comparator key: 1. Classification thresholds not defined; 2. Not compared to credible reference standard; 3. Not compared to other tests in use for same purpose.

^d Outcomes key: 1. Study does not directly assess a key health outcome; 2. Evidence chain or decision model not explicated; 3. Key clinical validity outcomes not reported (sensitivity, specificity, and predictive values); 4. Reclassification of diagnostic or risk categories not reported; 5. Adverse events of the test not described (excluding minor discomforts and inconvenience of venipuncture or noninvasive tests).

^e Follow-Up key: 1. Follow-up duration not sufficient with respect to natural history of disease (true-positives, true-negatives, false-positives, false-negatives cannot be determined).

Table 17. Study Design and Conduct Limitations of Clinical Validity Studies of Genetic Testing for CPVT

Study	Selection ^a	Blinding ^b	Test Delivery ^c	Selective Reporting ^d	Data Completeness ^e	Statistical ^f
Priori et al (2002) ⁴³ .	1,2: Not clear if all eligible patients were included	1: Blinding not described			1: No description of exclusions or indeterminate results	
Medeiros-Domingo et al (2009) ⁴⁴ .	1,2: Not clear if all eligible patients were included	1: Blinding not described			1: No description of exclusions or indeterminate results	
Bai et al (2009) ³¹ .		1: Blinding not described			1: No description of exclusions or indeterminate results	
Kapplinger et al (2018) ⁴⁵ .	1,2: Not clear if all eligible patients were included	1: Blinding not described			1: No description of exclusions or indeterminate results	

CPVT: catecholaminergic polymorphic ventricular tachycardia.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Selection key: 1. Selection not described; 2. Selection not random or consecutive (i.e., convenience).

^b Blinding key: 1. Not blinded to results of reference or other comparator tests.

^c Test Delivery key: 1. Timing of delivery of index or reference test not described; 2. Timing of index and comparator tests not same; 3. Procedure for interpreting tests not described; 4. Expertise of evaluators not described.

^d Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^e Data Completeness key: 1. Inadequate description of indeterminate and missing samples; 2. High number of samples excluded; 3. High loss to follow-up or missing data.

^f Statistical key: 1. Confidence intervals and/or p values not reported; 2. Comparison with other tests not reported.

The specificity of known pathogenic variants for CPVT is uncertain but is likely high. A 2013 publication from the National Heart, Lung, and Blood Institute ESP reported on sequence variants in a large number of patients without CPVT.⁴⁶ The ESP sequenced all genome regions of protein-coding in a sample of 6503 persons drawn from various populations who did not specifically have CPVT or other cardiac ion channelopathies. Exome data were systematically searched to identify missense variants previously associated with CPVT. Authors identified 11% previously described variants in the ESP population in 41 putative CPVT cases. These data suggested that false-positive results are low, but authors cautioned against attributing clinical CPVT to a single missense variant.

Short QT Syndrome

Limited data on the clinical validity of SQTs were identified in the peer-reviewed literature due to the rarity of the condition. A precise genetic testing yield is unknown.

Section Summary: Clinical Validity of Genetic Testing for the Diagnosis of a Specific Channelopathy

In probands with LQTS and CPVT, genetic testing has a yield for identifying a disease-causing variant of approximately 70% and 60%, respectively. In probands with BrS, genetic testing has a much lower yield probably ranging from about 15% to 30% depending on the genes included. The yield of genetic testing is not well established in SQTs.

Data on the clinical specificity are available for LQTS but there are limited data for CPVT. The specificity varies according to the type of variant identified. For LQTS nonsense variants, which have the highest rate of pathogenicity, there are very few false-positives among patients without LQTS, and therefore a high specificity. However, for missense variants, the rate is approximately 5% among patients without LQTS; therefore, the specificity for these types of variants is lower, and false-positive results do occur.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Long QT Syndrome

Long QT syndrome may lead to catastrophic outcomes (i.e., SCD) in otherwise healthy individuals. Diagnosis using clinical methods alone may lead to underdiagnosis of LQTS, thus exposing undiagnosed patients to the risk of SCA. For patients in whom the clinical diagnosis of LQTS is uncertain, genetic testing may be necessary to clarify whether LQTS is present. Patients who are identified as genetic carriers of LQTS variants have a non-negligible risk of adverse cardiac events even in the absence of clinical signs and symptoms of the disorder. Therefore, treatment is likely indicated for patients found to have an LQTS variant, with or without other signs or symptoms.

Treatment with β -blockers has been demonstrated to decrease the likelihood of cardiac events, including SCA.

Sodium-channel blockers (e.g., mexiletine) are sometimes used, particularly in those with SCN5A variants. Preliminary modeling studies by Zhu et al (2018) designed to predict LQT3 mutations with enhanced mexiletine sensitivity have been successfully validated in a small initial cohort of patients.⁴⁷

Treatment with an ICD is available for patients who fail or cannot take β -blockers.

Two studies evaluated the psychological effects of genetic testing for LQTS. Hendriks et al (2008) studied 77 patients with an LQTS variant and their 57 partners.⁴⁸ Psychologic testing was performed after the diagnosis of LQTS had been made and repeated twice over an 18-month period. Disease-related anxiety scores were increased in the index patients and their partners. This psychological distress decreased over time but remained elevated at 18 months.

Andersen et al (2008) conducted qualitative interviews with 7 individuals with LQTS variants.⁴⁹ They reported that affected patients had excess worry and limitations in daily life associated with the increased risk of sudden death, which was partially alleviated by acquiring knowledge about LQTS. The greatest concern was expressed for their family members, particularly children and grandchildren.

The evidence suggests that different LQTS subtypes may have variable prognoses, thus indicating that genetic testing may assist in risk stratification. Several reports have compared rates of cardiovascular events in subtypes of LQTS.^{50,51,52,53} These studies have reported that rates of cardiovascular events differ among subtypes, but there is no common pattern across all studies. Three of the 4 studies^{50,51,52} reported that patients with LQT2 have higher event rates than

patients with LQT1, while Zareba et al (1998)⁵³ reported that patients with LQT1 have higher event rates than patients with LQT2.

Some studies that have reported outcomes of treatment with β -blockers have also reported outcomes by specific subtypes of LQTS.^{50,52} Priori et al (2004) reported pre-post rates of cardiovascular events by LQTS subtypes following initiation of β -blocker therapy.⁵⁰ There was a decrease in event rates in all LQTS subtypes, with a similar magnitude of decrease in each subtype. Moss et al (2000) also reported pre-post event rates for patients treated with β -blocker therapy.⁵⁴ This study indicated a significant reduction in event rates for patients with LQT1 and LQT2 but not for LQT3. This analysis was limited by the small number of patients with LQT3 and cardiac events before β -blocker treatment (4/28). Sauer et al (2007) evaluated differential response to β -blocker therapy in a Cox proportional hazards analysis.⁵⁵ They reported an overall risk reduction in the first cardiac event of approximately 60% (hazard ratio [HR], 0.41; 95% CI, 0.27 to 0.64) in adults treated with β -blockers and an interaction effect by genotype. Efficacy of β -blocker treatment was worse in those with LQT3 genotype ($p=.04$) than in those with LQT1 or LQT2. There was no difference in efficacy between LQT1 and LQT2 genotypes.

Shimizu et al (2019) conducted an observational study on 1124 Japanese patients with LQTS and various pathogenic variants (e.g., nonpore membrane-spanning variants, pore site and segment 5 to segment 6 [S5-pore-S6] variants, and N/C-terminus variants) for LQT1, LQT2, and LQT3.⁵⁶ For patients with LQT1, the membrane-spanning pathogenic variant was associated with a higher risk of arrhythmic events compared to the N/C-terminus variant in female patients (HR, 1.60; 95% CI, 1.19 to 2.17; $p=.002$). Patients with LQT2 S5-pore-S6 variants were found to have a higher risk of arrhythmic events compared to others (HR, 1.88; 95% CI, 1.44 to 2.44; $p<.001$). In patients with LQT3, S5-pore-S6 variants were associated with lethal arrhythmic events compared with other (HR, 4.2; 95% CI, 2.09 to 8.36; $p<.001$). While these findings suggest that risk stratification of arrhythmic events may potentially be informed by specific pathogenic gene variants in LQTS, the study is limited by its retrospective analysis.

Biton et al (2019) studied LQTS patients ($n=212$) enrolled in the Rochester LQTS ICD registry who underwent ICD implantation for primary prevention of SCD.⁵⁷ During median follow-up duration of 9.2 ± 4.9 years, 42 patients experienced at least 1 appropriate shock. The cumulative probability of appropriate shock at 8 years was 22%. $QT_c \geq 550$ ms (HR, 3.94; 95% CI, 2.08 to 7.46; $p<.001$) and prior syncope on β -blockers (HR, 1.92, 95% CI, 1.01 to 3.65; $p=.047$) were associated with an increased risk of appropriate shock. Importantly, LQT2 genotype (HR, 2.10; 95% CI, 1.22 to 3.61; $p=.008$) and the presence of multiple mutations (HR, 2.87; 95% CI, 1.49 to 5.53; $p=.002$) were associated with an increased risk of recurrent shocks compared to LQT1 genotype, suggesting that both clinical and genetic variables may have utility in the risk stratification of high-risk patients undergoing evaluation for an ICD.

Cuneo et al (2020) conducted a multicenter retrospective analysis of 148 pregnancies from 103 families with the 3 most common heterozygous pathogenic LQTS genotypes (*KCNQ1*, *KCNH2*, or *SCN5A*).⁵⁸ Fetal death at >20 weeks gestation was 8 times more frequent compared to the general population. The likelihood of fetal death was found to be significantly greater with maternal versus paternal LQTS (24.4% vs. 3.5%; $p=.36$).

Brugada Syndrome

The diagnostic testing yield for BrS limits its clinical usefulness. A finding of a genetic variant is not diagnostic of the disorder but is an indicator of high risk for development of BrS. The diagnostic criteria for BrS do not presently include the presence of a genetic variant. Furthermore, treatment decisions are based on the presence of symptoms such as syncope or documented ventricular arrhythmias. Treatment is primarily with an implantable ICD, which is reserved for high-risk patients. However, for family members of patients with a known BrS variant, a negative test can rule out the disorder.

Rattanawong et al (2019) conducted a systematic review and meta-analysis of 7 cohort and case-control studies investigating the association of *SCN5A* mutations with major arrhythmic events (e.g., VT, ventricular fibrillation, appropriate implantable ICD shocks, aborted cardiac arrest, and SCD) in patients with BrS (n=1049).⁵⁹ *SCN5A* mutations were associated with major arrhythmic events in Asian patients (risk ratio, 2.03; 95% CI, 1.37 to 3.00; p=.0004; $I^2=0.0\%$), symptomatic patients (risk ratio, 2.66; 95% CI, 1.62 to 4.36; p=.0001; $I^2=23.0\%$), and patients with spontaneous Brugada type 1 ECG pattern (risk ratio, 1.84; 95% CI, 1.05 to 3.23; p=.03; $I^2=0.0\%$). The inclusion criteria did not specify criteria for establishing a clinical diagnosis of BrS, and therefore, the analysis was limited by heterogeneity in clinical, genetic, and outcome reporting among included studies. Reporting on specific major arrhythmic events relevant to health outcomes such as delivery of appropriate ICD shocks and aborted cardiac arrests was not individually reported. Therefore, the clinical utility of *SCN5A* genetic variant risk stratification in this population remains unclear.

Catecholaminergic Polymorphic Ventricular Tachycardia

The clinical utility for genetic testing in CPVT follows a similar chain of logic as that for LQTS. In patients for whom the clinical diagnosis can be made with certainty, there is a limited utility for genetic testing. However, there are some patients in whom signs and symptoms of CPVT are present, but for whom the diagnosis cannot be made with certainty. In this case, documentation of a pathogenic variant that is known to be associated with CPVT confirms the diagnosis. When the diagnosis is confirmed, treatment with β -blockers is indicated, and lifestyle changes are recommended. Although high-quality outcome studies are lacking to demonstrate a benefit of medication treatment, it is very likely that treatment reduces the risk of SCD. Therefore, there is a clinical utility.

There is currently no direct method of genotype-based risk stratification for management or prognosis of CPVT. However, testing can have important implications for all family members for presymptomatic diagnosis, counseling, or therapy. Asymptomatic patients with confirmed CPVT should also be treated with β -blockers and lifestyle changes. Also, CPVT has been associated with sudden infant death syndrome, and some investigators have considered testing at birth for prompt therapy in infants who are at-risk due to CPVT in close family members.

Short QT Syndrome

No studies were identified that provide evidence for the clinical utility of genetic testing for SQTs, consistent with the clinical rarity of the condition. Clinical sensitivity for the test is low, with laboratory test providers estimating a yield as low as 15%.

Section Summary: Clinical Utility of Genetic Testing for the Diagnosis of a Specific Channelopathy

The clinical utility of genetic testing for LQTS or CPVT is high when there is a moderate-to-high pretest probability and when the diagnosis cannot be made with certainty by other methods. A definitive diagnosis of either channelopathy leads to treatment with β -blockers in most cases, and sometimes to treatment with an ICD. As a result, confirming the diagnosis is likely to lead to a health outcome benefit by reducing the risk for ventricular arrhythmias and SCD. The clinical utility of testing is also high for close relatives of patients with known cardiac ion channel variants because these individuals should also be treated if they have a pathogenic variant.

For BrS, the clinical utility is less certain, but there is potential for genetic testing to change treatment decisions by stratifying patients for the need for ICD. A meta-analysis reported that the presence of *SCN5A* variants could not predict cardiac events; however, a registry study published after the meta-analysis reported that patients with the *SCN5A* variant experienced more cardiac events and experienced the first event at a younger age than patients who did not have the *SCN5A* variant. Studies have been conducted to further determine risk level by type of variant, but the studies have small sample sizes, so interpretation is limited.

For SQTs, the clinical utility is uncertain because there is no clear link between the establishment of a definitive diagnosis and a change in management that will improve outcomes.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Summary of Evidence**Long QT Syndrome**

For individuals with suspected congenital LQTS who receive genetic testing for variants associated with congenital LQTS, the evidence includes observational studies reporting on the testing yield. Relevant outcomes are OS, test validity, changes in reproductive decision making, and morbid events. A genetic variant can be identified in approximately 70% of those with LQTS. The clinical utility of genetic testing for LQTS is high when there is a moderate-to-high pretest probability. There is a chain of evidence to suggest that testing for variants associated with LQTS in individuals who are suspected to have these disorders leads to improved outcomes. A definitive diagnosis of LQTS leads to treatment with β -blockers in most cases, and sometimes to treatment with an ICD. As a result, confirming the diagnosis is likely to lead to a health outcome benefit by reducing the risk for ventricular arrhythmias and SCD. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with a close relative(s) with a known LQTS variant who receive genetic testing for variants associated with congenital LQTS, the evidence includes observational studies reporting on changes in management. Relevant outcomes are OS, test validity, changes in reproductive decision making, and morbid events. A positive genetic test for an LQTS variant leads to treatment with β -blockers in most cases, and sometimes to treatment with an ICD; a negative test would allow family members to defer further testing. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Brugada Syndrome

For individuals with suspected BrS who receive genetic testing for variants associated with BrS, the evidence includes observational studies reporting on testing yields. Relevant outcomes are OS, test validity, changes in reproductive decision making, and morbid events. The clinical validity of testing for BrS is low: a genetic variant can only be identified in approximately 15% to 35% of BrS. Management changes, primarily use of ICDs, are directed by clinical symptoms. It is not clear that a genetic diagnosis in the absence of other clinical signs and symptoms leads to a change in management that improves outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with a close relative(s) with a known BrS variant who receive genetic testing for variants associated with BrS, the evidence includes observational studies reporting on testing yields. Relevant outcomes are OS, test validity, changes in reproductive decision making, and morbid events. Brugada syndrome management changes, primarily use of ICDs, are directed by clinical symptoms. There is limited evidence on the effect of changes in management based on genetic testing in an individual with family members who have a known variant. However, a negative test would allow family members to defer further testing. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Catecholaminergic Polymorphic Ventricular Tachycardia

For individuals with suspected CPVT who receive genetic testing for variants associated with CPVT, the evidence includes observational studies reporting on testing yields. Relevant outcomes are OS, test validity, changes in reproductive decision making, and morbid events. A

genetic variant can be identified in approximately 60% of CPVT patients. There is a chain of evidence to suggest that testing for variants associated with CPVT in individuals who are suspected to have these disorders. Confirming the diagnosis of CPVT is likely to lead to a health outcome benefit by initiating changes in management that reduce the risk of ventricular arrhythmias and SCD. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with a close relative(s) with a known CPVT variant who receive genetic testing for variants associated with CPVT, the evidence includes observational studies reporting testing yields. Relevant outcomes are OS, test validity, changes in reproductive decision making, and morbid events. For close relatives of patients with known CPVT variants who are found to have a pathogenic variant, preventive treatment can be initiated. Also, a negative test in the setting of a known familial variant should have a high negative predictive value. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Short QT Syndrome

For individuals with suspected SQTS who receive genetic testing for variants associated with SQTS, the evidence includes limited data on testing yields. Relevant outcomes are OS, test validity, changes in reproductive decision making, and morbid events. The yield of genetic testing in SQTS is not well-characterized. Management changes, primarily use of ICDs, are directed by clinical symptoms. There is limited evidence on changes in management based on genetic testing in a symptomatic proband without a definitive diagnosis. It is not clear that a genetic diagnosis in the absence of other clinical signs and symptoms leads to a change in management that improves outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with a close relative(s) with a known SQTS variant who receive genetic testing for variants associated with SQTS, the evidence includes observational studies reporting on testing yields. Relevant outcomes are OS, test validity, changes in reproductive decision making, and morbid events. For patients with SQTS, management changes, primarily use of ICDs, are directed by clinical symptoms. There is limited evidence on changes in management based on genetic testing in an individual with family members who have a known variant. It is not clear that a genetic diagnosis in the absence of other clinical signs and symptoms leads to a change in management that improves outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with a close family member(s) who experienced SCD and a specific diagnosis has been made who receive genetic testing for variants associated with cardiac ion channelopathies, the evidence includes cohort studies that describe the genetic testing yield. In all studies identified, genetic testing was obtained only after a specific diagnosis was suspected based on history or ancillary testing. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2015 Input

In response to requests from Blue Cross Blue Shield Association, input was received from 3 specialty societies (4 reviewers) and 4 academic medical centers (9 reviewers) in 2015. Input was limited to the use of genetic testing for Brugada syndrome (BrS) and short QT syndrome (SQTS). There was a consensus that genetic testing for BrS is medically necessary to establish the diagnosis of BrS in an individual with a suspected but not definitive diagnosis of BrS and to evaluate family members of an individual with a known pathogenic genetic variant for BrS. There was less consensus on whether genetic testing for variants associated with SQTS is medically necessary to establish the diagnosis of SQTS in an individual with a suspected but not definitive diagnosis of SQTS, but there was consensus that testing for SQTS to evaluate family members of an individual with a known pathogenic genetic variant for SQTS is medically necessary. However, reviewers acknowledged that the rarity of SQTS somewhat limited conclusions that could be made.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in Supplemental Information if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American Heart Association

In 2021, the American Heart Association published a scientific statement on genetic testing for heritable cardiovascular diseases (including channelopathies) in children.⁶⁰ The statement recommends that genetic testing be performed when a cardiac channelopathy is likely to be present, including after a variant has been found in a family member. Testing to identify at-risk relatives can be considered. Brugada syndrome is difficult to identify since not all adults express genetic variants; therefore, identifying at-risk children may require clinical evaluation, electrocardiogram (ECG) testing, and/or pharmacologic challenge of all of the child's first-degree relatives. Genetic testing should also be performed in children who are resuscitated from cardiac arrest with no clear cause. Several factors can be considered when deciding the appropriate age for genetic testing of an individual child, including whether the disease is expected to present during childhood, whether the channelopathy can be fatal, whether therapies exist to mitigate mortality risk, and family preferences. Ongoing follow-up genetic testing can confirm pathogenicity of the variant over time.

In 2020, the American Heart Association authored a scientific statement on genetic testing for inherited cardiovascular disease.⁶¹ Prior guidelines from several international cardiovascular clinical organizations and published studies were reviewed. For BrS, the authors concluded that genetic testing supports the clinical diagnosis. For patients with catecholaminergic polymorphic ventricular tachycardia (CPVT) and long QT syndrome (LQTS), genetic testing is needed for diagnosis and subtype classification. Management of LQTS may also differ depending on the causative gene. Genetic testing for all of these conditions facilitates identifying at-risk family members. Specific genes with the strongest causative evidence for cardiac channelopathies are listed in Table 18.

Table 18. Specific Genes for Testing in Cardiac Channelopathies

Channelopathy	Genes with definitive evidence of a causal role in the disease
LQTS	<i>KCNQ1, KCNH2, SCN5A</i>
SQTS	<i>KCNH2, KCNQ1, KCNJ2</i>
BrS	<i>SCN5A</i>
CPVT	<i>RYR2, CASQ2</i>

BrS: Brugada syndrome; CPVT: catecholaminergic polymorphic ventricular tachycardia; LQTS: long QT syndrome; SQTS: short QT syndrome.

American Heart Association, American College of Cardiology, and Heart Rhythm Society

In 2017, the American Heart Association, American College of Cardiology, and the Heart Rhythm Society published guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death (SCD).⁶² Table 1 9 summarizes the recommendations relating to cardiac ion channelopathies.

Table 1 9. Recommendations for Genetic Testing in Cardiac Channelopathies

Consensus Recommendation	COR	LOE
In first-degree relatives of patients who have a causative mutation for LQTS, CPVT, SQTS, or BrS, genetic counseling and mutation-specific genetic testing are recommended.	I (strong)	B-NR
In patients with clinically diagnosed LQTS, genetic counseling and genetic testing are recommended. Genetic testing offers diagnostic, prognostic, and therapeutic information.	I (strong)	B-NR
In patients with CPVT and with clinical VT or exertional syncope, genetic counseling and genetic testing are reasonable. Genetic testing may confirm a diagnosis; however, therapy for these patients is not guided by genotype status.	IIa (moderate)	B-NR
In patients with suspected or established BrS, genetic counseling and genetic testing may be useful to facilitate cascade screening of relatives, allowing for lifestyle modification and potential treatment.	IIb (weak)	C-EO
In patients with SQTS, genetic testing may be considered to facilitate screening of first-degree relatives.	IIb (weak)	C-EO

B-NR: moderate level of evidence, nonrandomized studies; BrS: Brugada syndrome; C-EO: consensus of expert opinion based on clinical experience; COR: class of recommendation; CPVT: catecholaminergic polymorphic ventricular tachycardia; LOE: level of evidence; LQTS: long QT syndrome; SQTS: short QT syndrome; VT: ventricular tachycardia.

Heart Rhythm Society and Asia Pacific Heart Rhythm Society

In 2020, the Heart Rhythm Society and Asia Pacific Heart Rhythm Society authored an expert consensus statement on investigation of individuals who have died from sudden unexplained death, patients with sudden cardiac arrest (SCA), and their families.⁶³ Suspicion for a genetic cause of SCD or a resuscitated SCA warrants genetic testing and counseling. Genetic testing should include the most likely genes for the suspected phenotype and should include clinical and genetic evaluation of family members to identify other at-risk individuals. Testing of many genes can lead to uncertainty and misinterpretation of results and is generally discouraged. Genetic investigation should only be undertaken by multidisciplinary teams with expertise in cardiology, genetics, and pathology. The document provides detailed guidance on specific scenarios for which genetic testing is warranted but does not describe specific genes that should be tested.

Heart Rhythm Society, European Heart Rhythm Association, and Asia Pacific Heart Rhythm Society

In 2013, the Heart Rhythm Society, the European Heart Rhythm Association, and the Asia Pacific Heart Rhythm Society issued an expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes.⁶⁴ The consensus statement refers to the 2011 guidelines on genetic testing for channelopathies and cardiomyopathies discussed next for the indications for genetic testing in patients affected by inherited arrhythmias and their family members and for diagnostic, prognostic, and therapeutic implications of the results of genetic testing. The 2013 consensus statement provided guidance for the evaluation of patients with idiopathic ventricular fibrillation, sudden unexplained death syndrome, and sudden unexplained death in infancy. Guidance on genetic testing for these patients was included (see Table 20). Idiopathic ventricular fibrillation is defined as a resuscitated cardiac arrest victim, preferably with documentation of ventricular fibrillation, in whom known cardiac, respiratory, metabolic, and toxicologic etiologies have been excluded through clinical evaluation.

The guidelines defined several terms related to specific types of SCD, including sudden unexplained death syndrome, which refers to an unexplained sudden death in an individual older than 1 year of age, sudden arrhythmic death syndrome, which refers to a sudden

unexplained death syndrome case with negative pathologic and toxicologic assessment, and sudden unexplained death in infancy, which refers to an unexplained sudden death in an individual younger than 1 year of age with negative pathologic and toxicologic assessment.

Table 20. Recommendations for Genetic Testing in IVF, SUDS, and SUDI

	Consensus Recommendation	Class
IVF	Genetic testing in IVF can be useful when there is suspicion of a specific genetic disease following clinical evaluation of the IVF patient and/or family members.	IIa
	Genetic screening of a large panel of genes in IVF patients in whom there is no suspicion of an inherited arrhythmogenic disease after clinical evaluation should not be performed.	III
SUDS	Collection of blood and/or suitable tissue for molecular autopsy/postmortem genetic testing is recommended in all SUDS victims.	I
	Genetic screening of the first-degree relatives of a SUDS victim is recommended whenever a pathogenic mutation in a gene associated with increased risk of sudden death is identified by molecular autopsy in the SUDS victim.	I
SUDI	Collection of blood and/or suitable tissue for molecular autopsy is recommended in all SUDI victims.	I
	An arrhythmia syndrome-focused molecular autopsy/postmortem genetic testing can be useful for all SUDI victims.	IIa
	Genetic screening of the first-degree relatives of a SUDI victim is recommended whenever a pathogenic mutation in a gene associated with increased risk of sudden death is identified by molecular autopsy in the SUDI victim. Obligate mutations carriers should be prioritized.	I

IVF: idiopathic ventricular fibrillation; SUDI: sudden unexplained death in infancy; SUDS: sudden unexplained death syndrome.

In 2011, the Heart Rhythm Society and European Heart Rhythm Association jointly published an expert consensus statement on genetic testing for channelopathies and cardiomyopathies.²⁴ This document made the following specific recommendations on testing for LQTS, BrS, CPVT, and SQTS (see Table 2 1).

Table 2 1. Cardiac Ion Channelopathy Testing Recommendations

	Consensus Recommendation	Class ^a	LOE ^b
LQTS	<ul style="list-style-type: none"> Comprehensive or LQT1-3 (<i>KCNQ1</i>, <i>KCNH2</i>, <i>SCN5A</i>) targeted LQTS genetic testing is recommended for any patient in whom a cardiologist has established a strong clinical index of suspicion for LQTS based on examination of the patient's clinical history, family history, and expressed electrocardiographic (resting 12-lead ECGs and/or provocative stress testing with exercise or catecholamine infusion) phenotype. Comprehensive or LQT1-3 (<i>KCNQ1</i>, <i>KCNH2</i>, <i>SCN5A</i>) targeted LQTS genetic testing is recommended for any asymptomatic patient with QT prolongation in the absence of other clinical conditions that might prolong the QT interval (such as electrolyte abnormalities, hypertrophy, bundle branch block, etc., i.e., otherwise idiopathic) on serial 12-lead ECGs defined as QTc >480 ms (prepuberty) or >500 ms (adults). Mutation-specific genetic testing is recommended for family members and other appropriate relatives subsequently following the identification of the LQTS-causative mutation in an index case. 	I	C
	Comprehensive or LQT1-3 (<i>KCNQ1</i> , <i>KCNH2</i> , <i>SCN5A</i>) targeted LQTS genetic testing may be considered for any asymptomatic patient with otherwise idiopathic QTc values >460 ms (prepuberty) or >480 ms (adults) on serial 12-lead ECGs.	IIb	C
BrS	Mutation-specific genetic testing is recommended for family members and appropriate relatives following the identification of the BrS-causative mutation in an index case.	I	C
	Comprehensive or BrS1 (<i>SCN5A</i>) targeted BrS genetic testing can be useful for any patient in whom a cardiologist has established a clinical index of	IIa	C

	Consensus Recommendation	Class ^a	LOE ^b
	suspicion for BrS based on examination of the patient's clinical history, family history, and expressed electrocardiographic (resting 12-lead ECGs and/or provocative drug challenge testing) phenotype.		
	Genetic testing is not indicated in the setting of an isolated type 2 or type 3 Brugada ECG pattern.	III	C
CPVT	Comprehensive or <i>CPVT1</i> and <i>CVPT2</i> (<i>RYR2</i> , <i>CASQ2</i>) targeted CPVT genetic testing is recommended for any patient in whom a cardiologist has established a clinical index of suspicion for CPVT based on examination of the patient's clinical history, family history, and expressed electrocardiographic phenotype during provocative stress testing with cycle, treadmill, or catecholamine infusion. Mutation-specific genetic testing is recommended for family members and appropriate relatives following the identification of the CPVT-causative mutation in an index case.	I	C
SQTS	Mutation-specific genetic testing is recommended for family members and appropriate relatives following the identification of the SQTS-causative mutation in an index case.	I	C
	Comprehensive or SQT1-3 (<i>KCNH2</i> , <i>KCNQ1</i> , <i>KCNJ2</i>) targeted SQTS genetic testing may be considered for any patient in whom a cardiologist has established a strong clinical index of suspicion for SQTS based on examination of the patient's clinical history, family history, and electrocardiographic phenotype.	IIb	C

BrS: Brugada syndrome; CPVT: catecholaminergic polymorphic ventricular tachycardia; ECG: electrocardiogram; LOE: level of evidence; LQTS: long QT syndrome; QTc: corrected QT; SQTS: short QT syndrome.

^a Class I: "is recommended" when an index case has a sound clinical suspicion for the presence of a channelopathy with a high positive predictive value for the genetic test (>40%) with a signal-to-noise ratio of >10 and/or the test may provide diagnostic or prognostic information or may change therapeutic choices; Class IIa: "can be useful"; Class IIb: "may be considered"; Class III: "is not recommended" (the test fails to provide any additional benefit or could be harmful in the diagnostic process).

^b Only consensus opinion of experts, case studies or standard of care.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 22.

Table 22. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
Ongoing			
NCT04832126	Genetic Analysis of Heart Channelopathies in Brazilian Patients and Their Relatives	100	Jul 2024 (recruiting)
NCT03783975	A Community-Based Approach to Overcoming Barriers to Cascade Screening for Long QT Syndrome	500	Dec 2021 (recruiting)
NCT02439658	Genetics of QT Prolongation With Antiarrhythmics (DOFEGEN)	1000	Jul 2022 (recruiting)
NCT04232787	Discovering the Genetic Causes of Brugada Syndrome in Thais and Southeast Asian Population (SEA-BrS)	750	Jan 2023 (recruiting)
NCT02824822	Genetic Markers of Cardiovascular Diseases and the Potential Role in Sudden Unexpected Death in Epilepsy	600	Dec 2023 (recruiting)
NCT02014961	Worm Study: Identification of Modifier Genes in a Unique Founder Population With Sudden Cardiac Death	223	Apr 2025 (recruiting)
NCT03880708	China Inherited Ventricular Arrhythmias Registry, a Multicenter, Observational and Prospective Study	500	Oct 2027 (recruiting)
Unpublished			

NCT No.	Trial Name	Planned Enrollment	Completion Date
NCT01705925 ^a	Multicenter Evaluation of Children and Young Adults With Genotype Positive Long QT Syndrome	92	Dec 2018 (completed)
NCT02425189	The Canadian National Long QT Syndrome Registry (LOTSREG)	1051	Aug 2020 (completed)

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Appendix 1

Appendix Table 1. Categories of Genetic Testing Addressed in Policy No. 2.04.43

Category	Addressed
1. Testing of an affected individual's germline to benefit the individual	
1a. Diagnostic	X
1b. Prognostic	X
1c. Therapeutic	
2. Testing cancer cells from an affected individual to benefit the individual	
2a. Diagnostic	
2b. Prognostic	
2c. Therapeutic	
3. Testing an asymptomatic individual to determine future risk of disease	X
4. Testing of an affected individual's germline to benefit family members	
5. Reproductive testing	
5a. Carrier testing: preconception	
5b. Carrier testing: prenatal	
5c. In utero testing: aneuploidy	
5d. In utero testing: familial variants	
5e. In utero testing: other	
5f. Preimplantation testing with in vitro fertilization	

References

- Abriel H, Zaklyazminskaya EV. Cardiac channelopathies: genetic and molecular mechanisms. *Gene*. Mar 15 2013; 517(1): 1-11. PMID 23266818
- Modell SM, Bradley DJ, Lehmann MH. Genetic testing for long QT syndrome and the category of cardiac ion channelopathies. *PLoS Curr*. May 03 2012: e4f9995f69e6c7. PMID 22872816
- Huang MH, Marcus FI. Idiopathic Brugada-type electrocardiographic pattern in an octogenarian. *J Electrocardiol*. Apr 2004; 37(2): 109-11. PMID 15127377
- Brugada R, Campuzano O, Sarquella-Brugada G, et al. Brugada Syndrome. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2016.
- Tester DJ, Ackerman MJ. Genetic testing for potentially lethal, highly treatable inherited cardiomyopathies/channelopathies in clinical practice. *Circulation*. Mar 08 2011; 123(9): 1021-37. PMID 21382904
- Bennett MT, Sanatani S, Chakrabarti S, et al. Assessment of genetic causes of cardiac arrest. *Can J Cardiol*. Jan 2013; 29(1): 100-10. PMID 23200097
- Ackerman MJ, Marcou CA, Tester DJ. Personalized medicine: genetic diagnosis for inherited cardiomyopathies/channelopathies. *Rev Esp Cardiol*. Apr 2013;66(4):298-307. PMID 23484907
- Wilders R. Cardiac ion channelopathies and the sudden infant death syndrome. *ISRN Cardiol*. 2012; 2012: 846171. PMID 23304551
- Eddy CA, MacCormick JM, Chung SK, et al. Identification of large gene deletions and duplications in KCNQ1 and KCNH2 in patients with long QT syndrome. *Heart Rhythm*. Sep 2008; 5(9): 1275-81. PMID 18774102
- Chiang CE. Congenital and acquired long QT syndrome. Current concepts and management. *Cardiol Rev*. Jul-Aug 2004; 12(4): 222-34. PMID 15191637

11. Priori SG, Napolitano C, Schwartz PJ. Low penetrance in the long-QT syndrome: clinical impact. *Circulation*. Feb 02 1999; 99(4): 529-33. PMID 9927399
12. Sarquella-Brugada G, Fernandez-Falgueras A, Cesar S, et al. Clinical impact of rare variants associated with inherited channelopathies: a 5-year update. *Hum Genet*. Sep 21 2021. PMID 34546463
13. Beckmann BM, Scheiper-Welling S, Wilde AAM, et al. Clinical utility gene card for: Long-QT syndrome. *Eur J Hum Genet*. Dec 2021; 29(12): 1825-1832. PMID 34031550
14. Arking DE, Pulit SL, Crotti L, et al. Genetic association study of QT interval highlights role for calcium signaling pathways in myocardial repolarization. *Nat Genet*. Aug 2014; 46(8): 826-36. PMID 24952745
15. Alders M, Christiaans I. Long QT Syndrome. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2015.
16. Walsh R, Adler A, Amin AS, et al. Evaluation of gene validity for CPVT and short QT syndrome in sudden arrhythmic death. *Eur Heart J*. Sep 24 2021. PMID 34557911
17. Napolitano C, Priori SG, Bloise R. Catecholaminergic Polymorphic Ventricular Tachycardia. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2016.
18. Schwartz PJ, Moss AJ, Vincent GM, et al. Diagnostic criteria for the long QT syndrome. An update. *Circulation*. Aug 1993; 88(2): 782-4. PMID 8339437
19. Perrin MJ, Gollob MH. The genetics of cardiac disease associated with sudden cardiac death: a paper from the 2011 William Beaumont Hospital Symposium on molecular pathology. *J Mol Diagn*. Sep 2012; 14(5): 424-36. PMID 22749884
20. Wilde AA, Behr ER. Genetic testing for inherited cardiac disease. *Nat Rev Cardiol*. Oct 2013; 10(10): 571-83. PMID 23900354
21. Antzelevitch C, Brugada P, Borggrefe M, et al. Brugada syndrome: report of the second consensus conference: endorsed by the Heart Rhythm Society and the European Heart Rhythm Association. *Circulation*. Feb 08 2005; 111(5): 659-70. PMID 15655131
22. Benito B, Brugada J, Brugada R, et al. Brugada syndrome. *Rev Esp Cardiol*. Nov 2009; 62(11): 1297-315. PMID 19889341
23. Sumitomo N, Harada K, Nagashima M, et al. Catecholaminergic polymorphic ventricular tachycardia: electrocardiographic characteristics and optimal therapeutic strategies to prevent sudden death. *Heart*. Jan 2003; 89(1): 66-70. PMID 12482795
24. Ackerman MJ, Priori SG, Willems S, et al. HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies this document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Heart Rhythm*. Aug 2011; 8(8): 1308-39. PMID 21787999
25. Tristani-Firouzi M. The Long and Short of It: Insights Into the Short QT Syndrome. *J Am Coll Cardiol*. Apr 08 2014; 63(13): 1309-1310. PMID 24333498
26. Giustetto C, Di Monte F, Wolpert C, et al. Short QT syndrome: clinical findings and diagnostic-therapeutic implications. *Eur Heart J*. Oct 2006; 27(20): 2440-7. PMID 16926178
27. Gollob MH, Redpath CJ, Roberts JD. The short QT syndrome: proposed diagnostic criteria. *J Am Coll Cardiol*. Feb 15 2011; 57(7): 802-12. PMID 21310316
28. Asatryan B, Schaller A, Seiler J, et al. Usefulness of Genetic Testing in Sudden Cardiac Arrest Survivors With or Without Previous Clinical Evidence of Heart Disease. *Am J Cardiol*. Jun 15 2019; 123(12): 2031-2038. PMID 30975432
29. Chiu SN, Juang JJ, Tseng WC, et al. Impact of genetic tests on survivors of paediatric sudden cardiac arrest. *Arch Dis Child*. Jun 14 2021. PMID 34127479
30. Tester DJ, Will ML, Haglund CM, et al. Effect of clinical phenotype on yield of long QT syndrome genetic testing. *J Am Coll Cardiol*. Feb 21 2006; 47(4): 764-8. PMID 16487842
31. Bai R, Napolitano C, Bloise R, et al. Yield of genetic screening in inherited cardiac channelopathies: how to prioritize access to genetic testing. *Circ Arrhythm Electrophysiol*. Feb 2009; 2(1): 6-15. PMID 19808439
32. Kapa S, Tester DJ, Salisbury BA, et al. Genetic testing for long-QT syndrome: distinguishing pathogenic mutations from benign variants. *Circulation*. Nov 03 2009; 120(18): 1752-60. PMID 19841300

33. Refsgaard L, Holst AG, Sadjadieh G, et al. High prevalence of genetic variants previously associated with LQT syndrome in new exome data. *Eur J Hum Genet*. Aug 2012; 20(8): 905-8. PMID 22378279
34. Priori SG, Napolitano C, Gasparini M, et al. Clinical and genetic heterogeneity of right bundle branch block and ST-segment elevation syndrome: A prospective evaluation of 52 families. *Circulation*. Nov 14 2000; 102(20): 2509-15. PMID 11076825
35. Kapplinger JD, Tester DJ, Alders M, et al. An international compendium of mutations in the SCN5A-encoded cardiac sodium channel in patients referred for Brugada syndrome genetic testing. *Heart Rhythm*. Jan 2010; 7(1): 33-46. PMID 20129283
36. Hu D, Barajas-Martinez H, Pfeiffer R, et al. Mutations in SCN10A are responsible for a large fraction of cases of Brugada syndrome. *J Am Coll Cardiol*. Jul 08 2014; 64(1): 66-79. PMID 24998131
37. Behr ER, Savio-Galimberti E, Barc J, et al. Role of common and rare variants in SCN10A: results from the Brugada syndrome QRS locus gene discovery collaborative study. *Cardiovasc Res*. Jun 01 2015; 106(3): 520-9. PMID 25691538
38. Andorin A, Behr ER, Denjoy I, et al. Impact of clinical and genetic findings on the management of young patients with Brugada syndrome. *Heart Rhythm*. Jun 2016; 13(6): 1274-82. PMID 26921764
39. Chen C, Tan Z, Zhu W, et al. Brugada syndrome with SCN5A mutations exhibits more pronounced electrophysiological defects and more severe prognosis: A meta-analysis. *Clin Genet*. Jan 2020; 97(1): 198-208. PMID 30963536
40. Monasky MM, Micaglio E, Vicedomini G, et al. Comparable clinical characteristics in Brugada syndrome patients harboring SCN5A or novel SCN10A variants. *Europace*. Oct 01 2019; 21(10): 1550-1558. PMID 31292628
41. Sacilotto L, Scanavacca MI, Olivetti N, et al. Low rate of life-threatening events and limitations in predicting invasive and noninvasive markers of symptoms in a cohort of type 1 Brugada syndrome patients: Data and insights from the GenBra registry. *J Cardiovasc Electrophysiol*. Nov 2020; 31(11): 2920-2928. PMID 32870538
42. Milman A, Behr ER, Gray B, et al. Genotype-Phenotype Correlation of SCN5A Genotype in Patients With Brugada Syndrome and Arrhythmic Events: Insights From the SABRUS in 392 Proband. *Circ Genom Precis Med*. Oct 2021; 14(5): e003222. PMID 34461752
43. Priori SG, Napolitano C, Memmi M, et al. Clinical and molecular characterization of patients with catecholaminergic polymorphic ventricular tachycardia. *Circulation*. Jul 02 2002; 106(1): 69-74. PMID 12093772
44. Medeiros-Domingo A, Bhuiyan ZA, Tester DJ, et al. The RYR2-encoded ryanodine receptor/calcium release channel in patients diagnosed previously with either catecholaminergic polymorphic ventricular tachycardia or genotype negative, exercise-induced long QT syndrome: a comprehensive open reading frame mutational analysis. *J Am Coll Cardiol*. Nov 24 2009; 54(22): 2065-74. PMID 19926015
45. Kapplinger JD, Pundi KN, Larson NB, et al. Yield of the RYR2 Genetic Test in Suspected Catecholaminergic Polymorphic Ventricular Tachycardia and Implications for Test Interpretation. *Circ Genom Precis Med*. Feb 2018; 11(2): e001424. PMID 29453246
46. Jabbari J, Jabbari R, Nielsen MW, et al. New exome data question the pathogenicity of genetic variants previously associated with catecholaminergic polymorphic ventricular tachycardia. *Circ Cardiovasc Genet*. Oct 2013; 6(5): 481-9. PMID 24025405
47. Zhu W, Mazzanti A, Voelker TL, et al. Predicting Patient Response to the Antiarrhythmic Mexiletine Based on Genetic Variation. *Circ Res*. Feb 15 2019; 124(4): 539-552. PMID 30566038
48. Hendriks KS, Hendriks MM, Birnie E, et al. Familial disease with a risk of sudden death: a longitudinal study of the psychological consequences of predictive testing for long QT syndrome. *Heart Rhythm*. May 2008; 5(5): 719-24. PMID 18452877
49. Andersen J, Oyen N, Bjorvatn C, et al. Living with long QT syndrome: a qualitative study of coping with increased risk of sudden cardiac death. *J Genet Couns*. Oct 2008; 17(5): 489-98. PMID 18719982

50. Priori SG, Napolitano C, Schwartz PJ, et al. Association of long QT syndrome loci and cardiac events among patients treated with beta-blockers. *JAMA*. Sep 15 2004; 292(11): 1341-4. PMID 15367556
51. Priori SG, Schwartz PJ, Napolitano C, et al. Risk stratification in the long-QT syndrome. *N Engl J Med*. May 08 2003; 348(19): 1866-74. PMID 12736279
52. Schwartz PJ, Priori SG, Spazzolini C, et al. Genotype-phenotype correlation in the long-QT syndrome: gene-specific triggers for life-threatening arrhythmias. *Circulation*. Jan 02 2001; 103(1): 89-95. PMID 11136691
53. Zareba W, Moss AJ, Schwartz PJ, et al. Influence of the genotype on the clinical course of the long-QT syndrome. International Long-QT Syndrome Registry Research Group. *N Engl J Med*. Oct 01 1998; 339(14): 960-5. PMID 9753711
54. Moss AJ, Zareba W, Hall WJ, et al. Effectiveness and limitations of beta-blocker therapy in congenital long-QT syndrome. *Circulation*. Feb 15 2000; 101(6): 616-23. PMID 10673253
55. Sauer AJ, Moss AJ, McNitt S, et al. Long QT syndrome in adults. *J Am Coll Cardiol*. Jan 23 2007; 49(3): 329-37. PMID 17239714
56. Shimizu W, Makimoto H, Yamagata K, et al. Association of Genetic and Clinical Aspects of Congenital Long QT Syndrome With Life-Threatening Arrhythmias in Japanese Patients. *JAMA Cardiol*. Mar 01 2019; 4(3): 246-254. PMID 30758498
57. Biton Y, Rosero S, Moss AJ, et al. Primary prevention with the implantable cardioverter-defibrillator in high-risk long-QT syndrome patients. *Europace*. Feb 01 2019; 21(2): 339-346. PMID 29947754
58. Cuneo BF, Kaizer AM, Clur SA, et al. Mothers with long QT syndrome are at increased risk for fetal death: findings from a multicenter international study. *Am J Obstet Gynecol*. Mar 2020; 222(3): 263.e1-263.e11. PMID 31520628
59. Rattanawong P, Chenbhanich J, Mekraksakit P, et al. SCN5A mutation status increases the risk of major arrhythmic events in Asian populations with Brugada syndrome: systematic review and meta-analysis. *Ann Noninvasive Electrocardiol*. Jan 2019; 24(1): e12589. PMID 30126015
60. Landstrom AP, Kim JJ, Gelb BD, et al. Genetic Testing for Heritable Cardiovascular Diseases in Pediatric Patients: A Scientific Statement From the American Heart Association. *Circ Genom Precis Med*. Oct 2021; 14(5): e000086. PMID 34412507
61. Musunuru K, Hershberger RE, Day SM, et al. Genetic Testing for Inherited Cardiovascular Diseases: A Scientific Statement From the American Heart Association. *Circ Genom Precis Med*. Aug 2020; 13(4): e000067. PMID 32698598
62. Al-Khatib SM, Stevenson WG, Ackerman MJ, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: Executive summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *Heart Rhythm*. Oct 2018; 15(10): e190-e252. PMID 29097320
63. Stiles MK, Wilde AAM, Abrams DJ, et al. 2020 APHRS/HRS expert consensus statement on the investigation of decedents with sudden unexplained death and patients with sudden cardiac arrest, and of their families. *Heart Rhythm*. Jan 2021; 18(1): e1-e50. PMID 33091602
64. Priori SG, Wilde AA, Horie M, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes: document endorsed by HRS, EHRA, and APHRS in May 2013 and by ACCF, AHA, PACES, and AEPC in June 2013. *Heart Rhythm*. Dec 2013; 10(12): 1932-63. PMID 24011539
65. Blue Cross Blue Shield Association. Medical Policy Reference Manual, No. 2.04.43 (January 2022).

Documentation for Clinical Review

Please provide the following documentation:

- History and physical and/or cardiology consultation notes including:
 - Syndrome that is being tested and clinical justification for testing
 - Description of signs and symptoms

- o Schwartz score (if applicable)
- o Cardiac testing results (e.g., electrocardiogram, holter or event monitor report)
- o Family history specific to long QT syndrome, Brugada syndrome, Catecholaminergic Polymorphic Ventricular Tachycardia, or short QT syndrome, including relationship to patient

Post Service (in addition to the above, please include the following):

- Results/reports of tests performed
- Procedure report(s)

Coding

This Policy relates only to the services or supplies described herein. Benefits may vary according to product design; therefore, contract language should be reviewed before applying the terms of the Policy.

The following codes are included below for informational purposes. Inclusion or exclusion of a code(s) does not constitute or imply member coverage or provider reimbursement policy. Policy Statements are intended to provide member coverage information and may include the use of some codes for clarity. The Policy Guidelines section may also provide additional information for how to interpret the Policy Statements and to provide coding guidance in some cases.

Type	Code	Description
CPT®	0237U	Cardiac ion channelopathies (e.g., Brugada syndrome, long QT syndrome, short QT syndrome, catecholaminergic polymorphic ventricular tachycardia), genomic sequence analysis panel including ANK2, CASQ2, CAV3, KCNE1, KCNE2, KCNH2, KCNJ2, KCNQ1, RYR2, and SCN5A, including small sequence changes in exonic and intronic regions, deletions, duplications, mobile element insertions, and variants in non-uniquely mappable regions
	81403	Molecular Pathology Procedure Level 4
	81404	Molecular Pathology Procedure Level 5
	81405	Molecular Pathology Procedure Level 6
	81406	Molecular Pathology Procedure Level 7
	81407	Molecular Pathology Procedure Level 8
	81408	Molecular Pathology Procedure Level 9
	81413	Cardiac ion channelopathies (e.g., Brugada syndrome, long QT syndrome, short QT syndrome, catecholaminergic polymorphic ventricular tachycardia); genomic sequence analysis panel, must include sequencing of at least 10 genes, including ANK2, CASQ2, CAV3, KCNE1, KCNE2, KCNH2, KCNJ2, KCNQ1, RYR2, and SCN5A
81414	Cardiac ion channelopathies (e.g., Brugada syndrome, long QT syndrome, short QT syndrome, catecholaminergic polymorphic ventricular tachycardia); duplication/deletion gene analysis panel, must include analysis of at least 2 genes, including KCNH2 and KCNQ1	
HCPCS	S3861	Genetic testing, sodium channel, voltage-gated, type V, alpha subunit (SCN5A) and variants for suspected Brugada Syndrome

Policy History

This section provides a chronological history of the activities, updates and changes that have occurred with this Medical Policy.

Effective Date	Action
07/06/2012	New policy
04/04/2014	Policy title change from Genetic Testing for Congenital Long QT Syndrome. Policy revision with position change.
06/30/2015	Coding update
01/01/2016	Policy revision with position change
03/01/2017	Policy revision without position change
03/01/2018	Policy revision without position change
04/01/2019	Policy revision without position change
03/01/2020	Annual review. No change to policy statement. Literature review updated.
01/01/2021	Coding Update
03/01/2021	Annual review. No change to policy statement. Literature review updated.
03/01/2022	Annual review. No change to policy statement. Literature review updated.

Definitions of Decision Determinations

Medically Necessary: Services that are Medically Necessary include only those which have been established as safe and effective, are furnished under generally accepted professional standards to treat illness, injury or medical condition, and which, as determined by Blue Shield, are: (a) consistent with Blue Shield medical policy; (b) consistent with the symptoms or diagnosis; (c) not furnished primarily for the convenience of the patient, the attending Physician or other provider; (d) furnished at the most appropriate level which can be provided safely and effectively to the patient; and (e) not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of the Member's illness, injury, or disease.

Investigational/Experimental: A treatment, procedure, or drug is investigational when it has not been recognized as safe and effective for use in treating the particular condition in accordance with generally accepted professional medical standards. This includes services where approval by the federal or state governmental is required prior to use, but has not yet been granted.

Split Evaluation: Blue Shield of California/Blue Shield of California Life & Health Insurance Company (Blue Shield) policy review can result in a split evaluation, where a treatment, procedure, or drug will be considered to be investigational for certain indications or conditions, but will be deemed safe and effective for other indications or conditions, and therefore potentially medically necessary in those instances.

Prior Authorization Requirements (as applicable to your plan)

Within five days before the actual date of service, the provider must confirm with Blue Shield that the member's health plan coverage is still in effect. Blue Shield reserves the right to revoke an authorization prior to services being rendered based on cancellation of the member's eligibility. Final determination of benefits will be made after review of the claim for limitations or exclusions.

Questions regarding the applicability of this policy should be directed to the Prior Authorization Department at (800) 541-6652, or the Transplant Case Management Department at (800) 637-2066 ext. 3507708 or visit the provider portal at www.blueshieldca.com/provider.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. Blue Shield of California may consider published peer-reviewed scientific literature, national guidelines, and local standards of practice in developing its medical policy. Federal and state law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over medical policy and must be considered first in determining covered services. Member contracts may differ in their benefits. Blue Shield reserves the right to review and update policies as appropriate.

Appendix A

POLICY STATEMENT (No changes)	
BEFORE Red font: Verbiage removed	AFTER Blue font: Verbiage Changes/Additions
<p>Genetic Testing for Cardiac Ion Channelopathies 2.04.43</p> <p>Policy Statement: Long QT Syndrome Genetic testing to confirm a diagnosis of congenital long QT syndrome (LQTS) may be considered medically necessary when signs and/or symptoms of LQTS are present, but a definitive diagnosis cannot be made without genetic testing. This includes:</p> <ol style="list-style-type: none"> I. Individuals who do not meet the clinical criteria for LQTS (i.e., those with a Schwartz score less than 4): but have a moderate-to-high pretest probability (see Policy Guidelines section) based on the Schwartz score and/or other clinical criteria. <p>Genetic testing of asymptomatic individuals to determine future risk of LQTS may be considered medically necessary when at least one or more of the following criteria is met:</p> <ol style="list-style-type: none"> I. A close relative (i.e., first-, second-, or third-degree relative) with a known LQTS variant II. A close relative diagnosed with LQTS by clinical means whose genetic status is unavailable. <p>Genetic testing for LQTS for all other situations not meeting the criteria outlined above, including but not limited to determining prognosis and/or directing therapy in patients with known LQTS, is considered investigational.</p> <p>Brugada Syndrome Genetic testing to confirm a diagnosis of Brugada syndrome (BrS) may be considered medically necessary when signs and/or symptoms consistent with BrS (see Policy Guidelines section) are present, but a definitive diagnosis cannot be made without genetic testing.</p> <p>Genetic testing of asymptomatic individuals to determine future risk of BrS may be considered medically necessary when patients have a</p>	<p>Genetic Testing for Cardiac Ion Channelopathies 2.04.43</p> <p>Policy Statement: Long QT Syndrome Genetic testing to confirm a diagnosis of congenital long QT syndrome (LQTS) may be considered medically necessary when signs and/or symptoms of LQTS are present, but a definitive diagnosis cannot be made without genetic testing. This includes:</p> <ol style="list-style-type: none"> I. Individuals who do not meet the clinical criteria for LQTS (i.e., those with a Schwartz score less than 4) but have a moderate-to-high pretest probability (see Policy Guidelines section) based on the Schwartz score and/or other clinical criteria. <p>Genetic testing of asymptomatic individuals to determine future risk of LQTS may be considered medically necessary when at least one or more of the following criteria is met:</p> <ol style="list-style-type: none"> I. A close relative (i.e., first-, second-, or third-degree relative) with a known LQTS variant II. A close relative diagnosed with LQTS by clinical means whose genetic status is unavailable. <p>Genetic testing for LQTS for all other situations not meeting the criteria outlined above, including but not limited to determining prognosis and/or directing therapy in patients with known LQTS, is considered investigational.</p> <p>Brugada Syndrome Genetic testing to confirm a diagnosis of Brugada syndrome (BrS) may be considered medically necessary when signs and/or symptoms consistent with BrS (see Policy Guidelines section) are present, but a definitive diagnosis cannot be made without genetic testing.</p> <p>Genetic testing of asymptomatic individuals to determine future risk of BrS may be considered medically necessary when patients have a</p>

POLICY STATEMENT
(No changes)

<p>BEFORE Red font: Verbiage removed</p>	<p>AFTER Blue font: Verbiage Changes/Additions</p>
<p><u>close relative</u> (i.e., first-, second-, or third-degree relative) with a known BrS variant.</p> <p>Genetic testing for BrS for all other situations not meeting the criteria outlined above is considered investigational.</p> <p>Catecholaminergic Polymorphic Ventricular Tachycardia Genetic testing to confirm a diagnosis of catecholaminergic polymorphic ventricular tachycardia (CPVT) may be considered medically necessary when signs and/or symptoms of CPVT are present, but a definitive diagnosis cannot be made without genetic testing.</p> <p>Genetic testing of asymptomatic individuals to determine future risk of CPVT may be considered medically necessary when at least one or more of the following criteria is met:</p> <ol style="list-style-type: none"> I. A <u>close relative</u> (i.e., first-, second-, or third-degree relative) with a known CPVT variant II. A <u>close relative</u> diagnosed with CPVT by clinical means whose genetic status is unavailable. <p>Genetic testing for CPVT for all other situations not meeting the criteria outlined above is considered investigational.</p> <p>Short QT Syndrome Genetic testing of asymptomatic individuals to determine future risk of short QT syndrome (SQTS) may be considered medically necessary when patients have a <u>close relative</u> (i.e., first-, second-, or third-degree relative) with a known SQTS variant.</p> <p>Genetic testing for SQTS for all other situations not meeting the criteria outlined above is considered investigational.</p>	<p><u>close relative</u> (i.e., first-, second-, or third-degree relative) with a known BrS variant.</p> <p>Genetic testing for BrS for all other situations not meeting the criteria outlined above is considered investigational.</p> <p>Catecholaminergic Polymorphic Ventricular Tachycardia Genetic testing to confirm a diagnosis of catecholaminergic polymorphic ventricular tachycardia (CPVT) may be considered medically necessary when signs and/or symptoms of CPVT are present, but a definitive diagnosis cannot be made without genetic testing.</p> <p>Genetic testing of asymptomatic individuals to determine future risk of CPVT may be considered medically necessary when at least one or more of the following criteria is met:</p> <ol style="list-style-type: none"> I. A <u>close relative</u> (i.e., first-, second-, or third-degree relative) with a known CPVT variant II. A <u>close relative</u> diagnosed with CPVT by clinical means whose genetic status is unavailable. <p>Genetic testing for CPVT for all other situations not meeting the criteria outlined above is considered investigational.</p> <p>Short QT Syndrome Genetic testing of asymptomatic individuals to determine future risk of short QT syndrome (SQTS) may be considered medically necessary when patients have a <u>close relative</u> (i.e., first-, second-, or third-degree relative) with a known SQTS variant.</p> <p>Genetic testing for SQTS for all other situations not meeting the criteria outlined above is considered investigational.</p>