

8.01.49	Intensity-Modulated Radiotherapy: Abdomen and Pelvis		
Original Policy Date:	April 5, 2007	Effective Date:	August 1, 2022
Section:	8.0 Therapy	Page:	Page 1 of 28

Policy Statement

- I. Intensity-modulated radiotherapy (IMRT) may be considered **medically necessary** as an approach to delivering radiotherapy for patients with cancer of the anus and anal canal.
- II. IMRT may be considered **medically necessary** when dosimetric planning with standard 3-dimensional conformal radiotherapy predicts that the radiation dose to an adjacent organ would result in unacceptable normal tissue toxicity, as documented by **all** of the following:
 - A. IMRT is used to treat cancer of **one or more** of the following:
 1. Stomach (gastric)
 2. Hepatobiliary tract
 3. Pancreas
 4. Esophageal cancer
 5. Rectal locations
 6. Gynecologic tumors (to include cervical, endometrial, and vulvar cancers)
 7. Other pelvic or abdominal tumor not listed
 - B. Documentation of **one or more** of the following:
 1. The target volume is in close proximity to critical structures that must be protected and **both** of the following: * (see source below)
 - a. Planned 3D-CRT exposure to critical adjacent structures is above normal tissue constraints
 - b. Planned IMRT exposure to these critical adjacent structures does not exceed normal tissue constraints
 2. An immediately adjacent area has been previously irradiated and abutting portals must be established with high precision
- III. Intensity-modulated radiotherapy is considered **investigational** for all other uses in the abdomen and pelvis not addressed above.

NOTE: Refer to [Appendix A](#) to see the policy statement changes (if any) from the previous version.

Policy Guidelines

For intensity-modulated radiotherapy (IMRT) to provide outcomes superior to 3D-CRT, there must be a clinically meaningful decrease in the radiation exposure to normal structures with IMRT compared with 3D-CRT. There is no standardized definition for a clinically meaningful decrease in radiation dose. In principle, a clinically meaningful decrease would signify a significant reduction in anticipated complications of radiation exposure. To document a clinically meaningful reduction in dose, dosimetry planning studies should demonstrate a significant decrease in the maximum dose of radiation delivered per unit of tissue, and/or a significant decrease in the volume of normal tissue exposed to potentially toxic radiation doses. While radiation tolerance dose levels for normal tissues are well-established, the decrease in the volume of tissue exposed that is needed to provide a clinically meaningful benefit has not been standardized. Therefore, precise parameters for a clinically meaningful decrease cannot be provided.

Requests for the above exceptions and all other indications not discussed in this policy will be reviewed on a case-by-case basis.

*The Normal Tissue Constraint Guidelines are derived from the textbook: Radiation Oncology: A Question-Based Review published by Lippincott Williams & Wilkins, 2010 [author: Hristov et al., 2010]. According to the author, most dosages were derived from randomized studies or consensus guidelines; however, pediatric dose constraints will vary greatly from protocol to protocol. Sources used in the development of the guidelines included the American Brachytherapy Society (ABS); Clinical practice guidelines from Johns Hopkins Hospital (JHH); the International Journal of Radiation Oncology *Biology* Physics (IJROBP); the National Comprehensive Cancer Network (NCCN), Quantitative Analyses of Normal Tissue Effects in the Clinic (QUANTEC); and the Radiation Therapy Oncology Group (RTOG) protocols at the time of publication.

The following guidelines are only intended to serve as a guide and may not be applicable to all clinical scenarios.

Organ	Constraints
Central Nervous System (1.8-2.0 Gray/fraction [Gy/fx])	
• Spinal Cord	max 50 Gy (full cord cross-section); tolerance increases by 25% 6 mos after 1st course (for re-irradiation)
• Brain	max 72 Gy (partial brain); avoid >2 Gy/fx or hyperfractionation
• Chiasm/Optic Nerves	max 55 Gy
• Brainstem	Entire brainstem <54 Gy, V59 Gy <1-10 cc
• Eyes (globe)	mean <35 Gy, max 54 Gy
• Lens	max 7 Gy
• Retina	max 50 Gy
• Lacrimal Gland	max 40 Gy
• Inner ear/cochlea	mean <=45 Gy (consider constraining to <=35 Gy with concurrent cisplatin)
• Pituitary gland	max 45 Gy (for panhypopituitarism, lower for GH deficiency)
• Cauda equina	max 60 Gy
Central Nervous System (single fraction)	
• Spinal Cord	max 13 Gy (if 3 fxs, max 20 Gy)
• Brain	V12 Gy <5-10 cc
• Chiasm/Optic Nerves	max 10 Gy
• Brainstem	max 12.5 Gy
• Sacral plexus	V18 <0.035 cc, V14.4 <5 cc
• Cauda equina	V16 <0.035 cc, V14 <5 cc
Head and Neck (1.8-2.0 Gy/fx)	
• Parotid gland(s)	mean <25 Gy (both glands) or mean <20 Gy (1 gland)
• Submandibular gland(s)	mean <35 Gy
• Larynx	mean <=44 Gy, V50 <=27%, max 63-66 Gy (when risk of tumor involvement is limited)
• TMJ/mandible	max 70 Gy (if not possible, then V75 <1 cc)
• Oral cavity	Non-oral cavity cancer: mean <30 Gy, avoid hot spots >60 Gy Oral cavity cancer: mean <50 Gy, V55 <1 cc, max 65 Gy
• Esophagus (cervical)	V45 <33%
• Pharyngeal constrictors	mean <50 Gy
• Thyroid	V26 <20%
Thoracic (1.8-2.0 Gy/fx)	
• Brachial plexus	max 66 Gy, V60 <5%
• Lung (combined lung for lung cancer treatment)	mean <20-23 Gy, V20 <30%-35%
• Lung (ipsilateral lung for breast cancer treatment)	V25 <10%
• Single lung (after pneumonectomy)	V5 <60%, V20 <4-10%, MLD <8 Gy
• Bronchial tree	max 80 Gy

Organ	Constraints
• Heart (lung cancer treatment)	Heart V45 <67%; V60 <33%
• Heart (breast cancer treatment)	V25 <10%
• Esophagus	V50 <32% ;V60 <33%
Thoracic (hypofractionation)	
Note: the max dose limits refer to volumes >0.035 cc (~3 mm ³).	
• Spinal cord	1 fraction: 14 Gy 3 fractions: 18 Gy (6 Gy/fx) 4 fractions: 26 Gy (6.5 Gy/fx) 5 fractions: 30 Gy (6 Gy/fx)
• Esophagus	1 fraction: 15.4 Gy 3 fractions: 30 Gy (10 Gy/fx) 4 fractions: 30 Gy (7.5 Gy/fx) 5 fractions: 32.5 Gy (6.5 Gy/fx)
• Brachial plexus	1 fraction: 17.5 Gy 3 fractions: 21 Gy (7 Gy/fx) 4 fractions: 27.2 Gy (6.8 Gy/fx) 5 fractions: 30 Gy (6 Gy/fx)
• Heart/Pericardium	1 fraction: 22 Gy 3 fractions: 30 Gy (10 Gy/fx) 4 fractions: 34 Gy (8.5 Gy/fx) 5 fractions: 35 Gy (7 Gy/fx)
• Great vessels	1 fraction: 37 Gy 3 fractions: 39 Gy (13 Gy/fx) 4 fractions: 49 Gy (12.25 Gy/fx) 5 fractions: 55 Gy (11 Gy/fx)
• Trachea/Large Bronchus	1 fraction: 20.2 Gy 3 fractions: 30 Gy (10 Gy/fx) 4 fractions: 34.8 Gy (8.7 Gy/fx) 5 fractions: 40 Gy (8 Gy/fx)
• Rib	1 fraction: 30 Gy 3 fractions: 30 Gy (10 Gy/fx) 4 fractions: 32 Gy (7.8 Gy/fx) 5 fractions: 32.5 Gy (6.5 Gy/fx)
• Skin	1 fraction: 26 Gy 3 fractions: 30 Gy (10 Gy/fx) 4 fractions: 36 Gy (9 Gy/fx) 5 fractions: 40 Gy (8 Gy/fx)
• Stomach	1 fraction: 12.4 Gy 3 fractions: 27 Gy (9 Gy/fx) 4 fractions: 30 Gy (7.5 Gy/fx) 5 fractions: 35 Gy (7 Gy/fx)
Gastrointestinal (GI) (1.8–2.0 Gy/fx)	
• Stomach	TD 5/5 whole stomach: 45 Gy
• Small bowel	V45 <195 cc
• Liver (metastatic disease)	mean liver <32 Gy (liver = normal liver minus gross disease)
• Liver (primary liver cancer)	mean liver <28 Gy (liver = normal liver minus gross disease)
• Colon	45 Gy, max dose 55 Gy
• Kidney (bilateral)	mean <18 Gy, V28 <20%, V23 Gy <30%, V20 <32%, V12 <55%. If mean kidney dose to 1 kidney >18 Gy, then constrain remaining kidney to V6 <30%.
Gastrointestinal (GI) (single fraction)	
• Duodenum	V16 <0.035 cc, V11.2 <5 cc
• Kidney (Cortex)	V8.4 <200 cc
• Kidney (Hilum)	V10.6 <66%
• Colon	V14.3 <20 cc, V18.4 <0.035 cc
• Jejunum/Ileum	V15.4 <0.035 cc, V11.9 <5 cc
• Stomach	V16 <0.035 cc, V11.2 <10 cc
• Rectum	V18.4 <0.035 cc, V14.3 <20 cc

Organ	Constraints
Genitourinary (GU) (1.8-2.0 Gy/fx)	
• Femoral heads	V50 <5%
• Rectum	V75 <15% , V70 <20%, V65 <25%, V60 <35%, V50 <50%
• Bladder	V80 <15%, V75 <25%, V70 <35%, V65 <50%
• Testis	V3 <50%
• Penile bulb	Mean dose to 95% of the volume <50 Gy. D70 < /=70 Gy, D50 < /=50 Gy
Genitourinary (GU) (LDR prostate brachytherapy)	
• Urethra	Volume of urethra receiving 150% of prescribed dose (Ur150) <30%
• Rectum	Volume of rectum receiving 100% of prescribed dose (RV100) <0.5 cc
Gynecological (GYN)	
• Bladder point (cervical brachytherapy)	Max 80 Gy (LDR equivalent dose)
• Rectal point (cervical brachytherapy)	Max 75 Gy (LDR equivalent dose)
• Proximal vagina (mucosa) (cervical brachytherapy)	Max 120 Gy (LDR equivalent dose)
• Distal vagina (mucosa) (cervical brachytherapy)	Max 98 Gy (LDR equivalent dose)

Coding

The following CPT codes are used for simple and complex IMRT delivery:

- **77385:** Intensity modulated radiation treatment delivery (IMRT), includes guidance and tracking, when performed; simple
- **77386:** Intensity modulated radiation treatment delivery (IMRT), includes guidance and tracking, when performed; complex

The Centers for Medicare & Medicaid Services (CMS) did not implement these CPT codes and instead created HCPCS G codes with the language of the previous CPT codes. So the following codes may be used for IMRT:

- **G6015:** Intensity modulated treatment delivery, single or multiple fields/arcs, via narrow spatially and temporally modulated beams, binary, dynamic MLC, per treatment session
- **G6016:** Compensator-based beam modulation treatment delivery of inverse planned treatment using three or more high resolution (milled or cast) compensator, convergent beam modulated fields, per treatment session

Code 77301 remains valid:

- **77301:** Intensity modulated radiotherapy plan, including dose-volume histograms for target and critical structure partial tolerance specifications

The following CPT code may also be used and is to be reported only once per IMRT plan:

- **77338:** Multi-leaf collimator (MLC) device(s) for intensity modulated radiation therapy (IMRT), design and construction per IMRT plan

The following CPT and HCPCS codes may also be used:

- **77261:** Therapeutic radiology treatment planning; simple
- **77262:** Therapeutic radiology treatment planning; intermediate
- **77263:** Therapeutic radiology treatment planning; complex
- **77293:** Respiratory motion management simulation (List separately in addition to code for primary procedure)
- **77300:** Basic radiation dosimetry calculation, central axis depth dose calculation, TDF, NSD, gap calculation, off axis factor, tissue inhomogeneity factors, calculation of non-ionizing radiation surface and depth dose, as required during course of treatment, only when prescribed by the treating physician

- **77306:** Teletherapy isodose plan; simple (1 or 2 unmodified ports directed to a single area of interest), includes basic dosimetry calculation(s)
- **77307:** Teletherapy isodose plan; complex (multiple treatment areas, tangential ports, the use of wedges, blocking, rotational beam, or special beam considerations), includes basic dosimetry calculation(s)
- **77331:** Special dosimetry (e.g., TLD, microdosimetry) (specify), only when prescribed by the treating physician
- **77332:** Treatment devices, design and construction; simple (simple block, simple bolus)
- **77334:** Treatment devices, design and construction; complex (irregular blocks, special shields, compensators, wedges, molds or casts)
- **77370:** Special medical radiation physics consultation
- **77470:** Special treatment procedure (e.g., total body irradiation, hemibody radiation, per oral or endocavitary irradiation)
- **77336:** Continuing medical physics consultation, including assessment of treatment parameters, quality assurance of dose delivery, and review of patient treatment documentation in support of the radiation oncologist, reported per week of therapy
- **77427:** Radiation treatment management, 5 treatments
- **77014:** Computed tomography guidance for placement of radiation therapy fields
- **77417:** Therapeutic radiology port image(s)
- **77387:** Guidance for localization of target volume for delivery of radiation treatment, includes intrafraction tracking, when performed
- **G6001:** Ultrasonic guidance for placement of radiation therapy fields
- **G6002:** Stereoscopic x-ray guidance for localization of target volume for the delivery of radiation therapy
- **G6017:** Intra-fraction localization and tracking of target or patient motion during delivery of radiation therapy (e.g., 3D positional tracking, gating, 3D surface tracking), each fraction of treatment

Additional documentation will be required to confirm medical necessity when a given code is billed at a frequency greater than that allowed in the code set.

Allowable Codes and Frequencies for IMRT/Proton

Description	Code	Maximum per course of treatment	Notes
Clinical Treatment Planning	77261, 77262 or 77263	1	
Simulation	77280, 77285, 77290	0	May not be billed with 77301. 1 unit of 77290 + 1 boost is allowed for proton therapy when using 77295 instead
Verification Simulation	77280	0	One per simulation allowed
Respiratory Motion Management	77293	0	1 for breast, lung, and upper abdominal or thoracic cancer areas
3D CRT Plan	77295	0	May not be billed with 77301. 1 unit may be allowed for proton therapy.
IMRT Plan	77301	1	If comparison 3D plan is generated, it is included in 77301
Basic Dosimetry	77300	4+ 1 boost, up to a max of 10 with documentation	0 if billed with 77306, 77307, 77321, 0394T or 0395T
Teletherapy Isodose Plan, Simple	77306	1 for mid-Tx change in volume/contour	Not on the same day as 77300; may not bill 77306 and 77307 together; documentation of medical necessity is required for more than 1
Teletherapy Isodose Plan, Complex	77307	1 for mid-Tx change in volume/contour	Not on the same day as 77300; may not bill 77306 and 77307 together; documentation of medical necessity is required for more than 1

Description	Code	Maximum per course of treatment	Notes
Special Dosimetry Calculation	77331	0	Needs documentation for review
Treatment Devices, Designs, and Construction	77332, 77333, 77334	1, 5 or 10	-If billed w/ MLC (77338): 1 -If billed w/o MLC: 5 (any combination) -More may be allowed when documentation of medical necessity is provided (such as additional beams), maximum of 10
Multi-leaf Collimator (MLC)	77338	1	MLC may not be reported in conjunction with HCPCS G6016
Special Radiation Physics Consult	77370	0	May allow x 1; documentation of medical necessity required
Special MD Consultation (Special Tx Procedure)	77470	0	May allow x 1; documentation of medical necessity required
Medical Physics Management	77336	8	Allowed once per 5 courses of therapy
Radiation Treatment Management	77427	8	Allowed once per 5 courses of therapy
Radiation (IMRT or Proton) Delivery, prostate and breast cancer	IMRT 77385 or G6015;	Using IMRT or Proton: 28 for prostate cancer	Prostate cancer: Documentation of medical necessity needed for more than 28 treatments
	Proton 77520, 77522, 77523	Using IMRT only: -16 for breast cancer without boost -24 for breast cancer with boost (IMRT only)	Breast cancer: documentation of medical necessity needed for treatments beyond 16 IMRT delivery sessions without boost and/or 24 IMRT delivery sessions with boost.
Radiation (IMRT or Proton) Delivery, all other cancers	IMRT 77385, 77386; or G6015-G6016; Proton 77520, 77522, 77523, 77525	No limit	All cancers other than hypofractionated prostate or breast

Description

Radiotherapy may be an integral component of the treatment of cancers of the abdomen and pelvis. Intensity-modulated radiotherapy (IMRT) has been proposed as a method that allows adequate radiation to the tumor while minimizing the radiation dose to surrounding normal tissues and critical structures.

Related Policies

- Intensity-Modulated Radiotherapy of the Prostate
- Intensity-Modulated Radiotherapy: Cancer of the Head and Neck or Thyroid
- Intensity-Modulated Radiotherapy: Central Nervous System Tumors
- Radiation Oncology

Benefit Application

Benefit determinations should be based in all cases on the applicable contract language. To the extent there are any conflicts between these guidelines and the contract language, the

contract language will control. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

Some state or federal mandates (e.g., Federal Employee Program [FEP]) prohibits plans from denying Food and Drug Administration (FDA)-approved technologies as investigational. In these instances, plans may have to consider the coverage eligibility of FDA-approved technologies on the basis of medical necessity alone.

Regulatory Status

In general, IMRT systems include intensity modulators which control, block, or filter the intensity of radiation; and RT planning systems which plan the radiation dose to be delivered.

A number of intensity modulators have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process. Intensity modulators include the Innocure Intensity Modulating Radiation Therapy Compensators (Innocure), cleared in 2006, and the decimal tissue compensator (Southeastern Radiation Products), cleared in 2004. FDA product code: IXI. Intensity modulators may be added to standard linear accelerators to deliver IMRT when used with proper treatment planning systems.

RT planning systems have also been cleared for marketing by the FDA through the 510(k) process. They include the FOCUS Radiation Treatment Planning System (Computerized Medical Systems) in 2002, Prowess Panther™ (Prowess) in 2003, TiGRT (LinaTech) in 2009, the RayDose (RaySearch Laboratories) in 2008, and the Eclipse Treatment Planning System (Varian Medical Systems) in 2017. FDA product code: MUJ.

Fully integrated IMRT systems also are available. These devices are customizable and support all stages of IMRT delivery, including planning, treatment delivery, and health record management. Varian Medical Systems has several 510(k) marketing clearances for high-energy linear accelerator systems that can be used to deliver precision RT such as IMRT. FDA product code: IYE.

Rationale

Background

Radiotherapy Techniques

Radiation therapy may be administered externally (i.e., a beam of radiation is directed into the body) or internally (i.e., a radioactive source is placed inside the body, near a tumor).¹ External radiotherapy (RT) techniques include "conventional" or 2-dimensional (2D) RT, 3-dimensional (3D) conformal RT, and intensity-modulated radiation therapy (IMRT).

Conventional External-Beam Radiotherapy

Methods to plan and deliver RT have evolved that permit more precise targeting of tumors with complex geometries. Conventional 2D treatment planning utilizes X-ray films to guide and position radiation beams.¹ Bony landmarks bones visualized on X-ray are used to locate a tumor and direct the radiation beams. The radiation is typically of uniform intensity.

Three-Dimensional Conformal Radiotherapy

Radiation treatment planning has evolved to use 3D images, usually from computed tomography (CT) scans, to more precisely delineate the boundaries of the tumor and to discriminate tumor tissue from adjacent normal tissue and nearby organs at risk for radiation damage. Three-dimensional conformal RT (3D-CRT) involves initially scanning the patient in the position that will be used for the radiation treatment.¹ The tumor target and surrounding normal organs are then outlined in 3D on the scan. Computer software assists in determining the

orientation of radiation beams and the amount of radiation the tumor and normal tissues receive to ensure coverage of the entire tumor in order to minimize radiation exposure for at risk normal tissue and nearby organs. Other imaging techniques and devices such as multileaf collimators (MLCs) may be used to "shape" the radiation beams. Methods have also been developed to position the patient and the radiation portal reproducibly for each fraction and to immobilize the patient, thus maintaining consistent beam axes across treatment sessions.

Intensity-Modulated Radiotherapy

IMRT is the more recent development in external radiation. Treatment planning and delivery are more complex, time-consuming, and labor-intensive for IMRT than for 3D-CRT. Similar to 3D-CRT, the tumor and surrounding normal organs are outlined in 3D by a scan and multiple radiation beams are positioned around the patient for radiation delivery.¹ In IMRT, radiation beams are divided into a grid-like pattern, separating a single beam into many smaller "beamlets". Specialized computer software allows for "inverse" treatment planning. The radiation oncologist delineates the target on each slice of a CT scan and specifies the target's prescribed radiation dose, acceptable limits of dose heterogeneity within the target volume, adjacent normal tissue volumes to avoid, and acceptable dose limits within the normal tissues. Based on these parameters and a digitally reconstructed radiographic image of the tumor, surrounding tissues, and organs at risk, computer software optimizes the location, shape, and intensities of the beam ports to achieve the treatment plan's goals.

Increased conformality may permit escalated tumor doses without increasing normal tissue toxicity and is proposed to improve local tumor control, with decreased exposure to surrounding, normal tissues, potentially reducing acute and late radiation toxicities. Better dose homogeneity within the target may also improve local tumor control by avoiding underdosing within the tumor and may decrease toxicity by avoiding overdosing.

Other advanced techniques that may further improve RT treatment by improving dose distribution. These techniques are considered variations of IMRT. Volumetric modulated arc therapy delivers radiation from a continuous rotation of the radiation source. The principal advantage of volumetric modulated arc therapy is greater efficiency in treatment delivery time, reducing radiation exposure and improving target radiation delivery due to less patient motion. Image-guided RT involves the incorporation of imaging before and/or during treatment to more precisely deliver RT to the target volume.

Literature Review

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function-including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and the quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Note that the evidence for the following abdominal and pelvic cancers has not yet been reviewed and is beyond the scope of this review: bladder, kidney, and ureter cancer and sarcoma.

Intensity-Modulated Radiotherapy for Cancers of the Abdomen and Pelvis

Multiple-dose planning studies generate 3-dimensional conformal radiotherapy (3D-CRT) and intensity-modulated radiotherapy (IMRT) treatment plans from the same scans and then compare predicted dose distributions within the target area and adjacent organs. Results of such planning studies have shown that IMRT is better than 3D-CRT with respect to conformality to the target and dose homogeneity within the target. Results have also demonstrated that IMRT delivers less radiation to nontarget areas. Dosimetry studies using stationary targets generally confirm these predictions. However, because patients move during treatment, dosimetry with stationary targets only approximate actual radiation doses received. Based on these dosimetry studies, radiation oncologists expect IMRT to improve treatment outcomes compared with those of 3D-CRT.

Comparative studies of radiation-induced adverse events from IMRT versus alternative radiation delivery would constitute definitive evidence of establishing the benefit of IMRT. Single-arm series of IMRT can give insights into the potential for benefit, particularly if an adverse event expected to occur at high rates is shown to decrease significantly. Studies of treatment benefit are also important to establish that IMRT is at least as good as other types of delivery, but absent such comparative trials, it is likely that benefit from IMRT is at least as good as with other types of delivery.

In general, when the indication for IMRT is to avoid radiation to sensitive areas, dosimetry studies have been considered sufficient evidence to demonstrate that harm would be avoided using IMRT. For other indications, such as using IMRT to provide better tumor control, comparative studies of health outcomes are needed to demonstrate such a benefit.

Gastrointestinal Tract Cancers

Clinical Context and Therapy Purpose

The purpose of IMRT in patients who have gastrointestinal tract cancers is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does the use of IMRT for the treatment of patients with gastrointestinal tract cancers improve net health outcomes?

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is patients with gastrointestinal (GI) cancers (e.g., stomach, hepatobiliary, and pancreatic cancers) who are recommended for radiotherapy (RT).

Interventions

The therapy being considered is IMRT. Intensity-modulated radiotherapy uses computer software and magnetic resonance imaging for increased conformality, permitting the delivery of higher doses of radiation to the tumor while limiting the exposure of surrounding normal tissues.

Comparators

The following therapy is currently being used: 3D-CRT. This therapy uses 3-dimensional images typically from computed tomography to discriminate tumor tissue from adjacent normal tissue and nearby organs. Computer algorithms are used to estimate radiation doses being delivered to each treatment segment.

Outcomes

The general outcomes of interest are overall survival (OS), recurrence (locoregional control), quality of life, and treatment-related adverse events (e.g., toxicity). Toxicity can be assessed using the U.S. Department of Health and Human Services grading criteria for adverse events (1=mild, 2=moderate, 3=severe or medically significant, 4=life-threatening, and 5=death).

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Stomach

Systematic Review

Ren et al (2019) completed a systematic review and meta-analysis evaluating the efficacy and safety of IMRT versus 3D-CRT that included 9 controlled clinical trials enrolling 516 patients with gastric cancer.² Results revealed a slightly improved 3-year OS rate (risk ratio [RR], 1.16; 95% CI, 0.98 to 1.36) and a significantly better 2-year OS rate with IMRT (RR, 2.49; 95% CI, 1.18 to 5.25; $p=.02$) as compared to 3D-CRT. Additionally, the 3-year rate of locoregional recurrence was improved with IMRT versus 3D-CRT (RR, 0.62; 95% CI, 0.39 to 0.98; $p<.05$). Similar 3-year disease-free survival rates were noted between the techniques (RR, 1.16; 95% CI, 0.95 to 1.43; $p>.05$). No significant differences in liver, GI, and kidney toxicity were observed among patients receiving IMRT compared with 3D-CRT. Limitations of this analysis included the small number of enrolled subjects (i.e., the majority of studies had <100 subjects), retrospective nature of included studies, which increased the risk of selective reporting bias, and the heterogeneity of IMRT or 3D-CRT techniques in studies. Additionally, the detail and radiation fields of RT varied considerably among the studies, impacting the efficacy and toxicity seen by investigators.

Nonrandomized Comparative Studies

Boda-Heggemann et al (2009) evaluated the efficacy and safety of 2 different adjuvant chemoradiotherapy regimens using 3D-CRT or IMRT in 2 consecutive cohorts who underwent primarily D2 resection for gastric cancer.³ A subsequent report (2013) from this group included 27 3D-CRT patients and 38 IMRT patients.⁴ The cohorts were generally well-matched, with American Joint Committee on Cancer advanced stage (II-IV) disease. Most (96%) who received 3D-CRT were treated with 5-fluorouracil plus folinic acid. Patients in the IMRT cohort received capecitabine plus oxaliplatin (70%) or 5-fluorouracil plus folinic acid (30%). Radiation was delivered to a total prescribed dose of 45 gray (Gy) at 1.8 Gy per fraction. In the 3D-CRT cohort, 5 patients received less than 45 Gy because of treatment intolerance. Two patients in the IMRT cohort did not tolerate the full course, and 1 patient received 47 Gy. Overall, the IMRT plus chemotherapy regimen decreased renal toxicity with a trend toward improved survival (Table 1). However, interpretation of the safety and efficacy of IMRT in this study is limited by differences in the chemotherapy regimens.

Table 1. Outcomes for IMRT With Capecitabine Plus Oxaliplatin versus 3D-CRT With 5-FU Plus FA for Stomach Cancer

Outcomes	3-Dimensional Conformal Radiotherapy	Intensity-Modulated Radiotherapy	p
Sample	27	38	
Renal toxicity, n (%)	2 (8)	0	.021
Median disease-free survival, mo	14	35	.069

Median overall survival, mo	18	43	.060
Actutimes 2-y overall survival, %	37	67	
Actutimes 5-y overall survival, %	22	44	

Adapted from Boda-Heggemann et al (2009, 2013).^{3,4}

3D-CRT: 3-dimensional conformal radiotherapy; 5-FU: 5-fluorouracil; FA: folinic acid; IMRT: intensity-modulated radiotherapy.

Hepatobiliary

Fuller et al (2009) compared a retrospective series with a historical control cohort. Clinical results using image-guided IMRT (n=24) were compared with results using 3D-CRT (n=24) in patients with primary adenocarcinoma of the biliary tract.⁵ Most patients underwent postsurgical chemoradiotherapy with concurrent fluoropyrimidine-based regimens. Treatment plans prescribed 46 to 56 Gy to the planning target volume that included the tumor and involved lymph nodes, in daily fractions of 1.8 to 2 Gy. Both groups received boost doses of 4 to 18 Gy as needed. The IMRT cohort had a median OS of 17.6 months (range, 10.3 to 32.3 months), while the 3D-CRT cohort had a median OS of 9.0 months (range, 6.6 to 17.3 months). There was no significant differences between patient cohorts in acute GI toxicity. Generalization of results is limited by the small numbers of patients, use of retrospective chart review data, nonrepresentative case spectrum (mostly advanced/metastatic disease), and comparison to a nonconcurrent control RT cohort.

Pancreatic

Literature searches have identified a few comparative studies and case series on IMRT for pancreatic cancer. For example, Lee et al (2016) reported on a prospective comparative study of GI toxicity in patients treated with concurrent chemoradiotherapy plus IMRT or 3D-CRT for treatment of borderline resectable pancreatic cancer.⁶ Treatment selection was by patient choice after consultation with a radiation oncologist. Symptoms of dyspepsia, nausea or vomiting, and diarrhea did not differ between groups. Upper endoscopy revealed more patients with gastroduodenal ulcers in the 3D-CRT group than in the IMRT group (Table 2). Overall survival was longer in the IMRT group than in the 3D-CRT group but the interpretation of survival results was limited by the risk of bias in this nonrandomized study.

Prasad et al (2016) retrospectively reviewed charts of patients with locally advanced pancreatic cancer who were treated with IMRT (n=134) or 3D-CRT (n=71).⁷ Propensity score analysis was performed to account for potential confounding variables, including age, sex, radiation dose, RT field size, and concurrent RT. Grade 2 GI toxicity occurred in significantly more patients treated with 3D-CRT than with IMRT (propensity score odds ratio [OR], 1.26; 95% CI, 1.08 to 1.45; p=.001; Table 2). Hematologic toxicity and median survival were similar in the 2 groups.

Table 2. Outcomes for IMRT versus 3D-CRT for Pancreatic Cancer

Comparison	3D-CRT	IMRT	p
Lee et al (2016) ⁶	n=40	n=44	
Grade 1-2 gastroduodenal ulcers, n (%)	11 (42.3)	3 (9.1)	.003
Overall survival, mo	15.8	22.6	.006
Prasad et al (2016) ⁷	n=71	n=134	
Grade 2+ gastrointestinal toxicity, n (%)	24 (33.8)	21 (15.7)	.001
Overall survival whole population, mo	NR	NR	NS

3D-CRT: 3-dimensional conformal radiotherapy; IMRT: intensity-modulated radiotherapy; NR: not reported; NS: not significant.

Section Summary: Gastrointestinal Tract Cancers

The evidence on IMRT for GI tract cancers includes nonrandomized comparative studies, retrospective series, and a systematic review. Intensity-modulated radiotherapy has been compared with 3D-CRT for the treatment of stomach, hepatobiliary, and pancreatic cancers, with some data reporting longer OS and decreased toxicity with IMRT. For the treatment of stomach cancer, IMRT improved survival compared with 3D-CRT, with a comparable or

improved safety profile. The evidence on hepatobiliary cancer includes a series of historical controls that found an increase in median survival with no difference in toxicity. Two comparative studies (1 prospective, 1 retrospective) were identified on IMRT for pancreatic cancer. The prospective comparative study found an increase in survival with a reduction in GI toxicity, while the retrospective study found a decrease in GI toxicity. Although most studies were limited by their retrospective designs and changes in practice patterns over time, the available evidence would suggest that IMRT improves survival and decreases toxicity better than 3D-CRT in patients with GI cancers.

Gynecologic Cancers

Clinical Context and Therapy Purpose

The purpose of IMRT in patients who have gynecologic cancers is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does the use of IMRT for the treatment of patients with gynecologic cancers improve net health outcomes?

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is patients with gynecologic cancers (i.e., cervical and endometrial cancers) who are recommended for radiotherapy (RT).

Interventions

The therapy being considered is IMRT. Intensity-modulated radiotherapy uses computer software and magnetic resonance imaging for increased conformality, permitting the delivery of higher doses of radiation to the tumor while limiting the exposure of surrounding normal tissues.

Comparators

The following therapy is currently being used: 3D-CRT. This therapy uses 3-dimensional images typically from computed tomography to discriminate tumor tissue from adjacent normal tissue and nearby organs. Computer algorithms are used to estimate radiation doses being delivered to each treatment segment.

Outcomes

The general outcomes of interest are OS, recurrence (locoregional control), quality of life, and treatment-related adverse events (e.g., toxicity). Toxicity can be assessed using the U.S. Department of Health and Human Services grading criteria for adverse events (1=mild, 2=moderate, 3=severe or medically significant, 4=life-threatening, and 5=death).

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Review

Lin et al (2018) completed a meta-analysis of 6 studies enrolling 1008 subjects in order to compare the efficacy and safety of IMRT with 3D-CRT or 2D (2-dimensional)-RT for definitive treatment of cervical cancer.⁸ Results revealed a nonsignificant difference in 3-year OS (OR, 2.41; 95% CI, 0.62 to 9.39; p=.21) and disease-free survival rates (OR, 1.44; 95% CI, 0.69 to 3.01; p=.33) between IMRT and 3D-CRT or 2D-RT. However, IMRT was associated with a significantly

reduced rate of acute GI and genitourinary (GU) toxicity: Grade 2 GI: OR, 0.5; 95% CI, 0.28 to 0.89; $p=.02$; Grade 3 or higher GI: OR, 0.55; 95% CI, 0.32 to 0.95; $p=.03$; Grade 2 GU: OR, 0.41; 95% CI, 0.2 to 0.84; $p=.01$; Grade 3 or higher GU: OR, 0.31; 95% CI, 0.14 to 0.67; $p=.003$. Some chronic GU toxicity also occurred less frequently with IMRT (Grade 3: OR, 0.09; 95% CI, 0.01 to 0.67; $p=.02$). This analysis had several limitations including the fact that most included studies had relatively small sample sizes and were retrospective and nonrandomized in nature. Additionally, some of the included studies did not compare clinical outcomes between the RT techniques.

Randomized Controlled Trials

Klopp et al (2018) designed a randomized trial that measured the impact of pelvic IMRT versus standard 4-field RT on patient-reported toxicity and quality of life in 278 women with cervical and endometrial cancer.⁹ Results revealed that the mean Expanded Prostate Cancer Index Composite (EPIC) bowel score decreased significantly less in the IMRT as compared to the standard RT group from baseline to end of RT (18.6 vs. 23.6 points; $p=.048$). Additionally, both the mean EPIC urinary score (5.6 vs. 10.4 points; $p=.03$) and Trial Outcome Index score (8.8 vs. 12.8 points; $p=.06$) declined significantly less with IMRT as compared to standard RT. Frequent or almost constant diarrhea was also reported more frequently among women receiving standard RT versus IMRT at the end of RT (51.9% vs. 33.7%; $p=.01$) and significantly more women administered standard RT were taking antidiarrheal medications 4 or more times daily (20.4% vs. 7.8%; $p=.04$).

A trial by Naik et al (2016) randomized 40 patients with cervical cancer to IMRT or 3D-CRT.¹⁰ Both arms received concurrent chemotherapy (cisplatin) plus RT at 50 Gy in 25 fractions. Dosimetric planning showed higher conformality and lower doses to organs at risk with IMRT. With follow-up through 90 days posttreatment, vomiting and acute GI and GU toxicity were significantly higher in the 3D-CRT group (Table 3).

Gandhi et al (2013) reported on a prospective randomized study that compared whole-pelvis IMRT with whole-pelvis 2D-RT in 44 patients with locally advanced cervical cancer.¹¹ Each treatment arm had 22 patients. The OS rate at 27 months was 88% with IMRT and 76% with 2D-RT ($p=.645$). However, fewer grade 2, 3, or 4 GI toxicities were experienced in the IMRT group than in the conventional RT group (Table 3).

Table 3. Acute Toxicity of Grade 2, 3 or 4 With 3D-CRT versus IMRT for Cervical Cancer

Toxicity	3D-CRT, n (%)	IMRT, n (%)	95% CI for the Difference	p
Naik et al (2016) ¹⁰ .				
Hematologic	8 (40)	7 (35)	-0.219 to 0.119	.644
Leukopenia	3 (15)	2 (10)	-0.1479 to 0.479	.424
Vomiting	7 (35)	3 (15)	0.338 to 0.061	.007
Acute gastrointestinal toxicity	9 (45)	4 (20)	-0.408 to -0.091	.003
Acute genitourinary toxicity	7 (35)	4 (20)	-0.295 to -0.004	.058
Gandhi et al (2013) ¹¹ .				
Gastrointestinal, grade ≥ 2	14 (64)	7 (32)	0.002 to 0.604	.034
Gastrointestinal, grade ≥ 3	6 (27)	1 (5)	0.003 to 0.447	.047
Genitourinary, grade ≥ 2	7 (32)	5 (24)	-0.202 to 0.361	.404
Genitourinary, grade ≥ 3	3 (14)	0 (0)	-0.019 to 0.291	.125

3D-CRT: 3-dimensional conformal radiotherapy; CI: confidence interval; IMRT: intensity-modulated radiotherapy.

Nonrandomized Comparative Studies

Shih et al (2016) conducted a retrospective comparison of bowel obstruction following IMRT (n=120) or 3D-CRT (n=104) after hysterectomy for endometrial or cervical cancer.¹² Groups were generally comparable, except more patients in the 3D-CRT group had open hysterectomy (81% vs. 47%, $p<.001$). Patients received regular examinations throughout a median follow-up of 67 months, and the 5-year rate of bowel obstruction was 0.9% in the IMRT group compared with 9.3% in the 3D-CRT group ($p=.006$). A body mass index of 30 kg/m² or more was also associated

with less bowel obstruction. However, on multivariate analysis, the only significant predictor of less bowel obstruction was IMRT ($p=.022$).

Chen et al (2014) reported on 101 patients with endometrial cancer treated with hysterectomy and adjuvant RT.¹³ No significant differences between IMRT patients ($n=65$) and CRT patients ($n=36$) were found in 5-year OS (82.9% vs. 93.5%; $p=.26$), local failure-free survival (93.7% vs. 89.3%; $p=.68$), or disease-free survival (88.0% vs. 82.8%; $p=.83$). However, IMRT patients experienced fewer acute and late toxicities.

Section Summary: Gynecologic Cancers

The evidence on IMRT for gynecologic cancers includes a systematic review, 3 RCTs, and nonrandomized comparative studies. There is limited comparative evidence on survival outcomes following IMRT or 3D-CRT. However, available results have generally been consistent that IMRT reduces GI and GU toxicity. Based on evidence with other cancers of the pelvis and abdomen in close proximity to organs at risk, it is expected that OS with IMRT would be at least as good as 3D-CRT, with a decrease in toxicity.

Anorectal Cancer

Clinical Context and Therapy Purpose

The purpose of IMRT in patients who have anorectal cancer is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does the use of IMRT for the treatment of patients with anorectal cancer improve net health outcomes?

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is patients with anorectal cancer who are recommended for radiotherapy (RT).

Interventions

The therapy being considered is IMRT. Intensity-modulated radiotherapy uses computer software and magnetic resonance imaging for increased conformality, permitting the delivery of higher doses of radiation to the tumor while limiting the exposure of surrounding normal tissues.

Comparators

The following therapy is currently being used: 3D-CRT. This therapy uses 3-dimensional images typically from computed tomography to discriminate tumor tissue from adjacent normal tissue and nearby organs. Computer algorithms are used to estimate radiation doses being delivered to each treatment segment.

Outcomes

The general outcomes of interest are OS, recurrence (locoregional control), quality of life, and treatment-related adverse events (e.g., toxicity). Toxicity can be assessed using the U.S. Department of Health and Human Services grading criteria for adverse events (1=mild, 2=moderate, 3=severe or medically significant, 4=life-threatening, and 5=death).

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Randomized Controlled Trials

Rattan et al (2016) conducted a small (n=20) RCT assessing IMRT for the treatment of anal canal cancer.¹⁴ Grade 3 GI toxicity during treatment was not observed in any patients in the IMRT group but was seen in 60% of patients treated with 3D-CRT (p=.010). Hematologic grade 3 toxicity was not seen in any patients treated with IMRT but was noted in 20% of patients treated with 3D-CRT (p=.228). Other parameters indicating better tolerance to treatment with IMRT were reduced need for parenteral fluid (10% vs. 60%, p=.019) and blood transfusion (0% vs. 20%, p=.060).

Nonrandomized Comparative Studies

Sun et al (2017) reported on a comparative analysis of the National Cancer Database of IMRT with 3D-CRT for the treatment of rectal adenocarcinoma.¹⁵ A total of 7386 patients with locally advanced rectal carcinoma were treated with neoadjuvant chemoradiotherapy (45 to 54 Gy) from 2006 to 2013; 3330 (45%) received IMRT and 4065 (55%) received 3D-CRT. Use of IMRT increased from 24% in 2006 to 50% in 2013. Patient age, race, insurance status, Charlson-Deyo comorbidity score, hospital type, income and education status, and clinical disease stage were not predictive of which RT was used. The mean radiation dose was higher with IMRT (4735 centigray vs. 4608 centigray, p<.001) and the occurrence of sphincter loss surgery was higher in the IMRT group (Table 4). However, patients treated with IMRT had a higher risk of positive margins. Multivariate analysis found no significant differences between the treatments for pathologic downstaging, unplanned readmission, 30-day mortality, or long-term survival. This study used unplanned readmission as a surrogate measure of adverse events but could not assess acute or late toxicity.

Table 4. Outcomes Following Radiochemotherapy With 3D-CRT or IMRT for Rectal Cancer

Outcome	3D-CRT, %	IMRT, %	Adjusted Odds Ratio	95% CI	p
Pathologic downstaging	57.0	55.0	0.89	0.79 to 1.01	.051
Sphincter loss surgery	28.3	34.7	1.32	1.14 to 1.52	<.001
Positive resection margin	5.6	8.0	1.57	1.21 to 2.03	<.001
Unplanned readmission	7.9	6.4	0.79	0.61 to 1.02	.07
30-d mortality	0.8	0.6	0.61	0.24 to 1.57	.31
Survival at 5 y	64	64	1.06	0.89 to 1.28	.47

Adapted from Sun et al (2017).¹⁵

3D-CRT: 3-dimensional conformal radiotherapy; CI: confidence interval; IMRT: intensity-modulated radiotherapy.

Huang et al (2017) reported on a retrospective comparison of outcomes and toxicity for pre-operative image-guided IMRT and 3D-CRT in locally advanced rectal cancer.¹⁶ A total of 144 consecutive patients treated between 2006 and 2015 were analyzed. The 3D-CRT group was treated with 45 Gy in 25 fractions while the IMRT group was treated with 45 Gy in 25 fractions with a simultaneous integrated boost of 0.2 Gy per day for the primary tumor up to a total dose of 50 Gy. Statistical analysis was performed for grade 0, 1, 2, 3, or 4 toxicity and was significant only for acute GI toxicity (p=.039; Table 5). Four-year OS and disease-free survival did not differ between the 2 groups. Multivariate analysis found IMRT to be an independent predictor of local failure-free survival (hazard ratio, 0.35; 95% CI, 0.11 to 0.95; p=.042).

Table 5. Grade 3 or Greater Toxicity Following Chemoradiotherapy for Rectal Cancer

Comparison	3D-CRT (n=99), n (%)	IMRT (n=45), n (%)
Skin	3 (3)	1 (2.2)
Acute gastrointestinal	14 (14.1)	3 (6.7)
Acute genitourinary	3 (3)	0 (0)
Hematologic	2 (2.0)	0 (0)
Late gastrointestinal	10 (10.1)	2 (4.4)

Late genitourinary	3 (3.1)	0 (0)
--------------------	---------	-------

Adapted from Huang et al (2017).¹⁶

3D-CRT: 3-dimensional conformal radiotherapy; IMRT: intensity-modulated radiotherapy.

In a retrospective review of 89 consecutive patients (52 IMRT, 37 3D-CRT), Chuong et al (2013) found that 3-year OS, progression-free survival, locoregional control, and colostomy-free survival did not differ significantly between patients treated with IMRT or with 3D-CRT ($p > .1$).¹⁷ Adverse events with 3D-CRT were more frequent and severe and required more treatment breaks than IMRT (11 days vs. 4 days; $p = .006$) even though the median duration of treatment breaks did not differ significantly (12.2 days vs. 8.0 days; $p = .35$). Intensity-modulated radiotherapy patients had fewer acute grade 3 or 4 nonhematologic toxicity ($p < .001$), improved late grade 3 or 4 GI toxicity ($p = .012$), and fewer acute grade 3 or 4 skin toxicity ($p < .001$) than 3D-CRT patients. Dasgupta et al (2013) retrospectively reviewed 223 patients (45 IMRT, 178 CRT) to compare outcomes for anal cancer.¹⁸ They reported that 2-year OS, distant metastases-free survival, and locoregional recurrence-free survival did not differ significantly between patients in the IMRT and CRT groups.

Dewas et al (2012) retrospectively reviewed 51 patients with anal cancer treated with IMRT ($n = 24$) or 3D-CRT ($n = 27$).¹⁹ Outcomes also did not differ significantly between patients in the IMRT and 3D-CRT groups in 2-year OS, locoregional relapse-free survival, and colostomy-free survival. Grade 3 acute toxicity occurred in 11 IMRT patients and in 10 3D-CRT patients.

Case Series

A GI toxicity study by Devisetty et al (2009) reported on 45 patients who received concurrent chemotherapy plus IMRT for anal cancer.²⁰ Intensity-modulated radiotherapy was administered to a dose of 45 Gy in 1.8-Gy fractions, with areas of gross disease subsequently boosted with 9 to 14.4 Gy. Acute GU toxicity was grade 0 in 25 (56%) cases, grade 1 in 10 (22%) patients, grade 2 in 5 (11%) patients, with no grade 3 or 4 toxicities reported; 5 (11%) patients reported no GU tract toxicities. Grades 3 and 4 leukopenia were reported in 26 (56%) cases, neutropenia in 14 (31%), and anemia in 4 (9%). Acute GI toxicity included grade 0 in 2 (4%) patients, grade 1 in 11 (24%), grade 2A in 25 (56%), grade 2B in 4 (9%), grade 3 in 3 (7%), and no grade 4 toxicities. Univariate analysis of data from these patients suggested a statistical correlation between the volume of bowel that received 30 Gy or more of radiation and the risk for clinically significant (grade ≥ 2) GI toxicities.

Pepek et al (2010) retrospectively analyzed toxicity and disease outcomes associated with IMRT in 47 patients with anal cancer.²¹ Thirty-one patients had squamous cell carcinoma. Intensity-modulated radiotherapy was prescribed to a dose of at least 54 Gy to areas of gross disease at 1.8 Gy per fraction. Forty (89%) patients received concurrent chemotherapy with various agents and combinations. The 2-year OS for all patients was 85%. Eight (18%) patients required treatment breaks. Toxicities included grade 4 leukopenia (7%) and thrombocytopenia (2%); grade 3 leukopenia (18%) and anemia (4%); and grade 2 skin toxicity (93%). These rates were lower than those reported in previous trials of chemoradiation, where grade 3 or 4 skin toxicity was noted in about 50% of patients and grade 3 or 4 GI toxicity noted in about 35%. In addition, the rate of treatment breaks was lower than in many studies; and some studies of chemoradiation included a break from RT.

Section Summary: Anorectal Cancer

The evidence on IMRT for anorectal cancer includes a small RCT with 20 patients, non-randomized comparative studies, and case series. Survival outcomes have not differed significantly between IMRT and 3D-CRT. Studies have found that patients receiving IMRT plus chemotherapy for the treatment of anal cancer experience fewer acute and late adverse events than patients receiving 3D-CRT plus chemotherapy, primarily in GI toxicity.

Esophageal Cancer

Clinical Context and Therapy Purpose

The purpose of IMRT in patients who have esophageal cancer is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does the use of IMRT for the treatment of patients with esophageal cancer improve net health outcomes?

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is patients with esophageal cancer who are recommended for radiotherapy (RT).

Interventions

The therapy being considered is IMRT. Intensity-modulated radiotherapy uses computer software and magnetic resonance imaging for increased conformality, permitting the delivery of higher doses of radiation to the tumor while limiting the exposure of surrounding normal tissues.

Comparators

The following therapy is currently being used: 3D-CRT. This therapy uses 3-dimensional images typically from computed tomography to discriminate tumor tissue from adjacent normal tissue and nearby organs. Computer algorithms are used to estimate radiation doses being delivered to each treatment segment.

Outcomes

The general outcomes of interest are OS, recurrence (locoregional control), quality of life, and treatment-related adverse events (e.g., toxicity). Toxicity can be assessed using the U.S. Department of Health and Human Services grading criteria for adverse events (1=mild, 2=moderate, 3=severe or medically significant, 4=life-threatening, and 5=death).

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

Xu et al (2017) performed a systematic review and meta-analysis to compare IMRT and 3D-CRT in the treatment of esophageal cancer with regard to dosimetry and clinical outcomes (n=7 studies).²² For the dosimetric comparison of organs at risk, 5 studies were included. Results revealed that the mean dose of 3D-CRT was significantly higher as compared to IMRT for the lung (mean difference dose: 2.18; 95% CI, 0.83 to 3.53; p=.002), with patients treated with ≥ 20 Gy having significantly higher irradiated volumes for 3D-CRT than for IMRT. For the heart, the mean dose was not significantly different between 3D-CRT and IMRT (mean difference dose: 0.17; 95% CI, -3.73 to 4.07; p=.93); however, the heart in patients treated with 50 Gy had significantly higher irradiated volumes for 3D-CRT. The maximum dose in the spinal cord revealed no difference between the 2 RT techniques (p=.33). Evaluated clinical outcomes included OS (n=3 studies; 871 patients) and toxicity (n=2 studies; 205 patients). The 3-year OS was significantly improved with IMRT as compared to 3D-CRT (OR, 0.68; 95% CI, 0.52 to 0.90; p=.007). No difference between the 2 RT techniques was seen with regard to the incidence of radiation pneumonitis or radiation esophagitis, regardless of grade. Limitations of the review were the small number of studies

available for OS and toxicity outcome analyses and the retrospective nature of clinical outcomes studies.

Nonrandomized Comparative Studies

Lan et al (2020) retrospectively compared survival outcomes and symptomatic radiation pneumonitis in patients with esophageal cancer who received either IMRT (n=297) or 3D-CRT (n=91) from 2010 through 2017.²³ The median age of patients was 60 years and the median radiation dose for the entire cohort was 60 Gy. Results revealed significantly improved OS (p=.001), progression-free survival (PFS) (p=.008), and distant-metastases free survival (p=.011) with IMRT versus 3D-CRT; locoregional failure-free survival was not significantly different between the groups (p=.721). Intensity-modulated radiotherapy was also associated with significantly less grade ≥ 2 radiation pneumonitis as compared to 3D-CRT (5.4% vs. 23.1%; p<.001).

Ito et al (2017) retrospectively compared failure patterns and toxicities between IMRT (n=32) and 3D-CRT (n=48) in patients with cervical esophageal cancer.²⁴ All patients were administered systemic chemotherapy consisting of either induction chemotherapy or concurrent chemoradiotherapy, with or without adjuvant chemotherapy. The median follow-up of the entire cohort was 24.6 months and the median follow-up time for survivors was 35.9 months. Results revealed a 3 year OS of 81.6%, 57.2% (p=.037 vs. IMRT), and 66.6% for the IMRT, 3D-CRT, and total groups, respectively. However, there was no significant difference between IMRT and 3D-CRT with regard to complete response rate (75% vs. 68.9%; p=.62). Rates of locoregional control or PFS were not different between the groups as well. Overall, 47 patients developed recurrence of any type; there was no apparent difference in the failure pattern between the 2 RT techniques. The incidence of late toxicities was also not significantly different between IMRT and 3D-CRT. Ten patients in the IMRT groups were salvaged, and 60% survived without recurrence compared to 20% of the 3D-CRT group.

Haefner et al (2017) reported a retrospective analysis of 93 patients with esophageal cancer and compared outcomes and acute toxicity among patients receiving definitive CRT with either 3D-CRT (n=49) or IMRT (n=44).²⁵ The median follow-up for all patients was 20.1 months. The 1- and 3-year local relapse rates were 20.4% and 28.6% in the 3D-CRT group and 15.9% and 22.7% in the IMRT group, respectively (p=.62 for the 3-year rate). Median PFS and OS were not significantly different between the groups; 13.8 months 3D-CRT versus 16.6 months IMRT (p=.448) and 18.4 months 3D-CRT versus 42 months IMRT (p=.198), respectively. The incidence of acute toxicities (dysphasia, radio dermatitis, nausea/vomiting, mucositis, bleeding, pneumonitis) were also not significantly different between the 2 RT techniques.

Section Summary: Esophageal Cancer

The evidence on IMRT for esophageal cancer includes a systematic review and nonrandomized comparative studies. Survival outcomes from studies have been mixed with some concluding improved survival with IMRT and others demonstrating no difference from 3D-CRT. Similarly, some studies have concluded that IMRT is associated with a reduced dose for organs at risk and potentially less radiation-related toxicity as compared to 3D-CRT.

Summary of Evidence

For individuals who have GI tract cancers who receive IMRT, the evidence includes non-randomized comparative studies, retrospective series, and a systematic review. Relevant outcomes are OS, disease-specific survival, recurrence, quality of life, and treatment-related morbidity. IMRT has been compared with 3D-CRT for the treatment of stomach, hepatobiliary, and pancreatic cancers. Evidence has been inconsistent with the outcome of survival, with some studies reporting increased survival among patients receiving IMRT compared with patients receiving 3D-CRT, and other studies reporting no difference between groups. However, most studies found that patients receiving IMRT experienced significantly less GI toxicity compared with patients receiving 3D-CRT. The available comparative evidence, together with dosimetry studies of organs at risk, would suggest that IMRT decreases toxicity compared with

3D-CRT in patients who had GI cancers. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have gynecologic cancers who receive IMRT, the evidence includes a systematic review, 3 RCTs, and nonrandomized comparative studies. Relevant outcomes are OS, disease-specific survival, recurrence, quality of life, and treatment-related morbidity. There is limited comparative evidence on survival outcomes following IMRT or 3D-CRT. However, results are generally consistent that IMRT reduces GI and genitourinary toxicity. Based on evidence with other cancers of the pelvis and abdomen that are proximate to organs at risk, it is expected that OS with IMRT would be at least as good as 3D-CRT, with a decrease in toxicity. A reduction in GI toxicity is likely to improve the quality of life in patients with gynecologic cancer. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome. For individuals who have anorectal cancer who receive IMRT, the evidence includes a small RCT (n=20), nonrandomized comparative studies, and case series. Relevant outcomes are OS, disease-specific survival, recurrence, quality of life, and treatment-related morbidity. Survival outcomes have not differed significantly between patients receiving IMRT and 3D-CRT. However, studies have found that patients receiving IMRT plus chemotherapy for the treatment of anal cancer experience fewer acute and late adverse events than patients receiving 3D-CRT plus chemotherapy, primarily in GI toxicity. A reduction in GI toxicity is likely to improve the quality of life in patients with anorectal cancer. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have esophageal cancer who receive IMRT, the evidence includes a systematic review and nonrandomized comparative studies. Relevant outcomes are OS, disease-specific survival, recurrence, quality of life, and treatment-related morbidity. Survival outcomes have been mixed with some studies concluding that IMRT is associated with a significant improvement in OS, PFS, or distant-metastases free survival versus 3D-CRT and others reporting no difference between the RT techniques. Intensity-modulated radiotherapy appears to be associated with a reduced dose for organs at risk and may result in less radiation-induced toxicity. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Clinical Input From Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2012 Input

In response to requests from Blue Cross Blue Shield Association, input was received from 1 physician specialty society (4 reviewers) and 3 academic medical centers in 2012. Input was mixed but there was support for use of intensity-modulated radiotherapy (IMRT) in a number of cancers discussed herein. In general, this support was based on normal tissue constraints for radiation doses and whether these dose constraints could not be met without IMRT.

2010 Input

In response to requests from Blue Cross Blue Shield Association, input was received from 1 physician specialty society (2 reviewers) and 3 academic medical centers in 2010. There was support for use of IMRT in a number of cancers discussed herein. In general, this support was based on normal tissue constraints for radiation doses and whether these dose constraints could not be met without IMRT.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

National Comprehensive Cancer Network Guidelines

Gastrointestinal Tract Cancers

The NCCN guideline (v.2.2021) for gastric cancer indicates that IMRT "may be used in clinical settings where dose reduction to organs at risk is required, and cannot be achieved by 3D techniques."²⁶ In addition, target volumes need to be carefully defined and encompassed while taking into account variations in stomach filling and respiratory motion.

The NCCN guideline (v.2.2021) for hepatobiliary cancers states that "All tumors irrespective of the location may be amenable to external beam radiation therapy (3D conformal radiation therapy, intensity-modulated radiation therapy [IMRT], or stereotactic body radiation therapy [SBRT])."²⁷

Intensity-modulated radiotherapy is mentioned as an option in the NCCN guideline (v.2.2021) for pancreatic adenocarcinoma, stating that IMRT "is increasingly being applied for the therapy of locally advanced pancreatic adenocarcinoma and in the adjuvant setting with the aim of increasing radiation dose to the gross tumor while minimizing toxicity to surrounding tissues."²⁸ In addition, the guideline states that "there is no clear consensus on the appropriate maximum dose of radiation when IMRT technique is used."

Gynecologic Cancers

For cervical cancer, the NCCN guideline (v.1.2021) indicates IMRT "is helpful in minimizing the dose to the bowel and other critical structures in the post-hysterectomy setting and in treating the para-aortic nodes when necessary." This technique can also be useful "when high doses are required to treat gross disease in regional lymph nodes."²⁹ Intensity-modulated radiotherapy "should not be used as a routine alternative to brachytherapy for treatment of central disease in patients with an intact cervix." The guideline also mentions that "very careful attention to detail and reproducibility (including consideration of target and normal tissue definitions, patient and internal organ motion, soft tissue deformation, and rigorous dosimetric and physics quality assurance) is required for proper delivery of IMRT and related highly conformal technologies." The NCCN guideline (v.2.2021) on uterine neoplasms states that radiotherapy for uterine neoplasms includes external-beam radiotherapy and/or brachytherapy but that IMRT may be considered "for normal tissue sparing."³⁰

The NCCN guideline (v.1.2021) on ovarian cancer does not mention IMRT.³¹

Anorectal Cancers

The NCCN guideline (v.1.2021) for anal carcinoma states that IMRT "is preferred over 3D conformal RT [radiotherapy] in the treatment of anal carcinoma"; and that its use "requires expertise and careful target design to avoid reduction in local control by so-called 'marginal-miss'. "³²

The NCCN guideline (v.1.2021) on rectal cancer indicates that "...IMRT... should only be used in the setting of a clinical trial, in unique clinical situations such as reirradiation of previously treated patients with recurrent disease, or unique anatomical situations."³³

Esophageal Cancer

The NCCN guideline (v.2.2021) for esophageal and esophagogastric junction cancers states that IMRT..."is appropriate in clinical settings where reduction in dose to organs at risk (e.g., heart, lungs) is required that cannot be achieved by 3-D techniques."³⁴

American Society for Radiation Oncology

In 2020, the American Society for Radiation Oncology published a clinical practice guideline on RT for cervical cancer.³⁵ One key question within the guideline asked when it was appropriate to deliver IMRT for women administered definitive or postoperative RT for cervical cancer?

Recommendations regarding this clinical scenario included:

- "In women with cervical cancer treated with postoperative RT with or without chemotherapy, IMRT is recommended to decrease acute and chronic toxicity." This was a strong recommendation based on moderate quality evidence for acute toxicity and low quality evidence for chronic toxicity.
- "In women with cervical cancer treated with definitive RT with or without chemotherapy, IMRT is conditionally recommended to decrease acute and chronic toxicity." This was a conditional recommendation based on moderate quality evidence for acute and chronic toxicity.

The guideline also notes that there are "no data that IMRT improves disease-specific survival or OS over 2D/3D techniques."

In 2021, the American Society for Radiation Oncology published a clinical practice guideline on RT for rectal cancer.³⁶ Within this guideline, IMRT-specific recommendations include:

- "For patients with rectal cancer treated with RT, an IMRT/volumetric modulated arc therapy (VMAT) technique is conditionally recommended (low quality of evidence). IMRT/VMAT may be beneficial when the external iliac nodes and/or the inguinal nodes require treatment or when 3-D conformal techniques may confer a higher risk for toxicity.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 6.

Table 6. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT03239626	Postoperative Hypofractionated Intensity-Modulated Radiation Therapy in Cervical Cancer: A Prospective Exploratory Trial (POHIM_RT Trial)	120	Apr 2025
NCT03239613	Postoperative Hypofractionated Intensity-Modulated Radiation Therapy with Concurrent Chemotherapy in Cervical Cancer: A Prospective Exploratory Trial (POHIM_CCRT Trial)	84	Apr 2024
<i>Unpublished</i>			
NCT02964468	Multicenter Dose-escalation Trial of Radiotherapy in Patients with Locally Advanced Rectal Cancer	525	May 2020

NCT: national clinical trial.

References

1. Shinohara E, Whaley JT. Last reviewed: March 3, 2020. <https://www.oncolink.org/cancer-treatment/radiation/introduction-to-radiation-therapy/radiation-therapy-which-type-is-right-for-me>. Accessed June 3, 2021

2. Ren F, Li S, Zhang Y, et al. Efficacy and safety of intensity-modulated radiation therapy versus three-dimensional conformal radiation treatment for patients with gastric cancer: a systematic review and meta-analysis. *Radiat Oncol*. May 22 2019; 14(1): 84. PMID 31118042
3. Boda-Heggemann J, Hofheinz RD, Weiss C, et al. Combined adjuvant radiochemotherapy with IMRT/XELOX improves outcome with low renal toxicity in gastric cancer. *Int J Radiat Oncol Biol Phys*. Nov 15 2009; 75(4): 1187-95. PMID 19409725
4. Boda-Heggemann J, Weiss C, Schneider V, et al. Adjuvant IMRT/XELOX radiochemotherapy improves long-term overall- and disease-free survival in advanced gastric cancer. *Strahlenther Onkol*. May 2013; 189(5): 417-23. PMID 23558673
5. Fuller CD, Dang ND, Wang SJ, et al. Image-guided intensity-modulated radiotherapy (IG-IMRT) for biliary adenocarcinomas: Initial clinical results. *Radiother Oncol*. Aug 2009; 92(2): 249-54. PMID 19324442
6. Lee KJ, Yoon HI, Chung MJ, et al. A Comparison of Gastrointestinal Toxicities between Intensity-Modulated Radiotherapy and Three-Dimensional Conformal Radiotherapy for Pancreatic Cancer. *Gut Liver*. Mar 2016; 10(2): 303-9. PMID 26470767
7. Prasad S, Cambridge L, Huguet F, et al. Intensity modulated radiation therapy reduces gastrointestinal toxicity in locally advanced pancreas cancer. *Pract Radiat Oncol*. Mar-Apr 2016; 6(2): 78-85. PMID 26577010
8. Lin Y, Chen K, Lu Z, et al. Intensity-modulated radiation therapy for definitive treatment of cervical cancer: a meta-analysis. *Radiat Oncol*. Sep 14 2018; 13(1): 177. PMID 30217165
9. Klopp AH, Yeung AR, Deshmukh S, et al. Patient-Reported Toxicity During Pelvic Intensity-Modulated Radiation Therapy: NRG Oncology-RT0G 1203. *J Clin Oncol*. Aug 20 2018; 36(24): 2538-2544. PMID 29989857
10. Naik A, Gurjar OP, Gupta KL, et al. Comparison of dosimetric parameters and acute toxicity of intensity-modulated and three-dimensional radiotherapy in patients with cervix carcinoma: A randomized prospective study. *Cancer Radiother*. Jul 2016; 20(5): 370-6. PMID 27368915
11. Gandhi AK, Sharma DN, Rath GK, et al. Early clinical outcomes and toxicity of intensity modulated versus conventional pelvic radiation therapy for locally advanced cervix carcinoma: a prospective randomized study. *Int J Radiat Oncol Biol Phys*. Nov 01 2013; 87(3): 542-8. PMID 24074927
12. Shih KK, Hajj C, Kollmeier M, et al. Impact of postoperative intensity-modulated radiation therapy (IMRT) on the rate of bowel obstruction in gynecologic malignancy. *Gynecol Oncol*. Oct 2016; 143(1): 18-21. PMID 27486131
13. Chen CC, Wang L, Lu CH, et al. Comparison of clinical outcomes and toxicity in endometrial cancer patients treated with adjuvant intensity-modulated radiation therapy or conventional radiotherapy. *J Formos Med Assoc*. Dec 2014; 113(12): 949-55. PMID 24144528
14. Rattan R, Kapoor R, Bahl A, et al. Comparison of bone marrow sparing intensity modulated radiotherapy (IMRT) and three-dimensional conformal radiotherapy (3DCRT) in carcinoma of anal canal: a prospective study. *Ann Transl Med*. Feb 2016; 4(4): 70. PMID 27004217
15. Sun Z, Adam MA, Kim J, et al. Intensity-Modulated Radiation Therapy Is Not Associated with Perioperative or Survival Benefit over 3D-Conformal Radiotherapy for Rectal Cancer. *J Gastrointest Surg*. Jan 2017; 21(1): 106-111. PMID 27510332
16. Huang CM, Huang MY, Tsai HL, et al. A retrospective comparison of outcome and toxicity of preoperative image-guided intensity-modulated radiotherapy versus conventional pelvic radiotherapy for locally advanced rectal carcinoma. *J Radiat Res*. Mar 01 2017; 58(2): 247-259. PMID 27738080
17. Chuong MD, Freilich JM, Hoffe SE, et al. Intensity-Modulated Radiation Therapy vs. 3D Conformal Radiation Therapy for Squamous Cell Carcinoma of the Anal Canal. *Gastrointest Cancer Res*. Mar 2013; 6(2): 39-45. PMID 23745158
18. Dasgupta T, Rothenstein D, Chou JF, et al. Intensity-modulated radiotherapy vs. conventional radiotherapy in the treatment of anal squamous cell carcinoma: a propensity score analysis. *Radiother Oncol*. May 2013; 107(2): 189-94. PMID 23692961

19. Dewas CV, Maingon P, Dalban C, et al. Does gap-free intensity modulated chemoradiation therapy provide a greater clinical benefit than 3D conformal chemoradiation in patients with anal cancer?. *Radiat Oncol*. Nov 29 2012; 7: 201. PMID 23190693
20. Devisetty K, Mell LK, Salama JK, et al. A multi-institutional acute gastrointestinal toxicity analysis of anal cancer patients treated with concurrent intensity-modulated radiation therapy (IMRT) and chemotherapy. *Radiother Oncol*. Nov 2009; 93(2): 298-301. PMID 19717198
21. Pepek JM, Willett CG, Wu QJ, et al. Intensity-modulated radiation therapy for anal malignancies: a preliminary toxicity and disease outcomes analysis. *Int J Radiat Oncol Biol Phys*. Dec 01 2010; 78(5): 1413-9. PMID 20231064
22. Xu D, Li G, Li H, et al. Comparison of IMRT versus 3D-CRT in the treatment of esophagus cancer: A systematic review and meta-analysis. *Medicine (Baltimore)*. Aug 2017; 96(31): e7685. PMID 28767597
23. Lan K, Zhu J, Zhang J, et al. Propensity score-based comparison of survival and radiation pneumonitis after definitive chemoradiation for esophageal cancer: Intensity-modulated radiotherapy versus three-dimensional conformal radiotherapy. *Radiother Oncol*. Aug 2020; 149: 228-235. PMID 32474127
24. Ito M, Kodaira T, Tachibana H, et al. Clinical results of definitive chemoradiotherapy for cervical esophageal cancer: Comparison of failure pattern and toxicities between intensity-modulated radiotherapy and 3-dimensional conformal radiotherapy. *Head Neck*. Dec 2017; 39(12): 2406-2415. PMID 28960561
25. Haefner MF, Lang K, Verma V, et al. Intensity-modulated versus 3-dimensional conformal radiotherapy in the definitive treatment of esophageal cancer: comparison of outcomes and acute toxicity. *Radiat Oncol*. Aug 15 2017; 12(1): 131. PMID 28810885
26. National Comprehensive Cancer Network. Gastric Cancer. Version. 2.2019. Updated May 13, 2020. https://www.nccn.org/professionals/physician_gls/pdf/gastric.pdf. Accessed June 3, 2021.
27. National Comprehensive Cancer Network. Hepatobiliary Cancers. Version. 2.2021. Updated June 1, 2020. https://www.nccn.org/professionals/physician_gls/pdf/hepatobiliary.pdf. Accessed June 3, 2021.
28. National Comprehensive Cancer Network. Pancreatic Adenocarcinoma. Version. 2.2021. Updated February 25, 2021. https://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf. Accessed June 3, 2021.
29. National Comprehensive Cancer Network. Cervical Cancer. Version. 1.2020. Updated October 2, 2020. https://www.nccn.org/professionals/physician_gls/pdf/cervical.pdf. Accessed June 3, 2021.
30. National Comprehensive Cancer Network. Uterine Neoplasms. Version 2.2021. Updated March 6, 2020. https://www.nccn.org/professionals/physician_gls/pdf/uterine.pdf. Accessed June 3, 2021.
31. National Comprehensive Cancer Network. Ovarian Cancer. Version.1.2020. Updated March 11, 2020. https://www.nccn.org/professionals/physician_gls/pdf/ovarian.pdf. Accessed June 8, 2020
32. National Comprehensive Cancer Network. Anal Carcinoma. Version. 2.2020. Updated February 16, 2021. https://www.nccn.org/professionals/physician_gls/pdf/anal.pdf. Accessed June 3, 2021.
33. National Comprehensive Cancer Network. Rectal Cancer. Version. 4.2020. Updated May 21, 2020. https://www.nccn.org/professionals/physician_gls/pdf/rectal.pdf. Accessed June 8, 2020
34. National Comprehensive Cancer Network. Esophageal and esophagogastric junction cancers. Version 2.2021. Updated March 9, 2021. https://www.nccn.org/professionals/physician_gls/pdf/esophageal.pdf. Accessed June 4, 2021.

35. Chino J, Annunziata CM, Beriwal S, et al. Radiation Therapy for Cervical Cancer: Executive Summary of an ASTRO Clinical Practice Guideline. *Pract Radiat Oncol.* Jul 2020; 10(4): 220-234. PMID 32473857
36. Wo JY, Anker CJ, Ashman JB, et al. Radiation Therapy for Rectal Cancer: Executive Summary of an ASTRO Clinical Practice Guideline. *Pract Radiat Oncol.* Jan-Feb 2021; 11(1): 13-25. PMID 33097436

Documentation for Clinical Review

Please provide the following documentation:

- (click here >>>) [Fax Back Form for Radiation Oncology Services](#)

Coding

This Policy relates only to the services or supplies described herein. Benefits may vary according to product design; therefore, contract language should be reviewed before applying the terms of the Policy.

The following codes are included below for informational purposes. Inclusion or exclusion of a code(s) does not constitute or imply member coverage or provider reimbursement policy. Policy Statements are intended to provide member coverage information and may include the use of some codes for clarity. The Policy Guidelines section may also provide additional information for how to interpret the Policy Statements and to provide coding guidance in some cases.

Type	Code	Description
CPT®	77014	Computed tomography guidance for placement of radiation therapy fields
	77261	Therapeutic radiology treatment planning; simple
	77262	Therapeutic radiology treatment planning; intermediate
	77263	Therapeutic radiology treatment planning; complex
	77293	Respiratory motion management simulation (List separately in addition to code for primary procedure)
	77300	Basic radiation dosimetry calculation, central axis depth dose calculation, TDF, NSD, gap calculation, off axis factor, tissue inhomogeneity factors, calculation of non-ionizing radiation surface and depth dose, as required during course of treatment, only when prescribed by the treating physician
	77301	Intensity modulated radiotherapy plan, including dose-volume histograms for target and critical structure partial tolerance specifications
	77306	Teletherapy isodose plan; simple (1 or 2 unmodified ports directed to a single area of interest), includes basic dosimetry calculation(s)
	77307	Teletherapy isodose plan; complex (multiple treatment areas, tangential ports, the use of wedges, blocking, rotational beam, or special beam considerations), includes basic dosimetry calculation(s)
	77331	Special dosimetry (e.g., TLD, microdosimetry) (specify), only when prescribed by the treating physician
	77332	Treatment devices, design and construction; simple (simple block, simple bolus)
77334	Treatment devices, design and construction; complex (irregular blocks, special shields, compensators, wedges, molds or casts)	

Type	Code	Description
	77336	Continuing medical physics consultation, including assessment of treatment parameters, quality assurance of dose delivery, and review of patient treatment documentation in support of the radiation oncologist, reported per week of therapy
	77338	Multi-leaf collimator (MLC) device(s) for intensity modulated radiation therapy (IMRT), design and construction per IMRT plan
	77370	Special medical radiation physics consultation
	77385	Intensity modulated radiation treatment delivery (IMRT), includes guidance and tracking, when performed; simple
	77386	Intensity modulated radiation treatment delivery (IMRT), includes guidance and tracking, when performed; complex
	77387	Guidance for localization of target volume for delivery of radiation treatment, includes intrafraction tracking, when performed
	77417	Therapeutic radiology port image(s)
	77427	Radiation treatment management, 5 treatments
	77470	Special treatment procedure (e.g., total body irradiation, hemibody radiation, per oral or endocavitary irradiation)
HCPCS	G6001	Ultrasonic guidance for placement of radiation therapy fields
	G6002	Stereoscopic x-ray guidance for localization of target volume for the delivery of radiation therapy
	G6015	Intensity modulated treatment delivery, single or multiple fields/arcs, via narrow spatially and temporally modulated beams, binary, dynamic MLC, per treatment session
	G6016	Compensator-based beam modulation treatment delivery of inverse planned treatment using 3 or more high resolution (milled or cast) compensator, convergent beam modulated fields, per treatment session
	G6017	Intra-fraction localization and tracking of target or patient motion during delivery of radiation therapy (e.g., 3D positional tracking, gating, 3D surface tracking), each fraction of treatment

Policy History

This section provides a chronological history of the activities, updates and changes that have occurred with this Medical Policy.

Effective Date	Action
04/05/2007	BCBSA Medical Policy adoption
05/16/2008	Policy Title Revision, criteria revised Added medically necessary indication for prostate cancer
12/05/2008	Policy Revision Added medically necessary indications for head and neck cancer, CNS lesions, and prostate fiducial markers
04/02/2010	Policy revision with position change Coding update
08/02/2010	Administrative Review
04/01/2011	Policy revision with position change
10/12/2012	Policy revision with position change
03/29/2013	Policy revision with position change
01/30/2015	Coding Update
03/30/2015	Policy title change from Intensity Modulated Radiation Therapy (IMRT) BCBSA Medical Policy adoption Policy revision without position change
10/01/2016	Policy revision without position change

Effective Date	Action
09/01/2017	Policy revision without position change
09/01/2018	Policy revision without position change
10/01/2019	Policy revision without position change
06/01/2020	Administrative update. Policy statement and guidelines updated.
11/20/2020	Annual review. No change to policy statement. Policy guidelines and literature updated. Coding update.
08/01/2021	Annual review. No change to policy statement. Policy guidelines updated.
12/01/2021	Administrative update. Policy statement, guidelines and literature updated.
08/01/2022	Annual review. No change to policy statement.

Definitions of Decision Determinations

Medically Necessary: Services that are Medically Necessary include only those which have been established as safe and effective, are furnished under generally accepted professional standards to treat illness, injury or medical condition, and which, as determined by Blue Shield, are: (a) consistent with Blue Shield medical policy; (b) consistent with the symptoms or diagnosis; (c) not furnished primarily for the convenience of the patient, the attending Physician or other provider; (d) furnished at the most appropriate level which can be provided safely and effectively to the patient; and (e) not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of the Member's illness, injury, or disease.

Investigational/Experimental: A treatment, procedure, or drug is investigational when it has not been recognized as safe and effective for use in treating the particular condition in accordance with generally accepted professional medical standards. This includes services where approval by the federal or state governmental is required prior to use, but has not yet been granted.

Split Evaluation: Blue Shield of California/Blue Shield of California Life & Health Insurance Company (Blue Shield) policy review can result in a split evaluation, where a treatment, procedure, or drug will be considered to be investigational for certain indications or conditions, but will be deemed safe and effective for other indications or conditions, and therefore potentially medically necessary in those instances.

Prior Authorization Requirements (as applicable to your plan)

Within five days before the actual date of service, the provider must confirm with Blue Shield that the member's health plan coverage is still in effect. Blue Shield reserves the right to revoke an authorization prior to services being rendered based on cancellation of the member's eligibility. Final determination of benefits will be made after review of the claim for limitations or exclusions.

Questions regarding the applicability of this policy should be directed to the Prior Authorization Department at (800) 541-6652, or the Transplant Case Management Department at (800) 637-2066 ext. 3507708 or visit the provider portal at www.blueshieldca.com/provider.

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. Blue Shield of California may consider published peer-reviewed scientific literature, national guidelines, and local standards of practice in developing its medical policy. Federal and state law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over medical policy and must be considered first in determining covered services. Member contracts may differ in their benefits. Blue Shield reserves the right to review and update policies as appropriate.

Appendix A

POLICY STATEMENT	
BEFORE Red font: Verbiage removed	AFTER Blue font: Verbiage Changes/Additions
<p>Intensity-Modulated Radiotherapy: Abdomen and Pelvis 8.01.49</p> <p>Policy Statement: Intensity-modulated radiotherapy (IMRT) may be considered medically necessary as an approach to delivering radiotherapy for patients with cancer of the anus and anal canal.</p> <p>IMRT may be considered medically necessary when dosimetric planning with standard 3-dimensional conformal radiotherapy predicts that the radiation dose to an adjacent organ would result in unacceptable normal tissue toxicity, as documented by all of the following:</p> <ul style="list-style-type: none"> I. IMRT is used to treat cancer of one or more of the following: <ul style="list-style-type: none"> A. Stomach (gastric) B. Hepatobiliary tract C. Pancreas D. Esophageal cancer E. Rectal locations F. Gynecologic tumors (to include cervical, endometrial, and vulvar cancers) G. Other pelvic or abdominal tumor not listed II. Documentation of one or more of the following: <ul style="list-style-type: none"> A. The target volume is in close proximity to critical structures that must be protected and both of the following: * (see source below) <ul style="list-style-type: none"> 1. Planned 3D-CRT exposure to critical adjacent structures is above normal tissue constraints 2. Planned IMRT exposure to these critical adjacent structures does not exceed normal tissue constraints B. An immediately adjacent area has been previously irradiated and abutting portals must be established with high precision 	<p>Intensity-Modulated Radiotherapy: Abdomen and Pelvis 8.01.49</p> <p>Policy Statement:</p> <ul style="list-style-type: none"> I. Intensity-modulated radiotherapy (IMRT) may be considered medically necessary as an approach to delivering radiotherapy for patients with cancer of the anus and anal canal. II. IMRT may be considered medically necessary when dosimetric planning with standard 3-dimensional conformal radiotherapy predicts that the radiation dose to an adjacent organ would result in unacceptable normal tissue toxicity, as documented by all of the following: <ul style="list-style-type: none"> A. IMRT is used to treat cancer of one or more of the following: <ul style="list-style-type: none"> 1. Stomach (gastric) 2. Hepatobiliary tract 3. Pancreas 4. Esophageal cancer 5. Rectal locations 6. Gynecologic tumors (to include cervical, endometrial, and vulvar cancers) 7. Other pelvic or abdominal tumor not listed B. Documentation of one or more of the following: <ul style="list-style-type: none"> 1. The target volume is in close proximity to critical structures that must be protected and both of the following: * (see source below) <ul style="list-style-type: none"> a. Planned 3D-CRT exposure to critical adjacent structures is above normal tissue constraints b. Planned IMRT exposure to these critical adjacent structures does not exceed normal tissue constraints 2. An immediately adjacent area has been previously irradiated and abutting portals must be established with high precision

POLICY STATEMENT

BEFORE Red font: Verbiage removed	AFTER Blue font: Verbiage Changes/Additions
Intensity-modulated radiotherapy is considered investigational for all other uses in the abdomen and pelvis not addressed above.	III. Intensity-modulated radiotherapy is considered investigational for all other uses in the abdomen and pelvis not addressed above.