

<b>4.02.04</b>	<b>Reproductive Techniques</b>		
<b>Original Policy Date:</b>	August 31, 2015	<b>Effective Date:</b>	November 1, 2021
<b>Section:</b>	4.0 OB/Gyn/Reproduction	<b>Page:</b>	Page 1 of 25

## Policy Statement

The following reproductive techniques may be considered **medically necessary** for **any** of the following:

- I. Blastocyst transfer
- II. Cryopreservation of testicular tissue in adult men with [azoospermia](#) as part of an intracytoplasmic sperm injection procedure
- I. Intracytoplasmic sperm injection for male factor infertility
- II. Cryopreservation of embryos, oocytes, ovarian tissue, sperm or testicular tissue (in post-pubertal men) when there is risk of iatrogenic sterilization from chemotherapy or similar medically necessary medical or surgical treatment when **all** of the following criteria are met:
  - A. No prior elective sterilization
  - B. No known infertility already present
  - C. Post-pubertal and less than 45 years of age (or cryopreservation is no longer desired if younger than age 45)

The following reproductive techniques are considered **investigational**:

- I. [Co-culture](#) of embryos
- II. Cryopreservation of testicular tissue in prepubertal boys or ovarian tissue in prepubertal girls
- III. [Intracytoplasmic sperm injection](#) (ICSI) in the absence of male factor infertility

**NOTE:** Refer to [Appendix A](#) to see the policy statement changes (if any) from the previous version.

## Policy Guidelines

**Azoospermia** means no sperm in the seminal fluid, either from obstruction or lack of production.

**ICSI** takes a single sperm and injects it directly into the ovum during the IVF process. Traditional IVF places live sperm (50,000) near the ovum in a laboratory dish and allow one of the sperm to penetrate the ovum. ICSI has only shown benefit if male factor infertility (too few, or abnormal sperm function) is present. Obtaining sperm as part of the ICSI process can involve taking some testicular tissue from which the sperm are removed. Leftover tissue can be cryopreserved in case it is needed again later.

**Assisted hatching** refers to mechanically disrupting the membrane around the ovum (zona pellucida) which persists after fertilization around the embryo. It usually dissolves on its own during implantation. Mechanical disruption has been proposed to help with implantation.

**Co-culture** refers to trying to enhance the culture medium the embryo is put into during the 2 to 3 days prior to transferring to the uterus (after the **embryo matures into a blastocyst**). The hope is to have more embryos progress and then to have a higher implantation or pregnancy rate.

**Cryopreservation of oocytes** (immature eggs) is less successful than cryopreservation of a fertilized embryo. Oocytes are more fragile than embryos and more prone to damage both during freezing and thawing.

**Testicular tissue from pre-pubertal boys** would contain stem cells that would later create sperm. Freezing and later thawing this tissue has not yet been shown to result in usable sperm in humans. Mature, usable sperm is not available from pre-pubertal boys.

Ovarian tissue from pre-pubertal girls has been able to be used for successful conception in a few case reports. The patients best suited for this and the techniques to be used are still unclear, and success rates remain low.

### Coding

The following CPT codes describe procedures that would be routinely performed in all assisted reproductive technology (ART) procedures involving in vitro fertilization (IVF):

- **58970:** Follicle puncture for oocyte retrieval, any method

Either:

- **89250:** Culture of oocyte(s)/embryo(s), less than 4 days
- **89272:** Extended culture of oocyte(s)/embryo(s), 4-7 days

Either:

- **58974:** Embryo transfer, intrauterine
- **58976:** Gamete, zygote, or embryo intrafallopian transfer, any method
- **89255:** Preparation of embryo for transfer (any method)
- **89260:** Sperm isolation; simple prep (e.g., sperm wash and swim-up) for insemination or diagnosis with semen analysis
- **89261:** Sperm isolation; complex prep (e.g., Percoll gradient, albumin gradient) for insemination or diagnosis with semen analysis
- **89268:** Insemination of oocytes
- **89280:** Assisted oocyte fertilization, microtechnique; less than or equal to 10 oocytes
- **89281:** Assisted oocyte fertilization, microtechnique; greater than 10 oocytes

The following CPT codes describe procedures that would *not* be routinely performed in all ART procedures involving IVF:

- **89253:** Assisted embryo hatching, microtechniques (any method). Only performed in women over the age of 40, or in cases in which prior ART attempts resulted in failed implantation
- **89257:** Sperm identification from aspiration (other than seminal fluid). Only performed in patients with oligospermia who have undergone a prior testicular or epididymal aspiration; typically performed as a part of an intracytoplasmic sperm injection procedure (ICSI)
- **89258:** Cryopreservation; embryo(s)
- **89259:** Cryopreservation; sperm
- **89264:** Sperm identification from testis tissue, fresh or cryopreserved. Only performed in patients with oligospermia who have undergone a prior testicular biopsy; typically performed as a part of an ICSI procedure
- **89342:** Storage (per year); embryo(s)
- **89343:** Storage (per year); sperm/semen
- **89344:** Storage (per year); reproductive tissue, testicular/ovarian
- **89346:** Storage (per year); oocyte(s)
- **89352:** Thawing of cryopreserved; embryo(s)
- **89353:** Thawing of cryopreserved; sperm/semen, each aliquot
- **89354:** Thawing of cryopreserved; reproductive tissue, testicular/ovarian
- **89356:** Thawing of cryopreserved; oocytes, each aliquot

The following CPT codes describe procedures that would be routinely performed as part of an intrauterine or intracervical artificial insemination:

- **58321:** Artificial insemination; intra-cervical
- **58322:** Artificial insemination; intra-uterine
- **58323:** Sperm washing for artificial insemination

Note also that "S" codes are available (see Coding section) that describe in vitro fertilization (IVF) globally.

The following codes are available for cryopreservation of oocytes:

- **89337:** Cryopreservation, mature oocyte(s)

The following CPT code [replaced](#) 0357T:

- **89398:** Unlisted reproductive medicine laboratory procedure

**Effective January 1, 2021**, the following CPT code has been **deleted**:

- **0058T:** Cryopreservation; reproductive tissue, ovarian

## Description

A variety of techniques are available to establish a viable pregnancy for couples who have been diagnosed with infertility and for whom assisted insemination has been unsuccessful.

## Related Policies

- Preimplantation Genetic Testing

## Benefit Application

Benefit determinations should be based in all cases on the applicable contract language. To the extent there are any conflicts between these guidelines and the contract language, the contract language will control. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

Some state or federal mandates (e.g., Federal Employee Program [FEP]) prohibits plans from denying Food and Drug Administration (FDA)-approved technologies as investigational. In these instances, plans may have to consider the coverage eligibility of FDA-approved technologies on the basis of medical necessity alone.

## Regulatory Status

There are no medical devices or diagnostic tests related to ARTs that require U.S. Food and Drug Administration approval or clearance.

## Rationale

### Background

#### Infertility

Infertility can be due either to female factors (i.e., pelvic adhesions, ovarian dysfunction, endometriosis, prior tubal ligation), male factors (i.e., abnormalities in sperm production, function, or transport or prior vasectomy), a combination of male and female factors, or unknown causes.

#### Treatment

Various reproductive techniques are available to establish a viable pregnancy; different techniques are used depending on the reason for infertility. Assisted reproductive technologies (ARTs), as defined by the Centers for Disease Control and Prevention and other organizations, refer to fertility treatments in which eggs or embryos are handled.<sup>1</sup> Not included in assisted reproduction is assisted insemination (artificial insemination) using sperm from either a woman's partner or a sperm donor. In most instances, assisted reproduction will involve in vitro fertilization (IVF), a procedure in which oocytes harvested from the female are inseminated in vitro with sperm harvested from the male. Following the fertilization procedure, the zygote is cultured and

ultimately transferred back into the female's uterus or fallopian tubes. In some instances, the oocyte and sperm are collected but no IVF takes place, and the gametes are reintroduced into the fallopian tubes. Examples of ARTs include, but are not limited to, gamete intrafallopian transfer, transuterine fallopian transfer, natural oocyte retrieval with intravaginal fertilization, pronuclear stage tubal transfer, tubal embryo transfer, zygote intrafallopian transfer, gamete, and embryo cryopreservation, oocyte, and embryo donation, and gestational surrogacy.

The various components of ART and implantation into the uterus can be broadly subdivided into oocyte harvesting procedures, which are performed on the female partner; sperm collection procedures, which are performed on the male partner; and the in vitro component (i.e., the laboratory procedures), which are performed on the collected oocyte and sperm. The final step is the implantation procedure.

Most CPT codes describing the various steps in ART procedures are longstanding. They include codes for oocyte retrieval, sperm isolation, culture and fertilization of the oocyte, and embryo, zygote, or gamete transfer into the uterus or fallopian tubes. Only the relatively new reproductive techniques (i.e., intracytoplasmic sperm injection [ICSI], assisted hatching, co-culture of embryos) and cryopreservation of reproductive tissue (i.e., testicular, ovarian, oocytes) will be considered within this evidence summary.

### Literature Review

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function, including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

### Assisted Hatching

#### Clinical Context and Therapy Purpose

Implantation of the embryo in the uterus is a key component of success with in vitro fertilization (IVF). Although the exact steps in implantation are poorly understood, normal rupture of the surrounding zona pellucida with escape of the developing embryo (termed hatching) is crucial. Mechanical disruption of the zona pellucida (i.e., assisted hatching) has been proposed as a mechanism to improve implantation rates.

The purpose of IVF with assisted hatching in patients with infertility is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does IVF with assisted hatching treat infertility and improve the net health outcome?

The following PICO was used to select literature to inform this review.

**Populations**

The relevant population of interest is patients who are infertile.

**Interventions**

The therapy being considered is IVF with assisted hatching.

**Comparators**

The following practice is currently being used to make decisions about infertility: IVF without assisted hatching.

**Outcomes**

The general outcomes of interest are live birth rates and infant abnormalities.

Follow-up is measured in weeks to confirm a successful pregnancy and months to confirm a successful birth.

**Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

**Review of Evidence****Systematic Reviews**

A Cochrane review and meta-analysis by Carney et al (2012) identified 31 RCTs evaluating assisted hatching (N=5728).<sup>2</sup> Twelve studies included women with a poor fertility prognosis, 12 studies included women with a good fertility prognosis, and the remaining 7 studies did not report this factor. Fifteen studies used a laser for assisted hatching, 11 used chemical means, and 5 used mechanical means. Live birth rates were reported in 9 studies (n=1921 women). A pooled analysis of data from the 9 studies did not find a statistically significant difference between the groups receiving assisted hatching and a control condition (odds ratio [OR], 1.03; 95% confidence interval [CI], 0.85 to 1.26). The rate of live birth was 313 (31%) of 995 in the assisted hatching group and 282 (30%) of 926 in the control group. All 31 trials reported clinical pregnancy rates. In a meta-analysis of all trials, assisted hatching improved the pregnancy rate, but the estimate for the odds was of marginal statistical significance (OR, 1.13; 95% CI, 1.01 to 1.27).

**Randomized Controlled Trials**

Two RCTs not assessed in the Cochrane review have compared laser-assisted hatching with the standard of care. Shi et al (2016) evaluated 178 patients of advanced maternal age (age range, 35-42 years).<sup>3</sup> There were no statistically significant differences in implantation rates (32.5% in the assisted hatching group vs. 39.3% in the control group) or in clinical pregnancy rates (48.8% in the assisted hatching group vs. 50.4% in the control group; p values not reported). Kanyo et al (2016) assessed 413 women (mean age, 33 years).<sup>4</sup> In the overall study population, there was no statistically significant difference in the clinical pregnancy rate between the assisted hatching group (33.3%) and the control group (27.4%; p=.08). However, in the subgroup of patients ages 38 or older, the clinical pregnancy rate was significantly higher in the assisted hatching group (18.4%) than in the control group (11.4%; p=.03). There was no significant between-group difference in the clinical pregnancy rate among women younger than 38 years old. Neither trial reported live birth rates.

### Retrospective Studies

Knudtson et al (2017), in a retrospective cohort study, analyzed live birth rates in women who underwent first-cycle, autologous frozen embryo transfer.<sup>5</sup> From data reported between 2004 and 2013 to the Society for Assisted Reproductive Technology Clinic Outcomes Reporting System, 151,533 cycles were identified, 70,738 (46.7%) with assisted hatching and 80,795 (53.3%) without. Assisted hatching had a significantly lower live birth rate (34.2%) than nonassisted hatching (35.4%;  $p < .001$ ). Also, older patients (age  $\geq 38$  years) who received assisted hatching were associated with lower live birth rates ( $p \leq .05$ ). Results were similar in a 2019 study by McLaughlin et al that analyzed Society for Assisted Reproductive Technology Clinic Outcomes Reporting System data from 2007 to 2015 comparing assisted hatching ( $n=48,858$ ) with no assisted hatching ( $n=103,413$ ) in women undergoing first cycle, fresh IVF.<sup>6</sup> The study found assisted hatching associated with a significantly lower live birth rate than no assisted hatching (39.2% versus 43.9%; rate difference- 4.7%, 95% CI, -0.053 to -0.040).

Kissin et al (2014) retrospectively reviewed data on assisted hatching in the U.S. from 2000 to 2010.<sup>7</sup> Data were taken from the Centers for Disease Control and Prevention's National Assisted Reproductive Technology Surveillance System. The analysis of outcomes was limited to fresh autologous IVF cycles for which a transfer was performed on day 3 or 5. For the total patient population ( $N=536,852$ ), rates of implantation, clinical pregnancy, and live births were significantly lower when assisted hatching was used. For example, the live birth rate was 28.3% with assisted hatching and 36.5% without (adjusted odds ratio [AOR], 0.75; 95% CI, 0.70 to 0.81). Moreover, the rate of miscarriage was significantly higher when assisted hatching was used (18.0% vs. 13.5%; AOR=1.43; 95% CI, 1.34 to 1.52).

### Section Summary: Assisted Hatching

The available literature has generally not found better outcomes with assisted hatching than with standard of care. A 2012 Cochrane review of heterogeneous RCTs found that clinical pregnancy rates, but not the live birth rates, improved with assisted hatching. In subsequent RCTs, laser-assisted hatching did not improve the clinical pregnancy rate but, in 1 study, there was a higher rate of clinical pregnancy in the subgroup of women 38 years or older. In addition, analyses of a large national database found better outcomes (e.g., clinical pregnancy and live birth rates) when assisted hatching was not used.

### Embryo Co-Culture

In routine IVF procedures, the embryo is transferred to the uterus on day 2 or 3 of development, when it has between 4 and 8 cells. Embryo co-culture techniques, used successfully in domestic animals, represent an effort to improve the culture media for embryos such that a greater proportion of embryos will reach the blastocyst stage, in an attempt to improve implantation and pregnancy rates. In addition, if co-culture results in a higher implantation rate, fewer embryos could be transferred in each cycle, decreasing the incidence of multiple pregnancies. A variety of co-culture techniques have been investigated involving the use of feeder cell layers derived from a range of tissues, including the use of human reproductive tissues (i.e., oviducts) to nonhuman cells (i.e., fetal bovine uterine or oviduct cells) to established cell lines (i.e., Vero cells or bovine kidney cells).

### Clinical Context and Therapy Purpose

The purpose of IVF with embryo co-culture in patients with infertility is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does IVF with embryo co-culture to treat infertility improve the net health outcome?

The following PICO was used to select literature to inform this review.

**Patients**

The relevant population of interest is patients who are infertile.

**Interventions**

The therapy being considered is IVF with embryo co-culture.

**Comparators**

The following practice is currently being used to make decisions about infertility: IVF without embryo co-culture.

**Outcomes**

The general outcomes of interest are live birth rates and infant abnormalities. Follow-up is measured to confirm successful pregnancy up to successful birth.

**Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

**Review of Evidence****Randomized Controlled Trials**

Currently, no standardized method of co-culture has emerged, and clinical trials have generally not found that co-culture is associated with improved implantation or pregnancy rates.<sup>8,9,10,11,12,13</sup> For example, Wetzels et al (1998) reported on an RCT that assigned IVF treatments to co-culture with human fibroblasts or no culture.<sup>13</sup> Patients in the 2 groups were stratified by age (older or younger than 36 years) and prior IVF attempts (yes vs. no). The trialists reported that fibroblast co-culture did not affect the implantation or pregnancy rates. More recently, Ohl et al (2015) reported on a novel co-culture technique involving autologous endometrial cell co-culture.<sup>14</sup> In an interim analysis of 320 patients, the clinical pregnancy rate per embryo transfer was significantly higher in the co-culture group (53.4%) than in the control group (37.3%;  $p=.025$ ).

**Section Summary: Embryo Co-Culture**

There is no standardized method of co-culture, and few clinical trials have evaluated outcomes. Most have not found improved implantation or pregnancy rates after co-culture. A 2015 RCT has reported on a novel co-culture method, and an interim analysis of the trial found a higher clinical pregnancy rate with co-culture than with standard practice control group. Additional studies are needed to evaluate this novel co-culture technique. No studies have reported on the impact of co-culture on live birth rates.

**Blastocyst Transfer**

The most common days for embryo transfer in the clinical IVF setting are day 3 or day 5. Embryo transfer at the blastocyst stage on day 5 continues to be less common than cleavage-stage transfer on day 3. First introduced in clinical practice in 2005, use of blastocyst transfer is increasing in clinical practice. The rationale and reported advantages for blastocyst transfer are: higher implantation and clinical pregnancy rates, a more viable option for limiting to single embryo transfer, more appropriate endometrium-embryo synchronicity, optimization of embryo selection due to embryo development progression, and decreased potential for embryo trauma with biopsy obtained for preimplantation genetic testing. Advances in cell culture techniques and embryology assessments have facilitated increased use of blastocyst transfer and research into the technique. Critics of blastocyst transfer have raised concerns about the limitation on the

number of available embryos for transfer once the cleavage-stage is passed; critics also cite concerns due to uncertainties about the effects of the culture microenvironment, as well as early indicators of a higher rate of adverse pregnancy outcomes.

### **Clinical Context and Therapy Purpose**

The purpose of IVF with blastocyst transfer in patients with infertility is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does IVF with blastocyst transfer treat infertility and improve the net health outcome?

The following PICO was used to select literature to inform this review.

### **Populations**

The relevant population of interest is patients who are infertile.

### **Interventions**

The therapy being considered is IVF with blastocyst transfer.

### **Comparators**

The following practice is currently being used to make decisions about infertility: IVF without cleavage-stage transfer.

### **Outcomes**

The general outcomes of interest are live birth rates and infant abnormalities. Follow-up is measured to confirm successful pregnancy up to successful birth.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

### **Review of Evidence**

#### **Systematic Reviews**

Several systematic reviews of studies comparing outcomes associated with blastocyst-stage transfer with those of earlier stage transfer have been published. Only the Cochrane review by Glujovsky et al (2012) included RCTs.<sup>25</sup> The authors identified 23 RCTs, 12 of which reported on the rates of live births per couple. A pooled analysis of these trials found a significantly higher live birth rate with blastocyst transfer (292/751 [39%]) than with cleavage-stage transfer (237/759 [31%]). The odds for live birth was 1.40 (95% CI, 1.13 to 1.74). There was no significant difference in the rate of multiple pregnancies between the 2 treatment groups (16 RCTs; OR, 0.92; 95% CI, 0.71 to 1.19). In addition, there was no significant difference in the miscarriage rate (14 RCTs; OR, 1.14; 95% CI, 0.84 to 1.55). Glujovsky et al (2016), in their updated Cochrane review, placed more emphasis on whether blastocyst-stage (day 5-6) embryo transfers improved the live birth rates, and other associated outcomes, compared with cleavage-stage (day 2-3) embryo transfers.<sup>26</sup> Data from 4 new studies, 3 of which were published studies,<sup>27,28,29</sup> resulted in a total of 27 parallel-design RCTs that included 4031 couples or women. The data from a fourth study was only available in abstract form and reported on outcomes from a multicenter trial comparing blastocyst with day 2-3 transfer in intracytoplasmic sperm injection (ICSI) cycles for male factor infertility (MFI). There were no exclusions from the 2012 review. The live birth rate following fresh transfer was higher in the blastocyst transfer group (OR, 1.48; 95% CI, 1.20 to 1.82; 13 RCTs, 1630



women,  $I^2=45\%$ , low-quality evidence). There was no evidence of a difference between groups in rates of cumulative pregnancy per couple following fresh and frozen-thawed transfer after 1 oocyte retrieval (OR, 0.89; 95% CI, 0.64 to 1.22; 5 RCTs, 632 women,  $I^2=71\%$ , very low-quality evidence). The clinical pregnancy rate was also higher in the blastocyst transfer group, following fresh transfer (OR, 1.30; 95% CI, 1.14 to 1.47; 27 RCTs, 4031 women,  $I^2=56\%$ , moderate-quality evidence). Embryo freezing rates were lower in the blastocyst transfer group (OR, 0.48; 95% CI, 0.40 to 0.57; 14 RCTs, 2292 women,  $I^2=84\%$ , low-quality evidence). Failure to transfer any embryos was higher in the blastocyst transfer group (OR, 2.50; 95% CI, 1.76 to 3.55; 17 RCTs, 2577 women,  $I^2=36\%$ , moderate-quality evidence). The data for rates of multiple pregnancy and miscarriage were incomplete in 70% of the trials and limit conclusions concerning the following findings. There was no evidence of a difference between the groups in rates of multiple pregnancies (OR, 1.05, 95% CI, 0.83 to 1.33; 19 RCTs, 3019 women,  $I^2=30\%$ , low-quality evidence) or miscarriages (OR, 1.15, 95% CI, 0.88 to 1.50; 18 RCTs, 2917 women,  $I^2=0\%$ , low-quality evidence). Reviewers reported that the main limitation of the RCTs assessed was a high-risk of bias, which was associated with failure to describe acceptable methods of randomization and unclear or high-risk of attrition bias.

Maheshwari et al (2013) identified 8 observational studies analyzing singleton births following embryo transfer at the blastocyst or cleavage stage and reporting obstetric and/or perinatal outcomes.<sup>30</sup> Meta-analysis of 6 studies found a significantly higher rate of preterm delivery at less than 37 weeks after blastocyst-stage transfer compared with cleavage-stage transfer (relative risk, 1.27; 95% CI, 1.22 to 1.31); the absolute increase in risk was 2% (95% CI, 1% to 4%). Other pooled analyses of 2 to 3 studies each did not find significantly increased rates of low birth weight less than 1500 grams, congenital anomalies, or perinatal mortality following blastocyst-stage versus cleavage-stage embryo transfer.

### Observational Studies

A retrospective cohort study by Kallen et al (2010) reported on risks associated with blastocyst transfer.<sup>31</sup> Data were taken from the Swedish Medical Birth Register. There were 1311 infants born after blastocyst transfer and 12,562 born after cleavage-stage transfer. There were no significant differences in the rates of multiple births (10% after blastocyst transfer vs. 8.9% after cleavage-stage transfer). Among singleton births, the rate of preterm birth (<32 weeks) was 1.7% (18/1071) in the blastocyst transfer group and 1.35% (142/10513) in the cleavage-stage transfer group. In a multivariate analysis controlling for year of birth, maternal age, parity, smoking habits, and body mass index, the AOR was 1.44 (95% CI, 0.87 to 2.40). The rate of low birth weight singletons (<1500 g or <2500 g) did not differ significantly between the blastocyst transfer group and the cleavage-stage transfer group. There was a significantly higher rate of relatively severe congenital malformation (e.g., spina bifida, cardiovascular defects, cleft palate) after blastocyst transfer (61/1311 [4.7%]) than after cleavage-stage transfer (509/12,562 [4.1%]; AOR, 1.33; 95% CI, 1.01 to 1.75). The groups did not differ significantly in their rates of low Appearance, Pulse, Grimace, Activity and Respiration scores, intracranial hemorrhage rates, respiratory diagnoses, or cardiovascular malformations. Respiratory diagnoses were given to 94 (7.2%) of 1311 infants born after blastocyst transfer and to 774 (6.2%) of 12,562 after cleavage-stage transfer (OR, 1.15; 95% CI, 0.90 to 1.47).

Ginström Ernstad et al (2016) published another retrospective registry cohort study using data crosslinked across the Swedish Medical Birth Register, the Register of Birth Defects, and the National Patient Register.<sup>32</sup> All singleton deliveries after blastocyst transfer in Sweden from 2002 through 2013 were compared with deliveries after cleavage-stage transfer and deliveries after spontaneous conception. There were 4819 singletons born after blastocyst transfer, 25,747 after cleavage-stage transfer, and 1,196,394 after spontaneous conception. Singletons born after blastocyst transfer had no increased risk of birth defects compared with singletons born after the cleavage-stage transfer (AOR, 0.94; 95% CI, 0.79 to 1.13) or spontaneous conception (AOR, 1.09; 95% CI, 0.92 to 1.28). Perinatal mortality was higher in the blastocyst group versus the cleavage-stage group (AOR, 1.61; 95% CI, 1.14 to 2.29). When comparing singletons born after blastocyst transfer with singletons born after spontaneous conception, a higher risk of preterm birth (<37

weeks) was detected (AOR, 1.17; 95% CI, 1.05 to 1.31). Singletons born after blastocyst transfer had a lower rate of low birthweight (AOR, 0.83; 95% CI, 0.71 to 0.97) than singletons born after cleavage-stage transfer. The rate of being small for gestational age was also lower in singletons born after blastocyst transfer than after both cleavage-stage conception (AOR, 0.71; 95% CI, 0.56 to 0.88) and spontaneous conception (AOR, 0.70; 95% CI, 0.57 to 0.87). The risks of placenta previa and placental abruption were higher in pregnancies after blastocyst transfer than in pregnancies after cleavage stage (AOR, 2.08; 95% CI, 1.70 to 2.55; AOR, 1.62; 95% CI, 1.15 to 2.29, respectively) and after spontaneous conception (AOR, 6.38; 95% CI, 5.31 to 7.66; AOR, 2.31; 95% CI, 1.70 to 3.13, respectively).

A 2020 study by Spangmose et al focused on the comparative obstetric and perinatal harms of blastocyst transfer versus cleavage-stage transfer.<sup>33</sup> The study used combined data from Norway, Sweden and Denmark from 56,557 singleton pregnancies. Women undergoing blastocyst transfer were significantly more likely to have placenta previa (AOR, 2.11; 95% CI, 1.76 to 2.52) and marginally more likely to have a Cesarean section (AOR, 1.09; 95% CI, 1.01 to 1.18) relative to cleavage-stage transfer. Risk of labor induction was slightly lower with blastocyst transfer (AOR, 0.91; 95% CI, 0.83 to 0.99). There were no clear differences in perinatal outcomes, apart from risk of preterm birth which was slightly higher with blastocyst transfer (AOR, 1.14; 95% CI, 1.01 to 1.29).

### **Section Summary: Blastocyst Transfer**

An updated 2016 Cochrane review of 27 RCTs compared the effectiveness of blastocyst transfers with cleavage-stage transfers. The primary outcomes of live birth and cumulative clinical pregnancy rates were higher with fresh blastocyst transfer. There were no differences between groups in multiple pregnancies or early pregnancy loss (miscarriage). The main limitation of the RCTs evaluated in the Cochrane review was a high risk of bias associated with failure to describe acceptable methods of randomization and unclear or high risk of attrition bias. Differences in outcomes with the use of cryopreserved blastocysts and cleavage-stage embryos have been reported, and the mechanisms are not well understood. There are conflicting reports from retrospective studies on the incidence of pregnancy and neonatal adverse outcomes, including low birth weight and increased congenital anomalies.

### **Intracytoplasmic Sperm Injection for Male Factor Infertility**

Intracytoplasmic sperm injection is performed in cases of MFI when either insufficient numbers of sperm, abnormal sperm morphology, or poor sperm motility preclude unassisted IVF. Fertilization rates represent an intermediate outcome; the final outcome is the number of pregnancies per initiated cycle or per embryo transfer.

### **Clinical Context and Therapy Purpose**

The purpose of IVF with ICSI in patients with MFI is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does IVF with ICSI treat MFI and improve the net health outcome?

The following PICO was used to select literature to inform this review.

### **Populations**

The relevant population of interest is men with MFI.

### **Interventions**

The therapy being considered is IVF with ICSI.

### **Comparators**

The following practice is currently being used to make decisions about infertility: IVF without ICSI.

### Outcomes

The general outcomes of interest are live birth rates and infant abnormalities. Follow-up is measured in months to confirm successful birth.

### Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

### Review of Evidence

#### Case Series

The number of pregnancies per cycle and per embryo transfers, reported in relatively large series published in the mid-1990s, ranged between 45% and 50%.[34,35,36,37,38](#). At the time, those rates were very competitive with those of standard IVF.

More recently, Borges et al (2017) retrospectively analyzed ICSI outcomes for patients with MFI compared with isolated tubal factor infertility (TFI).[39](#). Nine hundred twenty-two ICSI cycles (743 for MFI, 179 for TFI) performed between 2010 and 2016 were identified. No significant differences were observed between the groups for rates of implantation (MFI=35.5% vs. TFI=32%;  $p=.34$ ), pregnancy (MFI=46.9% vs. TFI=40.9%;  $p=.184$ ), and miscarriage (MFI 10.3% vs. TFI 10.6%,  $p=.572$ ); rates remained similar even after women were stratified into groups by age ( $\leq 35$  years: MFI=531 vs. TFI=112;  $>35$  years: MFI=212 vs. TFI=67). The study was limited by its retrospective design and by the fact that MFI severity could not be determined because patients were not categorized by diagnosis.

Boulet et al (2015) published a large retrospective analysis of the outcomes following ICSI versus standard IVF (data captured from the Centers for Disease Control and Prevention's National Assisted Reproductive Technology Surveillance System from 2008 to 2012).[40](#). During that time, there were data on 494,907 fresh IVF cycles. A total of 74.6% of cycles used ICSI, with 92.9% of the cycles involving MFI and 64.5% of the cycles not. Among couples with MFI, there was a statistically significantly lower rate of implantation after ICSI (25.5%) than after standard IVF (25.6%;  $p=.02$ ); however, this difference between groups was not clinically significant. Rates of clinical intrauterine pregnancy and live birth did not differ significantly between ICSI and standard IVF. In couples without MFI, implantation, clinical pregnancy, and live birth rates were all significantly higher with standard IVF than with ICSI.

#### Adverse Events

A systematic review and meta-analysis by Massaro et al (2015) examined adverse events related to ICSI and standard IVF without ICSI.[41](#). Twenty-two observational studies were included; no RCTs were identified. Meta-analysis of 12 studies found a significantly increased odds of congenital genitourinary malformations in children conceived using ICSI versus standard IVF (pooled OR, 1.27; 95% CI, 1.02 to 1.58;  $p=.04$ ;  $I^2=0$ ). Five studies in this analysis were considered at high-risk of bias, and a pooled analysis of the 4 studies considered at low-risk of bias did not determine whether ICSI was associated with a statistically increased odds of genitourinary malformations.

#### Section Summary: Intracytoplasmic Sperm Injection for Male Factor Infertility

There is a lack of RCTs comparing ICSI with standard IVF. Observational studies have found similar rates of clinical pregnancy and live births after ICSI and standard IVF but those observational studies are subject to limitations (e.g., selection bias). A 2015 meta-analysis of observational studies found a significantly higher rate of congenital genitourinary malformations in children born after ICSI versus IVF, but there was no significant difference when only studies

with low-risk of bias were analyzed. Randomized controlled trials comparing health outcomes after ICSI for MFI with standard IVF would strengthen the evidence base.

### **Cryopreservation of Testicular Tissue in Adult Men With Azoospermia**

Testicular sperm extraction refers to the collection of sperm from testicular tissue in men with azoospermia. Extraction of testicular sperm may be performed during or subsequent to a diagnostic biopsy, specifically for the collection of spermatozoa. Spermatozoa may be isolated immediately and a portion used for an ICSI procedure during oocyte retrieval from the partner, with the remainder cryopreserved. Alternatively, the entire tissue sample can be cryopreserved with portion thawed and sperm isolation performed at subsequent ICSI cycles.

### **Clinical Context and Therapy Purpose**

The purpose of the cryopreservation of testicular tissue as part of ICSI in patients with azoospermia is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does the cryopreservation of testicular tissue as part of ICSI treat azoospermia and improve the net health outcome?

The following PICO was used to select literature to inform this review.

### **Populations**

The relevant population of interest is men who are infertile.

### **Interventions**

The therapy being considered is cryopreservation of testicular tissue as part of ICSI.

### **Comparators**

The following practice is currently being used to make decisions about infertility: IVF without cryopreservation of testicular tissue.

### **Outcomes**

The general outcomes of interest are live birth rates and infant abnormalities. Follow-up is measured in months to confirm successful birth.

### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

### **Review of Evidence**

#### **Case Series**

Testicular tissue extraction appears to be a well-established component of the overall ICSI procedure; cryopreservation of either the isolated sperm or the tissue sample eliminates the need for multiple biopsies to obtain fresh tissue in the event of a failed initial ICSI cycle.<sup>42</sup> However, clinical trials evaluating health outcomes after cryopreservation of testicular tissue in adult men with azoospermia were not identified.

### **Section Summary: Cryopreservation of Testicular Tissue in Adult Men With Azoospermia**

While cryopreservation of testicular tissue in adult men with azoospermia is a well-established component of the ICSI procedure, there is a lack of clinical trials to support this treatment.

### **Cryopreservation of Testicular Tissue in Prepubertal Boys With Cancer**

A potential application of cryopreservation of testicular tissue is its potential to preserve the reproductive capacity in prepubertal boys undergoing cancer chemotherapy; cryopreservation of ejaculate is not an option in these patients.

#### **Clinical Context and Therapy Purpose**

The purpose of the cryopreservation of testicular tissue in prepubertal boys with cancer is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does the cryopreservation of testicular tissue from prepubertal boys with cancer improve the net health outcome?

The following PICO was used to select literature to inform this review.

#### **Populations**

The relevant population of interest is prepubertal boys with cancer.

#### **Interventions**

The therapy being considered is the cryopreservation of testicular tissue.

#### **Comparators**

The following practice is currently being used to make decisions about infertility: no cryopreservation of testicular tissue.

#### **Outcomes**

The general outcomes of interest are live birth rates and infant abnormalities. Follow-up is measured in months to confirm successful birth.

#### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

#### **Review of Evidence**

##### **Modeling Studies**

It has been hypothesized that reimplantation of the frozen-thawed testicular stem cells will reinitiate spermatogenesis or, alternatively, spermatogenesis could be attempted in vitro, using frozen-thaw spermatogonia. While these strategies have been explored in animals, there are inadequate human studies.[43,44](#)

##### **Section Summary: Cryopreservation of Testicular Tissue in Prepubertal Boys With Cancer**

No clinical trials were identified evaluating the safety and efficacy of cryopreservation of testicular tissue in prepubertal boys undergoing cancer therapy.

##### **Potential Adverse Events to Offspring Conceived Via Assisted Reproduction**

Several systematic reviews have addressed the risk of birth defects.[45,46,47,48](#) The review with the most data is that by Hansen et al (2013).[47](#) They examined 45 cohort studies with outcomes in 92,671 infants born following assisted reproduction and 3,870,760 naturally conceived infants. In a pooled analysis, there was a higher risk of birth defects in infants born using reproductive techniques (relative risk, 1.32; 95% CI, 1.24 to 1.42). The risk of birth defects was also elevated

when the analysis was limited to the 6 studies conducted in the U.S. or Canada (relative risk, 1.38; 95% CI, 1.16 to 1.64). Another review, published by Davies et al (2012), included data on 308,974 live births in Australia, 6163 of which used ART.<sup>48</sup> There was a higher rate of birth defects after assisted conception (8.3%) compared with births to fertile women who did not use assisted reproduction (5.8%; unadjusted OR, 1.47; 95% CI, 1.33 to 1.62). The risk of birth defects was still significantly elevated but was lower in an analysis that adjusted for other factors that might increase risk (e.g., maternal age, parity, maternal ethnicity, maternal smoking during pregnancy, socioeconomic status; OR, 1.28; 95% CI, 1.16 to 1.41). A more recent review by Elias et al (2020) identified 14 cohort studies examining neonatal outcomes in ART.<sup>49</sup> The risk of preterm birth was significantly increased among both those undergoing fresh embryo transfer (OR, 1.64; 95% CI, 1.46 to 1.84) and frozen embryo transfer (OR, 1.39; 95% CI, 1.34 to 1.44) compared with spontaneous conceptions. Fresh embryo transfer was also associated with low birth weight (OR, 1.67; 95% CI, 1.52 to 1.85) and small for gestational age (OR, 1.46; 95% CI, 1.11 to 1.92) compared with standard conception while frozen embryo transfer increased the risk of large for gestational age (OR, 1.57; 95% CI, 1.48 to 1.68).

### Summary of Evidence

For individuals who have infertility who receive IVF with assisted hatching, the evidence includes RCTs, a systematic review, and retrospective studies. Randomized controlled trials have not shown that assisted hatching improves clinical pregnancy or live birth rate compared with standard care while large observational studies found assisted hatching associated with worse outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have infertility who receive IVF with embryo co-culture, the evidence includes RCTs and case series. Most clinical trials have not found improved implantation or pregnancy rates after co-culture, and studies have not reported live birth rates. Moreover, co-culture techniques have not been standardized. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have cancer who will undergo treatment that may lead to infertility who receive cryopreservation of ovarian tissue, the evidence includes case series that have reported on the technique as well as pregnancy and live birth rates after transplantation. The technique used has not been standardized, and there is a lack of controlled studies on health outcomes following cryopreservation of ovarian tissue. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have cancer who will undergo treatment that may lead to infertility who receive cryopreservation of oocytes, the evidence includes RCTs and a systematic review assessing the technique in related populations. The systematic review found that fertilization rates ranged from 71% to 79%, and the clinical pregnancy rates per transfer ranged from 36% to 61%. The available studies may not be generalizable to the population of interest, women with cancer. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have infertility who receive IVF with blastocyst transfer, the evidence includes RCTs and meta-analyses. The RCTs and meta-analyses have found that blastocyst transfer is associated with higher live birth rates than cleavage-stage transfer. One retrospective cohort study has reported a significantly higher rate of preterm birth after blastocyst-stage versus cleavage-stage transfer but did not find increased risks of other outcomes such as a low birth rate or perinatal mortality. A retrospective registry review of a similar population reported different findings. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have male factor infertility who receive IVF with ICSI, the evidence includes observational studies and a systematic review. The evidence includes observational studies that

found similar rates of clinical pregnancy and live birth after ICSI and standard IVF, and a meta-analysis of observational studies found a higher rate of genitourinary malformations in children born after ICSI (but only when lower quality studies were included in the analysis). The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have azoospermia who receive cryopreservation of testicular tissue as part of ICSI, the evidence includes no clinical trials. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are prepubertal boys with cancer who receive cryopreservation of testicular tissue, the evidence includes no clinical trials evaluating safety and efficacy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

### **Supplemental Information**

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

### **Clinical Input From Physician Specialty Societies and Academic Medical Centers**

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

In response to requests from Blue Cross Blue Shield Association, input was received from 4 physician specialty societies and 2 academic medical centers in 2012. There was general agreement that intracytoplasmic sperm injection and cryopreservation of testicular tissue in adult men with azoospermia as part of an intracytoplasmic sperm injection procedure may be considered medically necessary. Three of 5 reviewers who responded agreed that co-culture of embryos is considered investigational. In addition, 4 of 5 reviewers did not agree that blastocyst transfer is investigational; these reviewers considered blastocyst transfer to be medically necessary to decrease multiple gestations. Three of 6 reviewers agreed that cryopreservation of ovarian tissue or oocytes is investigational. The other 3 thought that cryopreservation of oocytes, but not ovarian tissue, is medically necessary. Clinical input on other policy statements was more variable.

### **Practice Guidelines and Position Statements**

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

### **American Society for Reproductive Medicine and Society for Assisted Reproductive Technology**

In 2019, the American Society for Reproductive Medicine (ASRM) released a 2019 committee opinion on fertility preservation in patients undergoing gonadotoxic therapy. The committee included several relevant opinions:

- Embryo, oocyte, and ejaculated or testicular sperm cryopreservation remain the principle established modalities for fertility preservation.
- Ovarian tissue cryopreservation is no longer considered experimental and can be used in prepubertal patients or when there is not time for ovarian stimulation.
- Testicular tissue cryopreservation in prepubertal males is still considered experimental and should be conducted under research protocols when no other options are feasible.

ASRM and joint ASRM/Society for Assisted Reproductive Technology (SART) opinions and recommendations on other assisted reproductive technologies are as follows:

- Planned oocyte cryopreservation (OC) for preserving future reproductive potential (2018): The committee states the process is ethical and “serves women’s legitimate interests in reproductive autonomy.” Women who choose OC should be informed of its efficacy, safety, benefits, and risks, and possible long-term health effects on the child. Providers should also provide their clinic’s statistics for successful freeze-thaw and live birth. Women should know that this relatively new technology is still emerging and not all benefits and harms are fully understood.<sup>50</sup>
- Assisted hatching (2014): Assisted hatching should not be used routinely for all patients undergoing IVF.<sup>51</sup>
- Blastocyst transfer (2013; reaffirmed in 2018): “Evidence supports blastocyst transfer in ‘good prognosis’ patients.”<sup>52,50</sup>

In 2020, ASRM developed joint guidelines with the American Urological Association (AUA) for male infertility diagnosis and treatment including recommendations for intracytoplasmic sperm injection.<sup>53,54</sup> Based on expert opinion, patients with low total motile sperm count should be advised to consider IVF with intracytoplasmic sperm injection.<sup>54</sup>

### American College of Obstetricians and Gynecologists

In 2014, the American College of Obstetricians and Gynecologists endorsed the 2013 ASRM-SART joint guidelines on mature OC.<sup>55</sup> The endorsement was affirmed in 2020.

### American Society of Clinical Oncology

In 2018, the American Society of Clinical Oncology updated its 2013 guidelines (with no changes to its recommendations) on fertility preservation for patients with cancer.<sup>56,57</sup> The guidelines included the following recommendations for males and females, respectively.

- "Recommendation 2.1. Sperm cryopreservation: Sperm cryopreservation is effective, and health care providers should discuss sperm banking with postpubertal males receiving cancer treatment.
- Recommendation 2.2. Hormonal gonad protection: Hormonal therapy in men is not successful in preserving fertility. It is not recommended.
- Recommendation 2.3. Other methods to preserve male fertility: Other methods, such as testicular tissue cryopreservation and reimplantation or grafting of human testicular tissue, should be performed only as part of clinical trials or approved experimental protocols..."
- "Recommendation 3.1. Embryo cryopreservation: Embryo cryopreservation is an established fertility preservation method, and it has routinely been used for storing surplus embryos after in vitro fertilization.
- Recommendation 3.2. Cryopreservation of unfertilized oocytes: Cryopreservation of unfertilized oocytes is an option, particularly for patients who do not have a male partner, do not wish to use donor sperm, or have religious or ethical objections to embryo freezing..."

### Agency for Healthcare Research and Quality

Myers et al (2008), in an evidence report conducted for the Agency for Healthcare Research and Quality, evaluated the effectiveness of assisted reproductive technology.<sup>58</sup> They reviewed evidence on the outcomes of interventions used in ovulation induction, superovulation, and IVF for the treatment of infertility. Reviewers concluded that:

"[i]nterventions for which there was sufficient evidence to demonstrate improved pregnancy or live birth rates (pertinent to this evidence review) included: ... (c) ultrasound-guided embryo transfer, and transfer on day 5 post-fertilization, in couples with a good prognosis; and (d) assisted hatching in couples with previous IVF failure. There was insufficient evidence regarding other interventions.

Infertility itself is associated with most of the adverse longer-term outcomes."



Reviewers concluded that "[d]espite the large emotional and economic burden resulting from infertility, there was relatively little high-quality evidence to support the choice of specific interventions." This conclusion was based primarily on studies that had pregnancy rates as the primary endpoint, not live births. In addition, studies used multiple assisted hatching techniques.

#### U.S. Preventive Services Task Force Recommendations

Not applicable.

#### Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

#### Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 1.

**Table 1. Summary of Key Trials**

NCT No.	Trial Name	Planned Enrollment	Completion Date
<b>Ongoing</b>			
<i>Ovarian tissue cryopreservation</i>			
NCT02646384	Ovarian Tissue Freezing For Fertility Preservation In Girls Facing A Fertility Threatening Medical Diagnosis Or Treatment Regimen	100	Dec 2021
NCT02846064	Development of Ovarian Tissue Autograft in Order to Restore Ovarian Function	50	Oct 2022
<i>Oocyte cryopreservation</i>			
NCT04616417	Investigational Oocyte Cryopreservation for Medical and Non Medical Indications	50	Jul 2025
<i>Blastocyst transfer</i>			
NCT03152643	Cumulative Live Birth Rates After Cleavage-stage Versus Blastocyst-stage Embryo Transfer: A Multicenter, Prospective, Randomized Controlled Trial	992	Nov 2021
NCT03764865	Day 3 vs Day 5 Embryo Transfer for Patients With Low Embryo Numbers Going Through in Vitro Fertilization	658	Feb 2024
<i>Intracytoplasmic Sperm Injection</i>			
NCT03298633	Intracytoplasmic Sperm Injection (ICSI) Versus Conventional in Vitro Fertilization (IVF) in Couples With Non-severe Male Infertility: a Randomized Controlled Trial	2346	Jul 2021
<i>Testicular tissue cryopreservation</i>			
NCT02872532	Testicular Tissue Cryopreservation for Fertility Preservation in Males Facing Fertility Threatening Diagnoses or Treatment Regimens	100	Aug 2022
NCT02972801	Testicular Tissue Cryopreservation for Fertility Preservation in Patients Facing Infertility-causing Diseases or Treatment Regimens	500	Jan 202 5

NCT: national clinical trial.

## References

- Centers for Disease Control. What is Assisted Reproductive Technology? [https://www.cdc.gov/art/whatis.html#:~:text=According%20to%20this%20definition%2C%20ART,donating%20them%20to%20another%20woman](https://www.cdc.gov/art/whatis.html#:~:text=According%20to%20this%20definition%2C%20ART,donating%20them%20to%20another%20woman.). Accessed June 28, 2021.
- Carney SK, Das S, Blake D, et al. Assisted hatching on assisted conception (in vitro fertilisation (IVF) and intracytoplasmic sperm injection (ICSI). Cochrane Database Syst Rev. Dec 12 2012; 12: CD001894. PMID 23235584

3. Shi W, Hongwei T, Zhang W, et al. A Prospective Randomized Controlled Study of Laser-Assisted Hatching on the Outcome of First Fresh IVF-ET Cycle in Advanced Age Women. *Reprod Sci*. Oct 2016; 23(10): 1397-401. PMID 27071963
4. Kanyo K, Zeke J, Kriston R, et al. The impact of laser-assisted hatching on the outcome of frozen human embryo transfer cycles. *Zygote*. Oct 2016; 24(5): 742-7. PMID 26957232
5. Knudtson JF, Failor CM, Gelfond JA, et al. Assisted hatching and live births in first-cycle frozen embryo transfers. *Fertil Steril*. Oct 2017; 108(4): 628-634. PMID 28863938
6. McLaughlin JE, Choi BY, Liu Q, et al. Does assisted hatching affect live birth in fresh, first cycle in vitro fertilization in good and poor prognosis patients?. *J Assist Reprod Genet*. Dec 2019; 36(12): 2425-2433. PMID 31713775
7. Kissin DM, Kawwass JF, Monsour M, et al. Assisted hatching: trends and pregnancy outcomes, United States, 2000-2010. *Fertil Steril*. Sep 2014; 102(3): 795-801. PMID 25044084
8. Kervancioglu ME, Saridogan E, Atasu T, et al. Human Fallopian tube epithelial cell co-culture increases fertilization rates in male factor infertility but not in tubal or unexplained infertility. *Hum Reprod*. Jun 1997; 12(6): 1253-8. PMID 9222012
9. Tucker MJ, Morton PC, Wright G, et al. Enhancement of outcome from intracytoplasmic sperm injection: does co-culture or assisted hatching improve implantation rates?. *Hum Reprod*. Nov 1996; 11(11): 2434-7. PMID 8981127
10. Veiga A, Torello MJ, Menezo Y, et al. Use of co-culture of human embryos on Vero cells to improve clinical implantation rate. *Hum Reprod*. Dec 1999; 14 Suppl 2: 112-20. PMID 10690807
11. Wiemer KE, Cohen J, Tucker MJ, et al. The application of co-culture in assisted reproduction: 10 years of experience with human embryos. *Hum Reprod*. Dec 1998; 13 Suppl 4: 226-38. PMID 10091073
12. Rubio C, Simon C, Mercader A, et al. Clinical experience employing co-culture of human embryos with autologous human endometrial epithelial cells. *Hum Reprod*. Dec 2000; 15 Suppl 6: 31-8. PMID 11261481
13. Wetzels AM, Bastiaans BA, Hendriks JC, et al. The effects of co-culture with human fibroblasts on human embryo development in vitro and implantation. *Hum Reprod*. May 1998; 13(5): 1325-30. PMID 9647567
14. Ohl J, de Mouzon J, Nicollet B, et al. Increased pregnancy rate using standardized coculture on autologous endometrial cells and single blastocyst transfer : a multicentre randomized controlled trial. *Cell Mol Biol (Noisy-le-grand)*. Dec 24 2015; 61(8): 79-88. PMID 26718434
15. Tryde Schmidt KL, Yding Andersen C, Starup J, et al. Orthotopic autotransplantation of cryopreserved ovarian tissue to a woman cured of cancer - follicular growth, steroid production and oocyte retrieval. *Reprod Biomed Online*. Apr 2004; 8(4): 448-53. PMID 15149569
16. Oktay K, Buyuk E, Veeck L, et al. Embryo development after heterotopic transplantation of cryopreserved ovarian tissue. *Lancet*. Mar 13 2004; 363(9412): 837-40. PMID 15031026
17. Meirow D, Levron J, Eldar-Geva T, et al. Pregnancy after transplantation of cryopreserved ovarian tissue in a patient with ovarian failure after chemotherapy. *N Engl J Med*. Jul 21 2005; 353(3): 318-21. PMID 15983020
18. Siegel-Itzkovich J. Woman gives birth after receiving transplant of her own ovarian tissue. *BMJ*. Jul 09 2005; 331(7508): 70. PMID 16002876
19. Donnez J, Dolmans MM, Demylle D, et al. Livebirth after orthotopic transplantation of cryopreserved ovarian tissue. *Lancet*. Oct 2004; 364(9443): 1405-10. PMID 15488215
20. Kim SS, Battaglia DE, Soules MR. The future of human ovarian cryopreservation and transplantation: fertility and beyond. *Fertil Steril*. Jun 2001; 75(6): 1049-56. PMID 11384626
21. Lobo RA. Potential options for preservation of fertility in women. *N Engl J Med*. Jul 07 2005; 353(1): 64-73. PMID 16000356
22. Johnson J, Patrizio P. Ovarian cryopreservation strategies and the fine control of ovarian follicle development in vitro. *Ann N Y Acad Sci*. Mar 2011; 1221: 40-6. PMID 21401628
23. Practice Committees of the American Society for Reproductive Medicine and the Society for Assisted Reproductive Technology. Mature oocyte cryopreservation: a guideline. *Fertil Steril*. Jan 2013; 99(1): 37-43. PMID 23083924

24. Levi Setti PE, Albani E, Morengi E, et al. Comparative analysis of fetal and neonatal outcomes of pregnancies from fresh and cryopreserved/thawed oocytes in the same group of patients. *Fertil Steril*. Aug 2013; 100(2): 396-401. PMID 23608156
25. Glujovsky D, Blake D, Farquhar C, et al. Cleavage stage versus blastocyst stage embryo transfer in assisted reproductive technology. *Cochrane Database Syst Rev*. Jul 11 2012; (7): CD002118. PMID 22786480
26. Glujovsky D, Farquhar C, Quinteiro Retamar AM, et al. Cleavage stage versus blastocyst stage embryo transfer in assisted reproductive technology. *Cochrane Database Syst Rev*. Jun 30 2016; (6): CD002118. PMID 27357126
27. Azimineko E, Mohseni Salehi MS, Kalantari V, et al. Pregnancy outcome after blastocyst stage transfer comparing to early cleavage stage embryo transfer. *Gynecol Endocrinol*. 2015; 31(11): 880-4. PMID 26437606
28. Fernandez-Shaw S, Cercas R, Brana C, et al. Ongoing and cumulative pregnancy rate after cleavage-stage versus blastocyst-stage embryo transfer using vitrification for cryopreservation: impact of age on the results. *J Assist Reprod Genet*. Feb 2015; 32(2): 177-84. PMID 25403438
29. Kaur P, Swarankar ML, Maheshwari M, et al. A comparative study between cleavage stage embryo transfer at day 3 and blastocyst stage transfer at day 5 in in-vitro fertilization/intra-cytoplasmic sperm injection on clinical pregnancy rates. *J Hum Reprod Sci*. Jul 2014; 7(3): 194-7. PMID 25395745
30. Maheshwari A, Kalampokas T, Davidson J, et al. Obstetric and perinatal outcomes in singleton pregnancies resulting from the transfer of blastocyst-stage versus cleavage-stage embryos generated through in vitro fertilization treatment: a systematic review and meta-analysis. *Fertil Steril*. Dec 2013; 100(6): 1615-21.e1-10. PMID 24083875
31. Kallen B, Finnstrom O, Lindam A, et al. Blastocyst versus cleavage stage transfer in in vitro fertilization: differences in neonatal outcome?. *Fertil Steril*. Oct 2010; 94(5): 1680-3. PMID 20137785
32. Ginstrom Erntad E, Bergh C, Khatibi A, et al. Neonatal and maternal outcome after blastocyst transfer: a population-based registry study. *Am J Obstet Gynecol*. Mar 2016; 214(3): 378.e1-378.e10. PMID 26928152
33. Spangmose AL, Ginstrom Erntad E, Malchau S, et al. Obstetric and perinatal risks in 4601 singletons and 884 twins conceived after fresh blastocyst transfers: a Nordic study from the CoNARTaS group. *Hum Reprod*. Apr 28 2020; 35(4): 805-815. PMID 32294185
34. Van Steirteghem AC, Liu J, Joris H, et al. Higher success rate by intracytoplasmic sperm injection than by subzonal insemination. Report of a second series of 300 consecutive treatment cycles. *Hum Reprod*. Jul 1993; 8(7): 1055-60. PMID 8408486
35. Palermo G, Joris H, Devroey P, et al. Pregnancies after intracytoplasmic injection of single spermatozoon into an oocyte. *Lancet*. Jul 04 1992; 340(8810): 17-8. PMID 1351601
36. Palermo G, Joris H, Derde MP, et al. Sperm characteristics and outcome of human assisted fertilization by subzonal insemination and intracytoplasmic sperm injection. *Fertil Steril*. Apr 1993; 59(4): 826-35. PMID 8458504
37. Van Steirteghem AC, Nagy Z, Joris H, et al. High fertilization and implantation rates after intracytoplasmic sperm injection. *Hum Reprod*. Jul 1993; 8(7): 1061-6. PMID 8408487
38. Fishel S, Timson J, Lisi F, et al. Micro-assisted fertilization in patients who have failed subzonal insemination. *Hum Reprod*. Mar 1994; 9(3): 501-5. PMID 8006142
39. Borges E, Zanetti BF, Braga DPAF, et al. Overcoming male factor infertility with intracytoplasmic sperm injection. *Rev Assoc Med Bras (1992)*. Aug 2017; 63(8): 697-703. PMID 28977108
40. Boulet SL, Mehta A, Kissin DM, et al. Trends in use of and reproductive outcomes associated with intracytoplasmic sperm injection. *JAMA*. Jan 20 2015; 313(3): 255-63. PMID 25602996
41. Massaro PA, MacLellan DL, Anderson PA, et al. Does intracytoplasmic sperm injection pose an increased risk of genitourinary congenital malformations in offspring compared to in vitro fertilization? A systematic review and meta-analysis. *J Urol*. May 2015; 193(5 Suppl): 1837-42. PMID 25813561

42. Dafopoulos K, Griesinger G, Schultze-Mosgau A, et al. Cumulative pregnancy rate after ICSI with cryopreserved testicular tissue in non-obstructive azoospermia. *Reprod Biomed Online*. Apr 2005; 10(4): 461-6. PMID 15901452
43. Hovatta O. Cryobiology of ovarian and testicular tissue. *Best Pract Res Clin Obstet Gynaecol*. Apr 2003; 17(2): 331-42. PMID 12758103
44. Tournaye H, Goossens E, Verheyen G, et al. Preserving the reproductive potential of men and boys with cancer: current concepts and future prospects. *Hum Reprod Update*. Nov-Dec 2004; 10(6): 525-32. PMID 15319377
45. Kettner LO, Henriksen TB, Bay B, et al. Assisted reproductive technology and somatic morbidity in childhood: a systematic review. *Fertil Steril*. Mar 2015; 103(3): 707-19. PMID 25624193
46. Farhi A, Reichman B, Boyko V, et al. Congenital malformations in infants conceived following assisted reproductive technology in comparison with spontaneously conceived infants. *J Matern Fetal Neonatal Med*. Aug 2013; 26(12): 1171-9. PMID 23451839
47. Hansen M, Kurinczuk JJ, Milne E, et al. Assisted reproductive technology and birth defects: a systematic review and meta-analysis. *Hum Reprod Update*. Jul-Aug 2013; 19(4): 330-53. PMID 23449641
48. Davies MJ, Moore VM, Willson KJ, et al. Reproductive technologies and the risk of birth defects. *N Engl J Med*. May 10 2012; 366(19): 1803-13. PMID 22559061
49. Elias FTS, Weber-Adrian D, Pudwell J, et al. Neonatal outcomes in singleton pregnancies conceived by fresh or frozen embryo transfer compared to spontaneous conceptions: a systematic review and meta-analysis. *Arch Gynecol Obstet*. Jul 2020; 302(1): 31-45. PMID 32445067
50. Practice Committee of the American Society for Reproductive Medicine and Practice Committee of the Society of Assisted Reproductive Technology. Blastocyst culture and transfer in a clinically assisted reproduction: a committee opinion. *Fertil Steril*. 2018;110(7):1246-1252.
51. Pfeifer S, Lobo R, Sokol R, et al. Role of assisted hatching in in vitro fertilization: a guideline. *Fertil Steril*. Aug 2014; 102(2): 348-51. PMID 24951365
52. Practice Committee of American Society for Reproductive Medicine. Blastocyst culture and transfer in clinical-assisted reproduction. *Fertil Steril*. Nov 2008; 90(5 Suppl): S174-7. PMID 19007621
53. Schlegel PN, Sigman M, Collura B, et al. Diagnosis and Treatment of Infertility in Men: AUA/ASRM Guideline Part I. *J Urol*. Jan 2021; 205(1): 36-43. PMID 33295257
54. Schlegel PN, Sigman M, Collura B, et al. Diagnosis and Treatment of Infertility in Men: AUA/ASRM Guideline PART II. *J Urol*. Jan 2021; 205(1): 44-51. PMID 33295258
55. ACOG: Committee Opinion No. 584: oocyte cryopreservation. *Obstet Gynecol*. Jan 2014; 123(1): 221-222. PMID 24463693
56. Loren AW, Mangu PB, Beck LN, et al. Fertility preservation for patients with cancer: American Society of Clinical Oncology clinical practice guideline update. *J Clin Oncol*. Jul 01 2013; 31(19): 2500-10. PMID 23715580
57. Oktay K, Harvey BE, Partridge AH, et al. Fertility Preservation in Patients With Cancer: ASCO Clinical Practice Guideline Update. *J Clin Oncol*. Jul 01 2018; 36(19): 1994-2001. PMID 29620997
58. Myers ER, McCrory DC, Mills AA, et al. Effectiveness of assisted reproductive technology (Evidence Report/Technology Assessment No. 167). Rockville, MD: Agency for Healthcare Research and Quality; 2008.
59. Blue Cross Blue Shield Association. Medical Policy Reference Manual, No. 4.02.04 (August 2021).

## Documentation for Clinical Review

### Please provide the following documentation:

- History and physical and/or consultation notes including:
  - Reason for cryopreservation if applicable (e.g., cancer with plans for radiation or chemotherapy)

- Previous history of fertility/infertility
- Previous treatment plan and response
- Previous procedures to address infertility
- Request for procedure per ongoing treatment plan
- Laboratory report including: specific name and test requested

**Post Service (in addition to the above, please include the following):**

- Operative/procedure notes (if applicable)

## Coding

*This Policy relates only to the services or supplies described herein. Benefits may vary according to product design; therefore, contract language should be reviewed before applying the terms of the Policy.*

*The following codes are included below for informational purposes. Inclusion or exclusion of a code(s) does not constitute or imply member coverage or provider reimbursement policy. Policy Statements are intended to provide member coverage information and may include the use of some codes for clarity. The Policy Guidelines section may also provide additional information for how to interpret the Policy Statements and to provide coding guidance in some cases.*

Type	Code	Description
CPT®	0058T	Cryopreservation; reproductive tissue, ovarian <b>(Deleted code effective 1/1/2021)</b>
	0255U	Andrology (infertility), sperm-capacitation assessment of ganglioside GM1 distribution patterns, fluorescence microscopy, fresh or frozen specimen, reported as percentage of capacitated sperm and probability of generating a pregnancy score <b>(Code effective 10/1/2021)</b>
	54500	Biopsy of testis, needle (separate procedure)
	54505	Biopsy of testis, incisional (separate procedure)
	54800	Biopsy of epididymis, needle
	55400	Vasovasostomy, vasovasorrhaphy
	55870	Electroejaculation
	58321	Artificial insemination; intra-cervical
	58322	Artificial insemination; intra-uterine
	58323	Sperm washing for artificial insemination
	58970	Follicle puncture for oocyte retrieval, any method
	58974	Embryo transfer, intrauterine
	58976	Gamete, zygote, or embryo intrafallopian transfer, any method
	89240	Unlisted miscellaneous pathology test
	89250	Culture of oocyte(s)/embryo(s), less than 4 days;
	89251	Culture of oocyte(s)/embryo(s), less than 4 days; with co-culture of oocyte(s)/embryos
	89253	Assisted embryo hatching, microtechniques (any method)
	89254	Oocyte identification from follicular fluid
	89255	Preparation of embryo for transfer (any method)
	89257	Sperm identification from aspiration (other than seminal fluid)
89258	Cryopreservation; embryo(s)	
89259	Cryopreservation; sperm	
89260	Sperm isolation; simple prep (e.g., sperm wash and swim-up) for insemination or diagnosis with semen analysis	

Type	Code	Description
	89261	Sperm isolation; complex prep (e.g., Percoll gradient, albumin gradient) for insemination or diagnosis with semen analysis
	89264	Sperm identification from testis tissue, fresh or cryopreserved
	89268	Insemination of oocytes
	89272	Extended culture of oocyte(s)/embryo(s), 4-7 days
	89280	Assisted oocyte fertilization, microtechnique; less than or equal to 10 oocytes
	89281	Assisted oocyte fertilization, microtechnique; greater than 10 oocytes
	89335	Cryopreservation, reproductive tissue, testicular
	89337	Cryopreservation, mature oocyte(s)
	89342	Storage (per year); embryo(s)
	89343	Storage (per year); sperm/semen
	89344	Storage (per year); reproductive tissue, testicular/ovarian
	89346	Storage (per year); oocyte(s)
	89352	Thawing of cryopreserved; embryo(s)
	89353	Thawing of cryopreserved; sperm/semen, each aliquot
	89354	Thawing of cryopreserved; reproductive tissue, testicular/ovarian
	89356	Thawing of cryopreserved; oocytes, each aliquot
	89398	Unlisted reproductive medicine laboratory procedure
HCPCS	S4011	In vitro fertilization; including but not limited to identification and incubation of mature oocytes, fertilization with sperm, incubation of embryo(s), and subsequent visualization for determination of development
	S4013	Complete cycle, gamete intrafallopian transfer (GIFT), case rate
	S4014	Complete cycle, zygote intrafallopian transfer (ZIFT), case rate
	S4015	Complete in vitro fertilization cycle, not otherwise specified, case rate
	S4016	Frozen in vitro fertilization cycle, case rate
	S4017	Incomplete cycle, treatment cancelled prior to stimulation, case rate
	S4018	Frozen embryo transfer procedure cancelled before transfer, case rate
	S4020	In vitro fertilization procedure cancelled before aspiration, case rate
	S4021	In vitro fertilization procedure cancelled after aspiration, case rate
	S4022	Assisted oocyte fertilization, case rate
	S4023	Donor egg cycle, incomplete, case rate
	S4025	Donor services for in vitro fertilization (sperm or embryo), case rate
	S4026	Procurement of donor sperm from sperm bank
	S4027	Storage of previously frozen embryos
	S4028	Microsurgical epididymal sperm aspiration (MESA)
	S4030	Sperm procurement and cryopreservation services; initial visit
	S4031	Sperm procurement and cryopreservation services; subsequent visit
	S4035	Stimulated intrauterine insemination (IUI), case rate
	S4037	Cryopreserved embryo transfer, case rate
	S4040	Monitoring and storage of cryopreserved embryos, per 30 days
S4042	Management of ovulation induction (interpretation of diagnostic tests and studies, nonface-to-face medical management of the patient), per cycle	

## Policy History

This section provides a chronological history of the activities, updates and changes that have occurred with this Medical Policy.

Effective Date	Action
08/31/2015	BCBSA Medical Policy adoption
11/01/2016	Policy revision without position change
10/01/2017	Policy revision without position change
10/01/2018	Policy revision without position change
12/01/2019	Policy revision without position change
03/01/2020	Coding update
11/01/2020	Annual review. No change to policy statement. Literature review updated.
01/01/2021	Coding update
11/01/2021	Annual review. No change to policy statement. Literature review updated.

## Definitions of Decision Determinations

**Medically Necessary:** Services that are Medically Necessary include only those which have been established as safe and effective, are furnished under generally accepted professional standards to treat illness, injury or medical condition, and which, as determined by Blue Shield, are: (a) consistent with Blue Shield medical policy; (b) consistent with the symptoms or diagnosis; (c) not furnished primarily for the convenience of the patient, the attending Physician or other provider; (d) furnished at the most appropriate level which can be provided safely and effectively to the patient; and (e) not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of the Member's illness, injury, or disease.

**Investigational/Experimental:** A treatment, procedure, or drug is investigational when it has not been recognized as safe and effective for use in treating the particular condition in accordance with generally accepted professional medical standards. This includes services where approval by the federal or state governmental is required prior to use, but has not yet been granted.

**Split Evaluation:** Blue Shield of California/Blue Shield of California Life & Health Insurance Company (Blue Shield) policy review can result in a split evaluation, where a treatment, procedure, or drug will be considered to be investigational for certain indications or conditions, but will be deemed safe and effective for other indications or conditions, and therefore potentially medically necessary in those instances.

## Prior Authorization Requirements (as applicable to your plan)

Within five days before the actual date of service, the provider must confirm with Blue Shield that the member's health plan coverage is still in effect. Blue Shield reserves the right to revoke an authorization prior to services being rendered based on cancellation of the member's eligibility. Final determination of benefits will be made after review of the claim for limitations or exclusions.

Questions regarding the applicability of this policy should be directed to the Prior Authorization Department at (800) 541-6652, or the Transplant Case Management Department at (800) 637-2066 ext. 3507708 or visit the provider portal at [www.blueshieldca.com/provider](http://www.blueshieldca.com/provider).

*Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. Blue Shield of California may consider published peer-reviewed scientific literature, national guidelines, and local standards of practice in developing its medical policy. Federal and state law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence*

#### **4.02.04 Reproductive Techniques**

Page 24 of 25

*over medical policy and must be considered first in determining covered services. Member contracts may differ in their benefits. Blue Shield reserves the right to review and update policies as appropriate.*



**Appendix A**

POLICY STATEMENT (No changes)	
BEFORE	AFTER
<p><b>Reproductive Techniques 4.02.04</b></p> <p><b>Policy Statement:</b> The following reproductive techniques may be considered <b>medically necessary</b> for <b>any</b> of the following:</p> <ol style="list-style-type: none"> <li>I. Blastocyst transfer</li> <li>II. Cryopreservation of testicular tissue in adult men with <a href="#">azoospermia</a> as part of an intracytoplasmic sperm injection procedure</li> <li>III. Intracytoplasmic sperm injection for male factor infertility</li> <li>IV. Cryopreservation of embryos, oocytes, ovarian tissue, sperm or testicular tissue (in post-pubertal men) when there is risk of iatrogenic sterilization from chemotherapy or similar medically necessary medical or surgical treatment when <b>all</b> of the following criteria are met:             <ol style="list-style-type: none"> <li>A. No prior elective sterilization</li> <li>B. No known infertility already present</li> <li>C. Post-pubertal and less than 45 years of age (or cryopreservation is no longer desired if younger than age 45)</li> </ol> </li> </ol> <p>The following reproductive techniques are considered <b>investigational</b>:</p> <ol style="list-style-type: none"> <li>I. <a href="#">Co-culture</a> of embryos</li> <li>II. Cryopreservation of testicular tissue in prepubertal boys or ovarian tissue in prepubertal girls</li> <li>III. <a href="#">Intracytoplasmic sperm injection</a> (ICSI) in the absence of male factor infertility</li> </ol>	<p><b>Reproductive Techniques 4.02.04</b></p> <p><b>Policy Statement:</b> The following reproductive techniques may be considered <b>medically necessary</b> for <b>any</b> of the following:</p> <ol style="list-style-type: none"> <li>I. Blastocyst transfer</li> <li>II. Cryopreservation of testicular tissue in adult men with <a href="#">azoospermia</a> as part of an intracytoplasmic sperm injection procedure</li> <li>V. Intracytoplasmic sperm injection for male factor infertility</li> <li>VI. Cryopreservation of embryos, oocytes, ovarian tissue, sperm or testicular tissue (in post-pubertal men) when there is risk of iatrogenic sterilization from chemotherapy or similar medically necessary medical or surgical treatment when <b>all</b> of the following criteria are met:             <ol style="list-style-type: none"> <li>A. No prior elective sterilization</li> <li>B. No known infertility already present</li> <li>C. Post-pubertal and less than 45 years of age (or cryopreservation is no longer desired if younger than age 45)</li> </ol> </li> </ol> <p>The following reproductive techniques are considered <b>investigational</b>:</p> <ol style="list-style-type: none"> <li>I. <a href="#">Co-culture</a> of embryos</li> <li>II. Cryopreservation of testicular tissue in prepubertal boys or ovarian tissue in prepubertal girls</li> <li>III. <a href="#">Intracytoplasmic sperm injection</a> (ICSI) in the absence of male factor infertility</li> </ol>