

Preventive Health Guidelines



The power of prevention

The power of prevention

**Put these preventive health guidelines into practice.
Your family's health could depend on it.**

You've probably heard the old saying, "An ounce of prevention is worth a pound of cure." Your doctors know what preventative and integral services, screenings, and tests are available to you, which often are the best first steps in preventing illness. They also allow for rapid response during the early onset of health problems, when treatments can be most effective.

To take advantage of the power of prevention, you and your family need to stay current with the recommended preventative and integral services, screenings, and tests appropriate for your age, gender, medical history, current health, and family history.

We have compiled the following best-practice guidelines to help you keep track of what's needed and when. Guidelines are recommended for your healthcare and are covered at no cost, unless you are enrolled in a grandfathered plan. Please refer to your Evidence of Coverage (EOC) or Certificate of Insurance (COI) for all covered services. You can call the number on the back of your ID card to verify your coverage as well.

For children ages 0 to 2

Topics you may want to discuss with your doctor

Safety

- Use a checklist to “babyproof” your home.
- Check your home for the presence of lead paint.

Nutrition

- Breastfeeding and iron-enriched formula and food for infants.

Dental health

- Do not put your baby or toddler to bed with a bottle containing juice, milk, or other sugary liquid. Do not prop a bottle in a baby’s or toddler’s mouth. Clean your baby’s gums and teeth daily.
- Use a clean, moist washcloth to wipe gums. Use a soft toothbrush with water only, beginning with eruption of first tooth.
- Age 6 months to preschool: Discuss with your dentist about taking an oral fluoride supplement if water is deficient in fluoride.
- Age 2: Begin brushing child’s teeth with pea-size amount of fluoride toothpaste.

Immunizations

Shot number in a series	1	2	3	4
DTaP (diphtheria, tetanus, acellular pertussis)	2	4	6	15–18 months
Flu, annual	For children 6 months and older ^{2,18}			
Hepatitis A	12–23 months (second dose at least 6 months after first)			
Hepatitis B	0 (birth)	1–2	6–18 months	
Hib (Haemophilus influenzae type b)	2	4	6	12–15 months
IPV (inactivated poliovirus vaccine)	2	4	6–18 months	
Meningococcal Age 2–18 months	For children with risk factors			
MMR (measles, mumps, rubella)	First dose at 12–15 months, second dose at ages 4–6			
Pneumococcal (pneumonia)	2	4	6	12–15 months
Rotarix (rotavirus), or	2	4 months		
RotaTeq (rotavirus)	2	4	6 months	
Varicella (chicken pox)	12–15 months, second dose at ages 4–6			

Screenings/counseling/services

Dental care	Discuss use or prescribe oral supplement for ages 6 months and older. Apply fluoride varnish to primary teeth of infants and children 6 months to 5 years old. ²⁴
Gonococcal ophthalmia	Topical eye medication administered during initial newborn care
Height and weight	Periodically
Newborn Screening Panel	Screening recommended for all disorders listed on the Recommended Uniform Screening Panel (RUSP) ²³
Skin Cancer	Behavioral counseling to minimize exposure to ultraviolet radiation for persons ages 6 months to 24 years old at high risk
Vision	Recommends screening to detect amblyopia or its risk factors ¹

Injury prevention for infants and young children

Decrease risk of SIDS

Sudden infant death syndrome (SIDS) is a leading cause of death for infants. Put infants to sleep on their backs to decrease the risk of SIDS.

Protect your children with car seats

Use the right car seat for your vehicle and for your child’s weight. Read the car seat and vehicle manufacturer’s instructions about installation and use. Use a rear-facing car seat until your child is at least 1 year old and weighs at least 20 pounds.

Babyproof your home

Take these steps to give your child a safe home environment:

- Keep medicines, cleaning solutions, and other dangerous substances in childproof containers, locked up and out of reach of children.
- Use safety gates across stairways (top and bottom) and guards on windows above the first floor.
- Keep hot-water heater temperatures below 120° F.
- Keep unused electrical outlets covered with plastic guards.
- Consider not placing your baby in a baby walker. If you do, provide constant supervision. Block the access to stairways and to objects that can fall (such as lamps) or cause burns (such as stoves or electric heaters).
- Keep objects and foods that can cause choking away from your child. This includes things like coins, balloons, small toy parts, hot dogs (whole or small bites), peanuts, and hard candy.
- Use fences that go all the way around pools and keep gates to pools locked.

For children ages 3 to 10

Topics you may want to discuss with your doctor

Safety

- Use a checklist to “child-proof” your home.
- Check your home for the presence of lead paint.

Exercise

- Participate in physical activity as a family, such as taking walks or playing at the playground.
- Limit screen time (such as mobile devices, computers, and television) to less than two hours a day.

Nutrition

- Eat a healthy diet. Limit fat and calories. Eat fruits, vegetables, beans, and whole grains every day.

Dental health

- Ask your dentist when and how to floss child’s teeth.
- Age 5: Talk to your dentist about dental sealants.

Other topics for discussion

- Well-child visits are a good time to talk to your doctor about any concerns you have with your child’s health, growth, or behavior.

Immunizations

DTaP (diphtheria, tetanus, acellular pertussis)	Fifth dose at ages 4-6 (Tdap for age 7 and above)
Flu, annual	Recommended ^{2,18}
Hepatitis A	For children not previously vaccinated and risk factors are present
Hepatitis B	For children who did not complete the immunization series between 0 and 18 months
Hib (Haemophilus influenzae type b)	For children not previously vaccinated and risk factors are present
IPV (inactivated poliovirus vaccine)	Fourth dose at ages 4–6
Meningococcal	For children with risk factors
MMR (measles, mumps, rubella)	Second dose at ages 4–6
Tdap (tetanus, diphtheria, pertussis)	For children not previously vaccinated with DTaP
Pneumococcal (pneumonia)	For children with risk factors ⁶ or an incomplete schedule (ages 2–5)
Varicella (chicken pox)	Second dose at ages 4–6

Screenings/counseling/services

Height, weight, body mass index (BMI), vision, hearing	At annual exam
Dental care	Discuss use or prescribe oral supplement for ages 6 months and older. Apply fluoride varnish to primary teeth of infants and children 6 months to 5 years old. ²⁴
Obesity	Screening, counseling, and behavioral interventions for children age 6 and older and offer or refer to comprehensive intensive behavioral intervention to promote improvements in weight status
Skin cancer	Behavioral counseling for minimizing exposure to ultraviolet radiation for persons ages 6 months to 24 years old at high risk
Tobacco use and cessation	Discuss education or brief counseling to prevent initiation of tobacco use amongst school-aged children and adolescents 5 years to 18 years old
Vision	Recommends screening to detect amblyopia or its risk factors ¹

Be aware of your child’s recommended weight: Use our online tools to calculate your child’s body mass index (BMI) by logging in to blueshieldca.com and searching for BMI.

Injury prevention:

For older children

- Children should use a booster seat in the car’s back seat until they are at least 8 years old or weigh at least 80 pounds.
- Older children should use car seat belts and sit in the back seat at all times.
- Teach your child traffic safety. Children under 9 years old need supervision when crossing streets.
- Make sure your child wears a helmet while rollerblading or riding a bicycle. Make sure your child uses protective equipment for rollerblading and skateboarding (helmet, wrist, and knee pads).
- Warn your child about the risk of using alcohol and drugs. Many driving and sports-related injuries are caused by the use of alcohol and drugs.

For all ages

- Use smoke and carbon monoxide alarms/detectors in your home. Change the batteries every year, and check once a month to see that they work.
- If you have a gun in your home, make sure that the gun and ammunition are locked up separately and kept out of children’s reach.
- Never drive after drinking alcohol or after marijuana use.
- Use car seat belts at all times.
- Post the number for the Poison Control Center – (800) 222-1222 – near your phone. Also, add the Poison Control Center number to your home “Important Information” list. The number is the same in every U.S. location. Do not try to treat poisoning until you have called the Poison Control Center.

For children ages 11 to 19

Topics you may want to discuss with your doctor

Exercise

- Regular physical activity (at least 30 minutes per day starting at age 11) can reduce the risks of coronary heart disease, osteoporosis, obesity, and diabetes.

Nutrition

- Eat a healthy diet. Limit fat and calories. Eat fruits, vegetables, beans, and whole grains every day.
- Optimal calcium intake for adolescents and young adults is estimated to be 1,200 to 1,500 mg/day.

Sexual health

- Sexually transmitted infection (STI)/ HIV prevention;⁶ practice safer sex (use condoms) or abstinence.
- Avoid unintended pregnancy; use contraception.
- Ongoing follow-up and monitoring must be covered without cost-sharing. Services include periodic HIV testing, serologic testing for hepatitis viruses B and C, periodic serum creatinine testing, periodic pregnancy testing, periodic screening for sexually transmitted bacterial infections, and adherence counseling.

Mental Health and Substance Use disorder

- Use of alcohol, tobacco (cigarettes, vaping, or chewing), inhalants, and other drugs among adolescents is a major concern for parents. Let the doctor know if you have any concerns about your child.
- Mental health and substance use disorders is defined as those conditions listed in the most recent edition of the World Health Organization (WHO) International Classification of Diseases or in the American Psychiatric Association Diagnostic and Statistical Manual of Mental Disorders.

Dental health

- Floss and brush with fluoride toothpaste daily. Seek dental care regularly.

Other topics for discussion

- It is a good idea to let your teenager have private time with the doctor to ask any questions he or she may not feel comfortable asking you.

Immunizations

Flu, annual	Recommended ²
Hepatitis A	Two-dose series for individuals not previously vaccinated and risk factors are present
Hepatitis B	Two-dose series for individuals not previously vaccinated; for individuals with risk factors seeking protection ²⁹
HPV (human papillomavirus)	Two- or three-dose series depending on age at initial vaccination. Recommended for all adolescents 11-12 years and through 18 years of age.
IPV (inactivated poliovirus vaccine)	For individuals not previously vaccinated. Not recommended for individuals 18 years or older
Meningococcal	Routine vaccination two-dose series. First dose at ages 11-12, second dose at age 16.
MMR (measles, mumps, rubella)	Two-dose series for children under 12 previously unvaccinated
Pneumococcal (pneumonia)	For children with risk factors ⁶
Tdap booster (tetanus, diphtheria, pertussis)	For children ages 11-12 who have completed the recommended DTaP immunization series ¹⁷
Varicella (chicken pox)	Two-dose series for children previously unvaccinated

Screenings/counseling/services

Alcohol misuse	Screening for unhealthy alcohol use and behavioral counseling as needed for individuals 12 years and older
Blood pressure, height, weight, BMI, vision, and hearing	At annual exam
Chlamydia and Gonorrhea	Screening for all sexually active women 12 years and older and for women at increased risk for infection ¹¹
Contraception	FDA-approved contraceptive methods for females, education, and counseling ³²
Depression	Screening for all adolescents for major depressive disorder (MDD)
Domestic violence and abuse	Screening for interpersonal and domestic violence for adolescents, women, and women of childbearing age ³⁴
Drug Misuse	Screening for unhealthy drug use for individuals 12 years and older ²⁵
Healthy diet and physical activity	Behavioral counseling ²⁷
Hepatitis C	Screening for HCV infection in persons at high risk of infection ³⁰
HIV	Screening for all adolescents and adults ages 12-65. Recommend preexposure prophylaxis (PrEP) to persons at high risk of HIV acquisition.
Hypertension	Screening for hypertension in adults 18 years and older with office blood pressure measurement (OBPM)
Obesity	Screening, counseling, and behavioral interventions and offer or refer to comprehensive intensive behavioral intervention to promote improvements in weight status
Sexually transmitted infections	Behavioral counseling as needed ²⁶
Skin cancer	Behavioral counseling for minimizing exposure to ultraviolet radiation for adolescents at high risk
Syphilis	Screening for individuals at increased risk for infection ¹²
Tobacco use and cessation	Provide education or brief counseling to prevent initiation of tobacco use among school-aged children and adolescents for individuals 12 years of age and older
Latent tuberculosis infection (LTBI)	Screening for individuals at increased risk of infection ³³

Promoting your preteen's and adolescent's social and emotional development

Parents need to offer open, positive communication while providing clear and fair rules and consistent guidance. Let your child find his or her own path while staying within the boundaries you have set.

- Be a good role model for handling disagreements – for example, talk calmly when disagreeing.
- Praise him or her for successfully avoiding a confrontation – for example, say, "I'm proud of you for staying calm."

- Supervise the websites and computer games that your child uses.
- Set limits on use of computers, telephones, texting, and TV after a set evening hour to help your child get regular sleep.
- Talk to your child about healthy relationships. Dating abuse does occur among preteens and teens.
- Be a role model for healthy eating and regular physical exercise.

For women ages 20 to 49

Topics you may want to discuss with your doctor

Exercise

- Regular physical activity (at least 30 minutes per day) can reduce the risks of coronary heart disease, osteoporosis, obesity, and diabetes.
- Over 40: Consult physician before starting new vigorous physical activity.

Nutrition

- Know your body mass index (BMI), blood pressure, and cholesterol level. Modify your diet accordingly.
- Eat a healthy diet. Limit fat and calories. Eat fruits, vegetables, beans, and whole grains every day.
- Optimal calcium intake for women between ages 25 and 50 is estimated to be 1,000 mg/day.
- Vitamin D is important for bone and muscle development, function, and preservation.

Sexual health

- Sexually transmitted infection (STI)/ HIV prevention;⁶ practice safer sex (use condoms) or abstinence.
- Avoid unintended pregnancy; use contraception.
- Ongoing follow-up and monitoring must be covered without cost-sharing. Services include periodic HIV testing, serologic testing for hepatitis viruses B and C, periodic serum creatinine testing, periodic pregnancy testing, periodic screening for sexually transmitted bacterial infections, and adherence counseling.

Mental Health and Substance Use disorder

- Stop smoking. Limit alcohol consumption. Avoid alcohol or drug use while driving.
- Mental health and substance use disorders is defined as those conditions listed in the most recent edition of the World Health Organization (WHO) International Classification of Diseases or in the American Psychiatric Association Diagnostic and Statistical Manual of Mental Disorders.

Dental health

- Floss and brush with fluoride toothpaste daily. Seek dental care regularly.

If you are pregnant, please refer to the "For pregnant women" page for pregnancy-related recommendations.

Immunizations

Flu, annual	Recommended ²
Hepatitis A	For individuals with risk factors; for individuals seeking protection ³
Hepatitis B	For individuals with risk factors; for individuals seeking protection ^{4,29}
HPV (human papillomavirus)	Recommended for all adults through 26 years of age. 2 or 3-dose series depending on age at initial vaccination
Meningococcal	For individuals with risk factors present ⁸
MMR (measles, mumps, rubella)	Once, without proof of immunity or if no previous second dose ⁵
Pneumococcal (pneumonia)	For individuals with risk factors ⁷
Td booster (tetanus, diphtheria)	Recommended once every 10 years ¹⁵
Varicella (chicken pox)	Recommended for adults without evidence of immunity; 2-dose series 4-8 weeks apart ¹⁰

Screenings/counseling/services

Alcohol misuse	Screening for unhealthy alcohol use and behavioral counseling as needed for individuals 12 years and older
Blood pressure, height, weight, BMI, vision, and hearing	At well visit, annually
BRCA risk assessment and genetic counseling/testing	Women with a positive result on the risk assessment tool or have a family history of breast, ovarian, tubal, prostate, pancreatic, or peritoneal cancer are recommended to receive genetic counseling and/or genetic testing. ²⁰
Breast cancer	Recommend mammogram every 1–2 years beginning at age 40 through 49
Breast cancer medication use	Recommends prescription or risk-reducing medications to women who are at increased risk for breast cancer and at low risk for adverse medication effects
Cardiovascular disease	Statin use for primary prevention in adults ³⁵
Cervical cancer	Recommended screening every 3 years with cervical cytology by Pap tests for individuals 21 to 65 years of age
Chlamydia and Gonorrhea	Screening for all sexually active women 12 years and older and for women at increased risk for infection ¹¹
Colorectal cancer	Screening for adults ages 45–75 ⁹
Contraception	FDA-approved female contraceptive methods, education, and counseling ³²
Depression	Screening for all adults
Diabetes	Screening for overweight or obese adults ages 40–70 ²²
Domestic violence and abuse	Screening for interpersonal and domestic violence for adolescents, women, and women of childbearing age ³⁴
Drug Misuse	Screening for unhealthy drug use for individuals 12 years and older ²⁵
Healthy diet and physical activity	Behavioral counseling ²⁷
Hepatitis C	Screening for HCV infection in persons at high risk of infection ages 18 to 79 ³⁰
HIV	Screening for all adolescents and adults ages 12–65. Recommend preexposure prophylaxis (PrEP) to persons at high risk of HIV acquisition.
Hypertension	Screening for hypertension in adults 18 years and older with office blood pressure measurement (OBPM)
Latent tuberculosis infection (LTBI)	Screening for individuals at increased risk for infection ³³
Obesity	Screening, counseling, and behavioral interventions and offer or refer to comprehensive intensive behavioral intervention to promote improvements in weight status
Osteoporosis	Screening for women at increased risk ¹³
Sexually transmitted infections	Behavioral counseling as needed ²⁶
Skin cancer	Behavioral counseling for minimizing exposure to ultraviolet radiation for young adults to age 24 at high risk
Syphilis	Screening for pregnant women and individuals at increased risk for infection ¹²
Tobacco use and cessation	Screening for tobacco use and cessation intervention for individuals 12 years and older

For men ages 20 to 49

Topics you may want to discuss with your doctor

Exercise

- Regular physical activity (at least 30 minutes per day) can reduce the risks of coronary heart disease, osteoporosis, obesity, and diabetes.
- Men over 40: Consult physician before starting new vigorous physical activity.

Nutrition

- Know your body mass index (BMI), blood pressure, and cholesterol level. Modify your diet accordingly.
- Vitamin D is important for bone and muscle development, function, and preservation.

Sexual health

- Sexually transmitted infection (STI)/HIV prevention;^{1,6} practice safer sex (use condoms) or abstinence.
- Ongoing follow-up and monitoring must be covered without cost-sharing. Services include periodic HIV testing, serologic testing for hepatitis viruses B and C, periodic serum creatinine testing, periodic screening for sexually transmitted bacterial infections, and adherence counseling.

Mental Health and Substance Use disorder

- Stop smoking. Limit alcohol consumption. Avoid alcohol or drug use while driving.
- Mental health and substance use disorders is defined as those conditions listed in the most recent edition of the World Health Organization (WHO) International Classification of Diseases or in the American Psychiatric Association Diagnostic and Statistical Manual of Mental Disorders.

Dental health

- Floss and brush with fluoride toothpaste daily. Seek dental care regularly.

“Know your numbers.” We encourage you to learn your “numbers” at your doctor visit and work toward the optimal goals through exercise and a healthy diet.

Immunizations

Flu, annual	Recommended ²
Hepatitis A	For individuals with risk factors; for individuals seeking protection ³
Hepatitis B	For individuals with risk factors; for individuals seeking protection ^{4,29}
HPV (human papillomavirus)	Recommended for all adults through 26 years of age. 2 or 3-dose series depending on age of initial vaccination
Meningococcal	For individuals with risk factors present ⁸
MMR (measles, mumps, rubella)	Once, without proof of immunity or if no previous second dose ⁵
Pneumococcal (pneumonia)	For individuals with risk factors ⁷
Td booster (tetanus, diphtheria)	Recommended once every 10 years ¹⁵
Varicella (chicken pox)	Recommended for adults without evidence of immunity; 2-dose series 4-8 weeks apart ¹⁰

Screenings/counseling/services

Alcohol misuse	Screening for unhealthy alcohol use and behavioral counseling as needed for individuals 12 years and older
Blood pressure, height, weight, BMI, vision, and hearing	At annual exam
Cardiovascular disease	Statin use for primary prevention in adults ³⁵
Colorectal cancer	Screening for adults ages 45-75 ⁹
Depression	Screening for all adults
Diabetes	Screening for overweight or obese adults ages 40-70 ²²
Drug Misuse	Screening for unhealthy drug use for individuals 12 years and older ²⁵
Healthy diet and physical activity	Behavioral counseling for 18 years to 79 years of age ²⁷
Hepatitis C	Screening for HCV infection in persons at high risk of infection ³⁰
HIV	Screening for all adolescents and adults ages 12-65. Recommend preexposure prophylaxis (PrEP) to persons at high risk of HIV acquisition.
Hypertension	Screening for hypertension in adults 18 years and older with office blood pressure measurement (OBPM)
Latent tuberculosis infection (LTBI)	Screening for individuals at increased risk for infection ³³
Obesity	Screening, counseling, and behavioral interventions and offer or refer to comprehensive intensive behavioral intervention to promote improvements in weight status
Sexually transmitted infections	Behavioral counseling as needed ²⁶
Skin cancer	Behavioral counseling for minimizing exposure to ultraviolet radiation for young adults to age 24 at high risk
Syphilis	Routine screening for individuals at increased risk for infection ¹²
Tobacco use and cessation	Screening for tobacco use and cessation intervention for individuals 12 years and older

Heart health factors

Optimal goals

Total cholesterol	Less than 200 mg/dL
LDL “bad” cholesterol	Less than 100 mg/dL
HDL “good” cholesterol	60 mg/dL or higher
Triglycerides	Less than 150 mg/dL
Blood pressure	Less than 120/80 mmHg
Fasting glucose	Less than 100 mg/dL
Body mass index (BMI)	Less than 25 kg/m ²
Exercise	Minimum of 30 minutes most days of the week

For men and women age 50 and older

Topics you may want to discuss with your doctor

Nutrition

- Eat a healthy diet. Limit fat and calories. Eat fruits, vegetables, beans, and whole grains every day.
- Optimal calcium intake is estimated to be 1,500 mg/day for postmenopausal women not on estrogen therapy.
- Vitamin D is important for bone and muscle development, function, and preservation.

Sexual health

- Sexually transmitted infection (STI)/HIV prevention;⁶ practice safer sex (use condoms) or abstinence.
- Ongoing follow-up and monitoring must be covered without cost-sharing. Services include periodic HIV testing, serologic testing for hepatitis viruses B and C, periodic serum creatinine testing, periodic screening for sexually transmitted bacterial infections, and adherence counseling.

Mental Health and Substance Use disorder

- Stop smoking. Limit alcohol consumption. Avoid alcohol or drug use while driving.
- Mental health and substance use disorders is defined as those conditions listed in the most recent edition of the World Health Organization (WHO) International Classification of Diseases or in the American Psychiatric Association Diagnostic and Statistical Manual of Mental

Dental health

- Floss and brush with fluoride toothpaste daily. Seek dental care regularly.

Other topics for discussion

- Fall prevention.
- Possible risks and benefits of hormone replacement therapy (HRT) for post-menopausal women.
- The dangers of drug interactions.
- Physical activity.
- Glaucoma eye exam by an eye care professional (i.e., an ophthalmologist, optometrist) for those age 65 and older.

For heart health, adults should exercise regularly (at least 30 minutes a day on most days), which can help reduce the risks of coronary heart disease, osteoporosis, obesity, and diabetes. Consult your physician before starting a new vigorous physical activity.

Immunizations

Flu, annual	Recommended ²
Hepatitis A	For individuals with risk factors; for individuals seeking protection ³
Hepatitis B	For individuals with risk factors; for individuals seeking protection ⁴
Meningococcal	For individuals with risk factors present ⁵
MMR (measles, mumps, rubella)	Once, without proof of immunity or if no previous second dose ⁵
Pneumococcal (pneumonia)	Recommended for individuals age 65 and older; and individuals under age 65 with risk factors ⁷
Td booster (tetanus, diphtheria)	Recommended once every 10 years ¹⁵
Varicella (chickenpox)	Recommended for adults without evidence of immunity; 2-dose series 4–8 weeks apart ¹⁰
Zoster (shingles)	Two-dose series of RZV 2-6 months apart

Screenings/counseling/services

AAA (abdominal aortic aneurysm)	For ages 65–75 who have ever smoked, one-time screening for AAA by ultrasonography
Alcohol misuse	Screening for unhealthy alcohol use and behavioral counseling as needed for individuals 12 years and older
Aspirin	Use for primary prevention of cardiovascular disease and colorectal cancer for adults who are at increased risk ¹⁹
Blood pressure, height, weight, BMI, vision, and hearing	At annual exam
BRCA risk assessment and genetic counseling/testing	Women with a positive result on the risk assessment tool or have a family history of breast, ovarian, tubal, prostate, pancreatic, or peritoneal cancer are recommended to receive genetic counseling and/or genetic testing. ²⁰
Breast cancer	Recommended biennial screening mammography for women aged 40 years and older
Breast cancer medication use	Recommended prescription or risk-reducing medications to women who are at increased risk for breast cancer and at low risk for adverse medication effects
Cardiovascular disease	Statin use for primary prevention in adults ³⁵
Cervical cancer	Recommended screening every 3 years with cervical cytology and every 5 years with high risk HPV testing
Chlamydia and Gonorrhea	Screening for individuals who are at increased risk for infection ¹¹
Colorectal cancer	Screening for adults ages 45-75 ^{9,21}
Depression	Screening for all adults
Diabetes	Screening for overweight or obese adults ages 40–70 ²²
Domestic violence and abuse	Screening for interpersonal and domestic violence for adolescents, women, and women of childbearing age ³⁴
Drug Misuse	Screening for unhealthy drug use for individuals 12 years and older ²⁵
Fall prevention	Recommended exercise interventions for adults ages 65 or older at increased risk ²⁸
Healthy diet and physical activity	Behavioral counseling ²⁷
Hepatitis C	Screening for HCV infection in persons at high risk of infection ³⁰
HIV	Screening for all adolescents and adults ages 12–65. Recommend preexposure prophylaxis (PrEP) to persons at high risk of HIV acquisition.
Hypertension	Screening for hypertension in adults 18 years and older with office blood pressure measurement (OBPM)
Latent tuberculosis infection (LTBI)	Screening for individuals at increased risk for infection ³³
Lung cancer	Screening for lung cancer annually in persons with smoking history using low-dose computed tomography ³¹
Obesity	Screening, counseling, and behavioral interventions and offer or refer to comprehensive intensive behavioral intervention to promote improvements in weight status
Osteoporosis	Recommended routine screening for osteoporosis with bone measurement testing for women 65 years and older ¹³
Sexually transmitted infections	Behavioral counseling as needed ²⁶
Syphilis	Screening for individuals at increased risk for infection ¹²
Tobacco use and cessation	Screening for tobacco use and cessation intervention

For pregnant women

Screenings/counseling/services

Alcohol misuse	Screening for unhealthy alcohol use and behavioral counseling as needed
Aspirin	Low-dose aspirin use for the prevention of morbidity and mortality from preeclampsia ¹⁴
Asymptomatic bacteriuria	Recommended screening using urine culture in pregnant women
Breast-feeding primary care interventions	Provide interventions during pregnancy and after birth to support breast-feeding
Chlamydia and Gonorrhea	Screening for all sexually active women under age 24 and for women at increased risk for infection ¹¹
Depression	Refer pregnant and postpartum persons who are at increased risk of depression to counseling interventions
Drug Misuse	Screening for unhealthy drug use for individuals 12 years and older ²⁵
Folic acid	Discuss use of 0.4 to 0.8 mg daily
Gestational diabetes	Recommended screening for gestational diabetes mellitus (GDM) in asymptomatic pregnant women after 24 weeks of gestation
Healthy weight gain	Recommended behavioral counseling and interventions to promote healthy weight gain and prevent excess gestational weight gain
Hepatitis B	First prenatal visit ²⁹
HIV	Screening for HIV infection in all pregnant persons, including those who are present in labor or at delivery whose HIV status is unknown
Maternal Mental Health	Covered services for individuals who present written documentation of being diagnosed with a maternal mental health condition. Completion of covered services may extend up to 12 months from date of diagnosis or from the end of the pregnancy, whichever occurs later.
Preeclampsia	Screening with blood pressure measurements
Rh (D) incompatibility	Recommended repeated Rh(D) antibody testing for all unsensitized Rh(D) – negative women at 24 to 28 weeks' gestation, unless the biological father is known to be Rh(D) – negative
Syphilis	Recommend early screening for infection in all pregnant women ¹²
Tdap	One dose of Tdap is recommended during each pregnancy, preferably in the early part of the gestational weeks 27–36
Tobacco use and cessation	Screening for tobacco use and tobacco-cessation intervention

Having a baby? Be aware that while almost all women get the “baby blues” after childbirth, as many as 10% will get postpartum depression. For more information, visit our website, [blueshieldca.com](https://www.blueshieldca.com), or see your healthcare provider.

Recommendations for a healthy pregnancy

Prenatal care

Begin within 14 days of confirming pregnancy.

Dietary supplements

Women of childbearing age should take 0.4 to 0.8 mg of folic acid daily to decrease the risk of fetal brain and spinal cord birth defects. The recommended calcium intake for pregnant or nursing women is 1,000 milligrams daily.

Screenings and diagnostics

Blood pressure and weight check at all visits; urine test; obstetrical history and physical; screenings for asymptomatic bacteriuria; chlamydia; gestational diabetes; Group B streptococcal bacteria; hepatitis B; syphilis; gonorrhea; hematocrit; rubella; varicella; Rh (D) incompatibility; HIV counseling and screening; ultrasonography;

screening for alpha fetoprotein; chorionic villus screening (CVS) or amniocentesis (for women age 35 and older); blood tests for certain birth defects; fundal height; fetal heart tones.

Discussion topics at prenatal care visits

Prior vaccinations (including flu shots), history of genital herpes, nutrition, smoking cessation, preterm labor risk, domestic abuse, mental health as an initial intervention service after screening for interpersonal and domestic violence, and other medication and drug use.

Postpartum care

To be performed within three to seven weeks following delivery. Postpartum exam to include weight, blood pressure, breast and abdomen exam, or pelvic exam.

Endnotes

1. All children 3 to 5 years are at risk of vision abnormalities and should be screened; specific risk factors include strabismus, refractive errors, and media opacity.
2. Annual vaccination against influenza is recommended for all persons age 6 months and older, including all adults. Healthy, nonpregnant adults under age 50 without high-risk medical conditions can receive either intranasally administered live, attenuated influenza vaccine, or inactivated vaccine. Other persons should receive the inactivated vaccine. Adults age 65 and older can receive the standard influenza vaccine or the high-dose influenza vaccine.
3. Risk factors for hepatitis A should be discussed with your provider.
4. Risk factors for hepatitis B should be discussed with your provider.
5. Measles component: Adults born before 1957 can be considered immune to measles. Adults born on or after 1957 should receive one or more doses of MMR, depending on their immune status. Also, a second dose of MMR may be necessary if exposed, traveling internationally, and other factors. Rubella component: Women with unreliable vaccination history should check with their provider. Check with your doctor for details regarding pregnancy.
6. Administer pneumococcal vaccine to children with certain underlying medical conditions, including a cochlear implant. A single revaccination should be administered after five years to children with functional or anatomic asplenia or an immunocompromising condition.
7. One dose for adults at risk, including those with chronic lung diseases (including asthma and COPD); cardiovascular diseases, diabetes mellitus, chronic liver disease, chronic renal failure, sickle cell disease, and immunocompromising conditions. Vaccination is also recommended in adults who smoke cigarettes and residents of nursing homes and long-term care facilities. Vaccination is not recommended in Alaska Native or American Indian persons unless they have another risk factor present. A second pneumococcal dose may be necessary for people age 65 and older who received the vaccine more than five years previously and were younger than 65 at the time of the primary vaccination. A one-time revaccination is recommended after five years for people with certain medical conditions, including immunosuppressive conditions and people who have undergone chemotherapy.
8. Individuals at risk for meningococcal disease include international travelers, college-bound students, or anyone with a damaged or removed spleen or with terminal complement component deficiency. These individuals should discuss the risks and benefits of vaccination with their doctor.
9. Colorectal cancer is the third leading cause of cancer death for both men and women. It is the most frequently diagnosed among persons aged 65 to 74 years. It is estimated that 10.5% of new colorectal cancer cases occur in persons younger than 50 years.
10. Individuals at risk for varicella infection include those who have close contact with persons at high risk for severe disease (healthcare workers and family contacts of immunocompromised persons) or are at high risk for exposure or transmission (e.g., teachers of young children; childcare employees; residents and staff members of institutional settings, including correctional institutions; college students; military personnel; adolescents and adults living in households with children; nonpregnant women of childbearing age; and international travelers).
11. Risk factors for chlamydia and gonorrhea infection include history of chlamydial or other sexually transmitted infections, new or multiple sexual partners, inconsistent condom use, commercial sex work, and drug use.
12. Risk factors for syphilis infection include all adolescents and adults who receive health care in a high-prevalence or high-risk clinical setting, men who have had sex with men, commercial sex workers, and those in adult correctional facilities. Individuals being treated for sexually transmitted diseases may be more likely than others to engage in high-risk behavior.
13. Osteoporotic fractures, particularly hip fractures, are associated with limitation of ambulation, chronic pain and disability, loss of independence and quality of life. Women have higher rates of osteoporosis than men at any given age.
14. Pregnant women who are at high risk for preeclampsia should use low-dose aspirin (81 mg/d) as preventive medication after 12 weeks of gestation.
15. People in contact with infants under 12 months of age and healthcare personnel can be given the Td vaccine as soon as feasible. It is recommended that Tdap should replace a single dose of Td for adults under age 65 if they have not previously received a dose of Tdap.
16. Sexually transmitted infections, also known as sexually transmitted diseases, include chlamydia, gonorrhea, herpes, HIV, HPV, syphilis, and others. See infection-specific notes for information on risk factors for sexually transmitted infections.
17. The Tdap (tetanus, diphtheria, acellular pertussis) booster is recommended in children ages 11 to 12 who have completed the childhood DTaP immunization series and have not yet received a tetanus and diphtheria (Td) booster dose.
18. Children through age 9 getting flu vaccine for the first time – or who received flu vaccine – should get two doses, at least four weeks apart.
19. Low-dose aspirin use for the primary prevention of cardiovascular disease (CVD) and colorectal cancer (CRC) in adults ages 50 to 59 years who have a 10% or greater 10-year CVD risk, are not at increased risk for bleeding, have a life expectancy of at least 10 years, and are willing to take low-dose aspirin daily for at least 10 years.
20. For breast cancer screening, BRCA mutation referral for genetic risk assessment and evaluation for breast and ovarian susceptibility is recommended for women with family history associated with an increased risk for deleterious mutations in BRCA1 or BRCA2 genes. Please see Blue Shield of California medical policy on Genetic Testing for Hereditary Breast and/or Ovarian Cancer.
21. Colorectal cancer screenings include fecal occult blood annually, sigmoidoscopy every five years, and colonoscopy every 10 years. Beginning age and screening interval can be reduced for patients at increased risk. Multi-targeted stool DNA testing every three years.
22. Diabetes screening should be performed for adults ages 35 to 70 who are overweight and obese. Intensive behavioral counseling interventions to promote a healthful diet and physical activity for patients with abnormal blood glucose.
23. Selection of conditions based on "Newborn Screening: Toward a Uniform Screening Panel and System" as authored by the American College of Medical Genetics (ACMG) and commissioned by the Health Resources and Service Administration (HRSA).
24. Fluoride oral supplement should be discussed at preventive care visit if primary water source is deficient in fluoride.
25. Unhealthy drug use screening should be implemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred.
26. Behavioral counseling to prevent sexually transmitted infections is for sexually active adolescents and adults who meet the following criteria: current sexually transmitted infections, sexually transmitted infections within the past year, multiple current sexual partners, and in non-monogamous relationships if they reside in a community with a high rate of sexually transmitted infections.
27. Intensive behavioral counseling to promote healthy diet and physical activity is recommended for all adults who have hyperlipidemia or have any known risk factors for cardiovascular and diet-related chronic disease. Diabetes Prevention Program – part of Wellvolution® – is Blue Shield's lifestyle medicine program focused on diabetes prevention.
28. Falls prevention counseling for older adults to exercise or physical therapy to prevent falls in community-dwelling adults 65 and older who are at increased risk for falls.
29. Hepatitis B screening for non-pregnant adolescents and adults for hepatitis B virus infection at high risk for infection; pregnant women at their first prenatal visit.
30. Hepatitis C screening for adults for hepatitis C virus infection at high risk for infection.
31. Lung cancer screening for adults ages 55 to 80 who have a smoking history.
32. For self-administered hormonal contraceptives, you may receive up to a 12-month supply.
33. Tuberculosis and latent tuberculosis infection (LTBI) for asymptomatic adults at increased risk for infection.
34. Screening and counseling for interpersonal and domestic violence is a covered service for adolescents, women, and women of childbearing age at least annually, and, when needed, those who screen positive are provided or referred to initial intervention services. Mental health is an initial intervention service after screening for interpersonal and domestic violence.
35. Statin use for the primary prevention of cardiovascular disease in adults – The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (i.e., symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the prevention of CVD events and mortality when all of the following criteria are met: 1) they are ages 40 to 75; 2) they have one or more CVD risk factors (i.e., dyslipidemia, diabetes, hypertension, or smoking); and 3) they have a calculated 10-year risk of a cardiovascular event of 10% or greater. Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipids screening in adults ages 40 to 75. Statin medications are a pharmacy benefit.

Wellvolution is a registered trademark of Blue Shield of California. Wellvolution and all associated digital and in-person health programs, services, and offerings are managed by Solera, Inc., a health company committed to changing lives by guiding people to better health in their communities.

These are Blue Shield of California's Preventive Health Guidelines, which are based on nationally recognized guidelines. Members must refer to their *Evidence of Coverage* or *Certificate of Insurance* or *Policy* for plan/policy coverage of preventive health benefits.

Blue Shield of California complies with applicable state laws and federal civil rights laws, and does not discriminate on the basis of race, color, national origin, ancestry, religion, sex, marital status, gender, gender identity, sexual orientation, age, or disability.

Blue Shield of California cumple con las leyes estatales y las leyes federales de derechos civiles vigentes, y no discrimina por motivos de raza, color, país de origen, ascendencia, religión, sexo, estado civil, género, identidad de género, orientación sexual, edad ni discapacidad.

Blue Shield of California 遵循適用的州法律和聯邦公民權利法律，並且不以種族、膚色、原國籍、血統、宗教、性別、婚姻狀況、性別認同、性取向、年齡或殘障為由而進行歧視。