

## HEDIS Provider Guide: Chlamydia Screening in Women (CHL)

## **Measure Description**

Women 16 to 24 years of age who were identified as sexually active and who had at least one chlamydia test during the measurement year.

Members identified as sexually active:

- Pregnancy
- Sexually Transmitted Infection (STI) screening
- Contraceptive prescribed
- Pregnancy test completed

## **Using Correct Billing Codes**

Description	CPT Codes
Chlamydia Screening	87110, 87270, 87320, 87490-87492, 87810

Contraceptives		
Desogestrel-ethinyl estradiol	Ethinyl estradiol-norethindrone	
Dienogest-estradiol (multiphasic)	Ethinyl estradiol-norgestimate	
Drospirenone-ethinyl estradinol	Ethinyl estradiol-norgestrel	
Drospirenone-ethinyl estradiol-levomefolate	Etonogestrel	
(biphasic)	Levonorgestrel	
Ethinyl estradiol-ethynodiol	Medroxyprogesterone	
Ethinyl estradiol-etonogestrel	Mestranol-norethindrone	
Ethinyl estradiol-levonorgestrel	Norethindrone	
Ethinyl estradiol-norelgestromin		

Diaphragm and Spermicide	
Diaphragm	Nonoxynol

## How to Improve HEDIS® Scores

- Perform chlamydia screening every year on every 16 to 24-year-old female identified as sexually active (use any visit opportunity).
- Add chlamydia screening as a standard lab for women 16 to 24 years old. Use adolescent well-care
  exams and well woman exams for this purpose.
- Remember that chlamydia screening can be performed through a urine test. Offer this option for your patients.
- Request a focused list of members who need chlamydia screening from your Blue Shield Promise Quality Program Manager.