

Promise Health Plan

letermovir injection (Prevymis)

Medical Benefit Drug Policy

Place of Service

Office Administration

Infusion Center Administration

Home Infusion Administration

Hospital Administration

Outpatient Facility Administration

Drug Details

USP Category: ANTIVIRALS

Mechanism of Action: CMV DNA terminase complex inhibitor

HCPCS:

C9399:Unclassified drugs or biologicals

J3490:Unclassified drugs

How Supplied:

- 240 mg/12 mL (20 mg/mL) single-dose vial (NDC 0006-5003-01)
- 480 mg/24 mL (20 mg/mL) single-dose vial (NDC 0006-5004-01)

Condition(s) listed in policy (see coverage criteria for details)

• Cytomegalovirus (CMV) Prophylaxis

Any condition not listed in this policy requires a review to confirm it is medically necessary. For conditions that have not been approved for intended use by the Food and Drug Administration (i.e., off-label use), the criteria outlined in the California Code of Regulations (CCR), Title 22, Section 51303 and 51313 must be met.

Special Instructions and Pertinent Information

Provider must submit documentation (such as office chart notes, lab results or other clinical information) to ensure the member has met all medical necessity requirements.

The member's specific benefit may impact drug coverage. Other utilization management processes, and/or legal restrictions may take precedence over the application of this clinical criteria.

Coverage Criteria

Effective: 05/01/2025

The following condition(s) require Prior Authorization/Preservice.

Cytomegalovirus (CMV) Prophylaxis

Meets medical necessity if all the following are met:

- 1. Being used as prophylaxis of CMV infection and disease
- 2. Either of the following:
 - a. Recipient of an allogeneic HSCT and all the following:
 - i. Diagnosis of CMV-seropositive

letermovir injection (Prevymis)

Page 1 of 3

- ii. Either of the following:
 - 1. Being used within 100 days post-transplant
 - 2. Being used through day 200 post-HSCT in patients at risk for late CMV infection and disease
- b. Recipient of a kidney transplant and all of the following:
 - i. High risk (Donor CMV seropositive/Recipient CMV seronegative [D+/R-])
 - ii. Being used within 200 days post-transplant
- 3. Medical reason why the oral formulation of Prevymis cannot be used

Covered Doses:

Up to 480 mg given as an IV infusion once daily

Coverage Period:

HSCT: 100 days post-transplant

HSCT and at risk for late CMV infection: 200 days post-transplant

Kidney transplant: 200 days post-transplant

ICD-10:

Z94.81

Additional Information

Pediatric dosing for HSCT recipients

Recommended daily IV dosage of Prevymis in pediatric HSCT recipients 6 months to less than 12 years of age or 12 yeasts of age and older and weighing less than 30 kg (from Prescribing Information)

Body weight	Daily IV dose
30 kg and above	480 mg
15 kg to less than 30 kg	120 mg
7.5 kg to less than 15 kg	60mg
6 kg to less than 7.5 kg	40 mg

Recommended daily oral dosage of Prevymis in pediatric HSCT recipients 6 months to less than 12 years of age or 12 years of age and older and weighing less than 30 kg (from Prescribing Information)

Body weight	Daily oral	Tablets	Oral Pellets <i>(product</i>
	dose		availability pending)
30 kg and above	480 mg	One 480 mg tablet or two 240 mg tablets	Four 120 mg packets of oral pellets

letermovir injection (Prevymis) Page 2 of 3

15 kg to less than 30 kg	120 mg	One 240 mg tablet	Two 120 mg packets of oral pellets
7.5 kg to less than 15 kg	60mg	Not recommended	One 120 mg packet of oral pellets
6 kg to less than 7.5 kg	40 mg	Not recommended	Four 20 mg packets of oral pellets

References

- 1. AHFS. Available by subscription at http://www.lexi.com
- 2. DrugDex. Available by subscription at http://www.micromedexsolutions.com/home/dispatch
- 3. National Comprehensive Cancer Network. Prevention and Treatment of Cancer-Related Infections (Version 3.2024). Available at http://www.nccn.org.
- 4. Prevymis (letermovir) Prescribing Information. Merck & Co., Rahway, NJ: 2024.

Review History

Effective: 05/01/2025

Date of Last Annual Review: 1Q2025 Changes from previous policy version:

• No clinical change following annual review.

Blue Shield of California Medication Policy to Determine Medical Necessity Reviewed by P&T Committee

letermovir injection (Prevymis)
Page 3 of 3