

1.01.17 Pelvic Floor Stimulation as a Treatment of Urinary and Fecal Incontinence

Original Policy Date:	February 27, 2015	Effective Date:	November 1, 2024
Section:	1.0 Durable Medical Equipment	Page:	Page 1 of 18

Policy Statement

- I. Electrical or magnetic stimulation of the pelvic floor muscles (pelvic floor stimulation) is considered **investigational** as a treatment for **either** of the following:
 - A. Urinary incontinence
 - B. Fecal incontinence

NOTE: Refer to [Appendix A](#) to see the policy statement changes (if any) from the previous version.

Policy Guidelines

Coding

See the [Codes table](#) for details.

Description

Pelvic floor stimulation is proposed as a nonsurgical treatment option for women and men with urinary or fecal incontinence. This approach involves either electrical stimulation of pelvic floor musculature or extracorporeal pulsed magnetic stimulation.

Related Policies

- Percutaneous and Subcutaneous Tibial Nerve Stimulation
- Sacral Nerve Neuromodulation/Stimulation

Benefit Application

Benefit determinations should be based in all cases on the applicable contract language. To the extent there are any conflicts between these guidelines and the contract language, the contract language will control. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

Some state or federal mandates (e.g., Federal Employee Program [FEP]) prohibits plans from denying Food and Drug Administration (FDA)-approved technologies as investigational. In these instances, plans may have to consider the coverage eligibility of FDA-approved technologies on the basis of medical necessity alone.

Regulatory Status

Several electrical stimulators have been cleared by the U.S. Food and Drug Administration (FDA). In 2006, the MyoTrac Infinity™ (Thought Technology) and in 2015, the Apex®M (now Apex®; InControl Medical) nonimplanted electrical stimulators for treating urinary incontinence were cleared for marketing by the FDA through the 510(k) process. Predicate devices are also used to treat urinary incontinence, including the Pathway™ CTS 2000 (Prometheus Group) and the InCare® PRS (Hollister). In 2011, the itouch Sure Pelvic Floor Exerciser (TensCare) was cleared for marketing.

In 2000, the NeoControl[®] Pelvic Floor Therapy System (Neotonus) was cleared through the FDA 510(k) process for treating urinary incontinence in women. This device, formerly known as the Neotonus Model 1000 Magnetic Stimulator, provides noninvasive electromagnetic stimulation of pelvic floor musculature. The magnetic system is embedded in a chair seat; patients sit on the chair fully clothed and receive the treatment. The magnetic fields are controlled by a separate power unit.

In 2014, the InTone[®] MV (InControl Medical), a nonimplantable device that provides electrical stimulation and/or biofeedback via manometry, was cleared by the FDA. The device is intended to treat male and female urinary and fecal incontinence.

FDA product code: KPI.

Rationale

Background

Pelvic Floor Stimulation

Pelvic floor stimulation (PFS) involves electrical stimulation of pelvic floor muscles using either a probe wired to a device for controlling the electrical stimulation or, more recently, extracorporeal electromagnetic (also called magnetic) pulses. Stimulation of the pudendal nerve to activate the pelvic floor musculature may improve urethral closure. In addition, PFS is thought to improve partially denervated urethral and pelvic floor musculature by enhancing the process of reinnervation. Methods of electrical PFS have varied in location (e.g., vaginal, rectal), stimulus frequency, stimulus intensity or amplitude, pulse duration, pulse to rest ratio, treatments per day, number of treatment days per week, length of time for each treatment session, and overall time period for device use between clinical and home settings. Variations in the amplitude and frequency of the electrical pulse are used to mimic and stimulate the different physiologic mechanisms of the voiding response, depending on the etiology of the incontinence (i.e., either detrusor instability, stress incontinence, or a mixed pattern). Magnetic PFS does not require an internal electrode; instead, patients sit fully clothed on a specialized chair with an embedded magnet.

Patients receiving electrical PFS may undergo treatment in a physician's office or physical therapy facility, or patients may undergo initial training in a physician's office followed by home treatment with a rented or purchased pelvic floor stimulator. Magnetic PFS may be administered in the physician's office.

Literature Review

Evidence reviews assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function—including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be

used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Promotion of greater diversity and inclusion in clinical research of historically marginalized groups (e.g., People of Color [African-American, Asian, Black, Latino and Native American]; LGBTQIA (Lesbian, Gay, Bisexual, Transgender, Queer, Intersex, Asexual); Women; and People with Disabilities [Physical and Invisible]) allows policy populations to be more reflective of and findings more applicable to our diverse members. While we also strive to use inclusive language related to these groups in our policies, use of gender-specific nouns (e.g., women, men, sisters, etc.) will continue when reflective of language used in publications describing study populations.

Electrical Pelvic Floor Stimulation for Urinary Incontinence Clinical Context and Therapy Purpose

The purpose of pelvic floor stimulation (PFS) in individuals who have urinary incontinence is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with urinary incontinence. Types of urinary incontinence include stress incontinence, urgency incontinence, and mixed (both stress and urgency).

Urinary incontinence is a common condition and can have a substantial impact on quality of life. Estimates from the National Center for Health Statistics have suggested that, among noninstitutionalized persons 65 years of age and older, 44% have reported issues with urinary incontinence.¹ Urinary incontinence in women is common, with some estimates citing a 50% incidence. Factors that increase a woman's risk include older age, obesity, parity, vaginal delivery, and family history.

Urinary incontinence is less common in men, with estimates ranging from 11% to 16%² in men greater than 65 years. Factors that increase a man's risk include older age, prostate disease, urinary tract infection history, impaired activities of daily living, neurologic disease, constipation, diabetes, and sleep apnea.

Interventions

The therapy being considered is electrical PFS for urinary incontinence. In an electrical PFS procedure, a probe delivers electrical pulses to stimulate the pudendal nerve, which activates the pelvic floor musculature. Activation of this musculature is believed to improve urethral closure.

Comparators

The following therapies are currently being used to make decisions about urinary incontinence: magnetic PFS or neuromodulation, behavioral therapies (e.g., monitoring fluid intake, bladder, and pelvic floor muscle training, diet), and medications.

Outcomes

The general outcomes of interest include a reduction in symptoms (e.g., number of incontinence episodes) and improvements in quality of life and cure rates. Short-term results can be measured at 6 months.³ Longer-term follow-up may be necessary to determine if treatment has durable effects.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- a. To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;

- b. In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- c. To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- d. Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Urinary Incontinence in Women

Systematic Reviews

A TEC Assessment (2000) concluded there was insufficient evidence that electrical PFS improved health outcomes compared with placebo or other behavioral therapies in women with stress, urge, or mixed incontinence.⁴

Subsequently, several systematic reviews of the literature with pooled study findings have been published.

Leonardo et al (2022) published a systematic review and meta-analysis of 8 RCTs (N=562) which evaluated the comparative effectiveness of biofeedback-assisted pelvic floor muscle training (PFMT) versus PFS versus a control group (PFMT alone, bladder training, or lifestyle recommendations only) in women with overactive bladder.⁵ Outcomes assessed included quality of life, number of episodes of incontinence, and number of patients who improved or were cured. The PFS group exhibited significant differences in quality of life (mean difference, 7.41; 95% confidence interval [CI], 7.90 to 12.92; $p=.008$), episodes of incontinence (mean difference, -1.33; 95% CI, -2.50 to -0.17; $p=.02$), and the number of patients who improved or were cured (risk ratio, 1.46; 95% CI, 1.14 to 1.87; $p=.003$) compared to the control group. The biofeedback-assisted PFMT group did not have significant differences in any of these outcomes compared to the control group. Limitations of the study include high heterogeneity for some analyses and differences in the qualitative and quantitative assessments utilized in the included RCTs which limits the direct comparability among the studies.

A 2017 Cochrane review evaluated the effect of PFS on self-reported incontinence.⁶ The review found no difference between PFS and PFMT in the likelihood of cure of stress incontinence at 6 months based on the results of 4 RCTs (N=143; relative risk [RR], 0.51; 95% CI, 0.15 to 1.63). There was also no difference between groups in adverse event rates based on an imprecise estimate (RR, 5.00; 95% CI, 0.25 to 99). Quality of life was not reported. The same review included studies comparing PFS + PFMT versus PFMT alone, finding no difference between groups in incontinence rates based on 3 trials ($n=99$; RR, 0.76; 95% CI, 0.38 to 1.52). The review found a small benefit of PFS + PFMT on incontinence-related quality of life when compared with PFMT alone (standard mean difference, -0.77; 95% CI, -1.11 to -0.42). The review deemed the evidence for PFS alone or in combination with PFMT versus PFMT alone inconclusive for incontinence and quality of life outcomes.

An Agency for Healthcare Research and Quality comparative effectiveness review prepared by Shamliyan et al (2012) identified 9 RCTs evaluating electrical intravaginal stimulation in women with urgency, stress, or mixed incontinence.⁷ Eight of the 9 studies were published in 2000 or earlier; nearly all used a sham treatment as the control. A pooled analysis of continence rates in 8 RCTs comparing electrical PFS with no active treatment yielded an RR of 2.86 (95% CI, 1.57 to 5.23). A pooled analysis of the reduction in incontinence symptoms yielded an RR of 2.01 (95% CI, 1.28 to 3.15). Reviewers concluded that a high level of evidence suggested electrical PFS is associated with increased continence rates, and that such stimulation improved urinary incontinence.

Moroni et al (2016) published a systematic review of conservative treatment for stress urinary incontinence.⁸ Five trials (N=221 women) were identified comparing intravaginal electrical PFS with control. There were insufficient data on cure rates (e.g., continence rates). A pooled analysis of 4 studies reporting urine quantity with a pad weight test found a significantly greater reduction in pad weight in the treatment versus control groups (mean difference, -9.15; 95% CI, -17.22 to -1.08). A

pooled analysis of 2 studies found significantly greater improvement in the incontinence-specific quality of life in the electrical PFS group than in the control group (mean difference, -1.44; 95% CI, -1.94 to -0.95). Three studies were included in a pooled analysis of a number of incontinence episodes; the findings were not reported. Reviewers stated that, among all conservative treatments assessed, evidence was strongest in support of PFS, with or without biofeedback, for treatment of stress urinary incontinence.

Men With Postprostatectomy Urinary Incontinence

Systematic Reviews

Sciarra et al (2021) conducted meta-analyses comparing the effect of PFS with PFMT and biofeedback on urinary incontinence in men following radical prostatectomy.⁹ The review included 5 RCTs of PFS, the most recent of which was published in 2018. PFS devices, frequency, and duration varied among the trials. At 3 months, the effect size for continence recovery (based on pad-free event rate) was 0.57 (95% CI, 0.46 to 0.69) for PFS, 0.40 (95% CI, 0.30 to 0.49) for PFMT, and 0.54 (95% CI, 0.32 to 0.75) for biofeedback ($p=.01$ for both PFS and biofeedback vs. PFMT). At 6 and 12 months, PFS effect sizes were 0.78 (95% CI, 0.59 to 0.98) and 0.82 (95% CI, 0.65 to 0.99), respectively, and there was no longer a statistically significant difference between any treatment group and rate of continence recovery.

A Cochrane review by Berghmans et al (2013) identified 6 RCTs on electrical PFS with nonimplanted electrodes for postprostatectomy urinary incontinence in men.¹⁰ The trials varied by intervention used, study protocols, study populations, and outcome measures. In a pooled analysis of 4 RCTs comparing the combination of electrical stimulation and pelvic floor muscle exercises with pelvic floor muscle exercises alone, there was no statistically significant difference between groups in the proportion of men with urinary incontinence at 3 months (RR, 0.93; 95% CI, 0.82 to 1.06). Findings from studies evaluating electrical PFS alone were not pooled. A 2023 Cochrane review on conservative interventions for managing urinary incontinence after prostate surgery found no studies on electrical or magnetic stimulation compared with no treatment, sham, or verbal/written instructions that reported on key outcomes.¹¹

Zhu et al (2012) conducted a meta-analysis and reported similar findings for electrical PFS to treat postprostatectomy urinary incontinence.¹² Reviewers identified 4 RCTs (N=210 men) that provided sufficient data on clinical outcomes. A pooled analysis of data from 3 trials did not find a statistically significant benefit of electrical PFS on continence levels compared with controls within 3 months of prostatectomy (RR, 1.21; 95% CI, 0.96 to 1.54). Similarly, a pooled analysis of data from all 4 trials did not show a statistically significant benefit of electrical PFS on continence levels 6 to 12 months after prostatectomy (RR, 1.03; 95% CI, 0.88 to 1.20).

Randomized Controlled Trials

Representative trials of men with postprostatectomy urinary incontinence include the RCT by Goode et al (2011) comparing behavioral therapy alone with behavioral therapy plus biofeedback and electrical PFS.¹³ The trial included 208 men with urinary incontinence persisting at least 1 year after radical prostatectomy. Men with preprostatectomy incontinence were excluded. Participants were randomized to 1 of 3 groups: 8 weeks of behavioral therapy (PFMT plus bladder control exercises; $n=70$), behavioral therapy plus biofeedback and electrical stimulation ($n=70$), and a delayed-treatment control group ($n=68$). The biofeedback plus electrical stimulation intervention (called "behavior-plus") consisted of in-office electrical stimulation with biofeedback using an anal probe and daily home electrical PFS. After 8 weeks, patients in the 2 active treatment groups were given instructions for a maintenance program of pelvic floor exercises and fluid control; they were then given follow-up at 6 and 12 months. The primary efficacy outcome was a reduction in the number of incontinent episodes at 8 weeks, as measured by a 7-day bladder diary. A total of 176 (85%) of 208 randomized men completed the 8 weeks of treatment. In an intention-to-treat analysis of the primary outcome, the mean reduction in incontinent episodes was 55% (28 to 13 episodes per week) in the behavioral therapy group, 51% (from 26 to 12 episodes per week) in the behavior-plus group, and

24% (from 25 to 20 episodes per week) in the control group. The overall difference between groups was statistically significant ($p=.001$), but the behavior-plus intervention did not result in a significantly better outcome than behavioral therapy alone. Findings were similar for other outcomes. For example, at the end of 8 weeks, there was a significantly higher rate of complete continence in the active treatment groups (11/70 [16%] in the behavior group vs. 12/70 [17%] in the behavior-plus group) than in the control group (4/68 [6%]); however, the group receiving biofeedback and electrical PFS did not have a significantly higher continence rate than the group receiving behavioral therapy alone. The trial did not isolate the effect of electrical PFS, and the combined behavior-plus intervention did not result in better outcomes than behavioral therapy alone.

Yamanishi et al (2010) published findings of an RCT comparing electrical stimulation with a sham control group.¹⁴ This trial, conducted in Japan, was double-blinded; in it, 56 men with severe postprostatectomy urinary incontinence were randomized to active ($n=26$) or sham ($n=30$) electrical PFS. All the men performed PFMT. Active or sham electrical PFS was performed until incontinence was resolved or until the end of the study at 12 months. Forty-seven patients (22 in the active stimulation group, 25 in the sham group) completed the trial. The continence rate (defined as loss of ≤ 8 grams of urine during a 24-hour pad test) was the primary efficacy outcome. There was a statistically higher rate of continence at 1, 3, and 6 months in the active stimulation group than in the sham group but the between-group difference was not statistically significant at 12 months. The numbers of men reported as continent in the active electrical PFS group were 8 (36%), 14 (63%), 18 (81%), and 19 (86%) at 1, 3, 6, and 12 months, respectively. Corresponding rates in the sham group were 1 (4%), 4 (16%), 11 (44%), and 17 (86%), respectively. Differences in the amount (number of grams) of daily leakage as measured by 24-hour pad tests differed significantly between groups at 1 month; however, the difference disappeared at the 12-month follow-up. For example, after 1 month, the mean amount of leakage was 210 grams in the active treatment group and 423 grams in the sham group ($p>.05$). Change in the amount of daily leakage from baseline differed significantly between groups at 1 month (-528 grams in the active treatment group vs. -257 grams in the sham group, $p<.01$) but not at the other follow-up time points.

Section Summary: Electrical Pelvic Floor Stimulation for Urinary Incontinence

A majority of RCTs on electrical PFS for treatment of women with urinary incontinence have been published before 2001. Meta-analyses of RCTs have had inconsistent findings on the impact of electrical intravaginal stimulation on urinary incontinence in women compared with sham treatment. Individual RCTs have evaluated electrical PFS as a treatment of postprostatectomy urinary incontinence in men. These studies reported improvements in some outcomes with electrical PFS but also have limitations, such as failure to isolate the effect of electrical PFS, and/or failure to find a sham comparator or an accepted treatment comparator. Pooled analyses from 3 systematic reviews found inconsistent evidence on the effect of PFS on continence at 3 months (1 found significant benefit and 2 did not), and found no clear benefit of PFS at 6- and 12-month follow-up.

Electrical Pelvic Floor Stimulation for Fecal Incontinence

Clinical Context and Therapy Purpose

The purpose of PFS in individuals who have fecal incontinence is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with fecal incontinence. Fecal incontinence can have a substantial impact on quality of life. Estimates from the National Center for Health Statistics have suggested that among noninstitutionalized persons, 65 years of age or older, 17% have reported issues with fecal incontinence. Risk factors for fecal incontinence are similar in men and women: older age, diarrhea, fecal urgency, urinary incontinence, and diabetes.

Interventions

The therapy being considered is electrical PFS for fecal incontinence.

Comparators

The following therapies are currently being used to make decisions about fecal incontinence: nonsurgical treatment options and behavioral therapies. Nonsurgical treatment options for incontinence may include pharmacologic therapy, bowel training exercises, and magnetic stimulation. Behavioral therapies include pelvic floor muscle training and diet.

Outcomes

The general outcomes of interest include a reduction in symptoms (e.g., number of incontinence episodes) and improvements in quality of life and cure rates. Electrical PFS therapy generally continues for 6 to 8 weeks.^{15,16}

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- a. To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- b. In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- c. To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- d. Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Systematic Reviews

A systematic review by Vonthein et al (2013) searched for studies on the impact of biofeedback and/or electrical PFS for treating fecal incontinence in adults.¹⁷ They identified 13 RCTs that used 1 or both of these treatments and reported health outcomes (e.g., remission or response rates using validated scales). A pooled analysis of trial results did not find statistically significant differences in rates of remission when comparing electrical PFS with a control intervention (RR, 0.47; 95% CI, 0.13 to 1.72). A pooled analysis of studies comparing electrical PFS plus biofeedback with electrical PFS alone found a significantly higher rate of remission with the combination intervention (RR, 22.97; 95% CI, 1.81 to 291.69). The latter analysis focused on the efficacy of biofeedback and not electrical PFS.

Additionally, the confidence interval was very wide, indicating an imprecise estimate of the treatment effect. The Vonthein et al (2013) review included only 2 RCTs on electrical PFS^{18,19} that were published after a Cochrane review (below). These 2 trials included the combination of amplitude-modulated medium-frequency stimulation and biofeedback. Electrical PFS was not evaluated in the absence of biofeedback.

A Cochrane review by Hosker et al (2007) identified 4 RCTs evaluating electrical stimulation as a treatment of fecal incontinence in adults.²⁰ One trial was sham-controlled, another compared electrical PFS with levatorplasty, and 2 used electrical PFS as an adjunct treatment. Reviewers did not pool study findings; they concluded that there is insufficient evidence to draw conclusions on the efficacy of electrical PFS for treating fecal incontinence.

Randomized Controlled Trials

An RCT by Cohen-Zubary et al (2015) allocated 42 women with fecal incontinence to 6 weeks of electrical stimulation (n=22) or biofeedback training (n=20).¹⁵ Biofeedback sessions were conducted in-clinic and electrical PFS sessions at home following an initial training in-clinic. Thirty-six (86%) women completed the trial and were included in the analysis; the analysis was not intention-to-treat. The trial's primary endpoints were improvements in frequency of fecal, urine, and gas incontinence, assessed using visual analog scale scores. There were no statistically significant differences between

groups for the primary outcomes. The mean visual analog scale score (standard deviation [SD]) for solid stool incontinence at baseline in the stimulation group was 2.9 (2.8), which decreased to 0.9 (0.9) at follow-up. In the biofeedback group, the baseline visual analog scale score was 1.1 (2.1) and 0.3 (0.5) at follow-up. The between-group difference for this outcome was not statistically significant. For within-group changes, the electrical stimulation group improved significantly on solid stool incontinence—but not on liquid stool or gas incontinence and the biofeedback group did not improve significantly on any of the fecal incontinence outcomes.

Norton et al (2006) in the U.K. published a sham-controlled randomized trial that included 90 adults with fecal incontinence.¹⁶ Patients used a home electric PFS device for 8 weeks. Patients allocated to active treatment had the stimulation set at 35 Hz, with a 0.5-second ramped pulse. The sham stimulator looked identical but stimulation was set at 1 Hz below the level tested for therapeutic effect. Patients were blinded to the treatment group; although nurses who trained patients on device use were not. The primary outcome was patient self-report of efficacy, using a rating scale ranging from -5 to +5 to indicate symptom change. Seventy (78%) of the 90 patients completed the trial. In an intention-to-treat analysis (assigning patients who dropped out a value of 0), there was no statistically significant difference between groups in patient ratings of symptom change. On a scale of -5 to +5, there was a median rating of 0 in each group ($p=.92$). In a completer analysis, the median change in symptoms was 2 in the active treatment group and 1 in the sham group ($p=.74$). Groups did not differ significantly on other secondary outcomes such as the frequency of urge or passive incontinence after treatment.

Section Summary: Electrical Pelvic Floor Stimulation for Fecal Incontinence

Several RCTs have evaluated electrical stimulation for treating fecal incontinence. Only 1 was sham-controlled, and it did not find that active stimulation produced better results than sham stimulation. Systematic reviews of RCTs have not found that electrical stimulation is superior to control interventions for treating fecal incontinence.

Magnetic Pelvic Floor Stimulation for Urinary Incontinence

Clinical Context and Therapy Purpose

The purpose of magnetic PFS in individuals who have urinary incontinence is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with urinary incontinence. Types of urinary incontinence include stress incontinence, urgency incontinence, and mixed (both stress and urgency). Urinary incontinence in women is common, with some estimates citing a 50% incidence. Factors that increase a woman's risk include older age, obesity, parity, vaginal delivery, and family history. Urinary incontinence is less common in men, with estimates ranging from 11% to 34% in men greater than 65 years. Factors that increase a man's risk include older age, prostate disease, urinary tract infection history, impaired activities of daily living, neurologic disease, constipation, diabetes, and sleep apnea.

Interventions

The therapy being considered is magnetic PFS for urinary incontinence. The mechanism of action of a magnetic PFS procedure is similar to the electrical procedure, though using magnetic pulses to activate the pelvic floor musculature. The magnetic pulses are delivered without a probe, with patients sitting fully clothed in a specialized chair with an embedded magnet.

Comparators

The following therapies are currently being used to make decisions about urinary incontinence: electrical PFS and behavioral therapies (e.g., monitoring fluid intake, pelvic floor muscle training, diet), and medications.

Outcomes

The general outcomes of interest include a reduction in symptoms (e.g., number of incontinence episodes) and improvements in quality of life and cure rates. Treatment is for approximately 8 weeks, and follow-up is generally up to 6 months.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

Women with Urinary Incontinence

Systematic Reviews

A systematic review of RCTs on magnetic stimulation for the treatment of urinary incontinence was published by Lim et al (2015).²¹ Reviewers identified 8 blinded, sham-controlled trials (N=484). Treatment protocols (e.g., frequency, duration of magnetic PFS) varied among trials. The primary outcome was cure rate; only 1 trial reported this outcome, so data were not pooled. A meta-analysis of 3 studies reporting improvements in the continence rates found significantly greater improvement in the treatment group than in the sham group (RR, 2.29; 95% CI, 1.60 to 3.29). Due to the variability across trials in types of incontinence treated and/or outcome reporting, data were not pooled for other outcomes. Reviewers noted that the evidence was limited by low-quality trials with short-term follow-up.

Randomized Controlled Trials

Yamanishi et al (2014) published an industry-sponsored evaluation of magnetic PFS provided to women with urinary urgency using an armchair-type stimulator.²² The device was produced by a Japanese company and does not have FDA approval. Patients received active (n=101) or sham (n=50) stimulation, 2 times a week for 6 weeks. The level of stimulation was tailored to each patient's maximum tolerable intensity; sham stimulation was set at a lower level than active treatment. Because noises differed between the 2 procedures, patients were isolated from the sounds to maintain blinding. Study personnel were not blinded. A total of 143 (95%) of 151 patients were included in the efficacy analysis. The primary endpoint was a change in the number of urinary incontinence episodes per week, as reported in a patient diary. The decrease in the weekly number (SD) of incontinence episodes was 13 (11) in the active treatment group compared with 9 (13) in the sham group ($p=.038$). Patients in the active stimulation group had significantly better results on some secondary outcomes (e.g., number of urgency episodes per 24 hours) but not others (e.g., number of voids per 24 hours).

A sham-controlled randomized trial evaluating magnetic PFS using the NeoControl chair did not find evidence that PFS improved outcomes. In this trial by Gillig et al (2009) in New Zealand, sham treatment involved inserting a thin aluminum plate in the chair to prevent penetration of the magnetic field.²³ The trial included 70 women, 35 in each group, with stress or mixed urinary incontinence. Both groups received 3 treatment sessions per week for 6 weeks. There was no significant difference between the active and sham treatment groups for the primary outcome measure, change from baseline in the 20-minute pad test result to 8 weeks after the start of treatment (2 weeks after finishing treatment). At 8 weeks, the mean change in the 20-minute pad test was 20.1 mL in the treatment group and 7.5 mL in the control group. The groups also did not differ significantly in the 20-minute pad weight or quality of life measure at the 6-month follow-up.

Data from 29 (83%) women in the active treatment group and 26 (74%) women in the sham group were available at 6 months; all participants appear to be included in the 8-week outcomes analysis. Lim et al (2017) randomized 120 women with stress urinary incontinence to treatment with magnetic PFS (QRS® -1010 PelviCenter) or sham treatment.²⁴ Patients received 2 sessions per week for 8 weeks (16 sessions). Patients who were unsatisfied after 2 months were allowed 16 additional active sessions in an open-label phase. All participating study centers were located in Malaysia. The primary endpoint of response was defined as a 5-point reduction on the International Consultation on Incontinence Questionnaire for Urinary Incontinence-Short Form (ICIQ-UI SF). A total of 45 (75.0%) patients responded at 2 months in the active treatment group compared with 13 (21.7%) patients in the sham group (RR, 3.46; 95% CI, 2.09 to 5.72; $p < .001$). At long-term follow-up (14 months), the patients who received 16 or more active sessions had improved response rates than those who received none (response rates of 68.3% to 75.0% vs. 21.1%). The study is limited by the small sample size and the limited demographic heterogeneity.

Men With Postprostatectomy Urinary Incontinence Systematic Review

A 2023 Cochrane review on conservative interventions for managing urinary incontinence after prostate surgery found no studies on electrical or magnetic stimulation compared with no treatment, sham, or verbal/written instructions that reported on key outcomes.¹¹

Randomized Controlled Trial

One RCT was identified on magnetic PFS for treating postprostatectomy urinary incontinence. Yokoyama et al (2004) reported findings from a 3-arm randomized trial.²⁵ Thirty-six men (12 in each group) were randomized to extracorporeal magnetic PFS (NeoControl chair), functional electrical PFS, or pelvic floor exercises. The primary outcome was pad weight testing for up to 6 months after the 1-month treatment period. At 1 month after catheter removal, pad weight was significantly lower in the electrical PFS group than in the control group; at 2 months after catheter removal, pad weight was significantly lower in the magnetic PFS group compared with the control group; and, beginning at 3 months after catheter removal, there were no significant differences across arms in pad weight. Additionally, there were no significant differences between groups in quality of life measurements at any follow-up point. The trial lacked a sham magnetic stimulation group and therefore a placebo effect cannot be ruled out as an explanation for the short-term reduction in pad weight in the magnetic PFS treatment group.

Section Summary: Magnetic Pelvic Floor Stimulation for Urinary Incontinence

A systematic review of RCTs evaluating the use of magnetic PFS for urinary incontinence in women concluded that the evidence was insufficient due to the small number of trials with short-term follow-up, methodologic limitations, and heterogeneity in terms of patient populations, interventions, and outcome reporting.

One RCT evaluated magnetic PFS for the treatment of men with postprostatectomy urinary incontinence. There was a greater improvement in pad weight at 2 months in the magnetic PFS group than in the pelvic floor muscle exercises group but there were no significant differences between groups beginning at 3 months. Other outcomes also did not favor the magnetic PFS group.

A 2023 systematic review was unable to identify studies on magnetic PFS evaluating outcomes of interest.

Magnetic Pelvic Floor Stimulation for Fecal Incontinence Clinical Context and Therapy Purpose

The purpose of PFS in individuals who have fecal incontinence is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with fecal incontinence. Risk factors for fecal incontinence are similar in men and women: older age, diarrhea, fecal urgency, urinary incontinence, and diabetes. For women, current and past use of hormone therapy is an added risk factor. Fecal incontinence can have a substantial impact on quality of life. Estimates from the National Center for Health Statistics have suggested that among noninstitutionalized persons, 65 years of age or older, 17% have reported issues with fecal incontinence.

Interventions

The therapy being considered is magnetic PFS for fecal incontinence. The mechanism of action of a magnetic PFS procedure is similar to the electrical procedure, though using magnetic pulses to activate the pelvic floor musculature. The magnetic pulses are delivered without a probe, with patients sitting fully clothed in a specialized chair with an embedded magnet.

Comparators

The following therapies are currently being used to make decisions about fecal incontinence: nonsurgical treatment options and behavioral therapies. Nonsurgical treatment options for incontinence may include pharmacologic therapy, bowel training exercises, and electrical stimulation.

Behavioral therapies include pelvic floor muscle training and diet.

Outcomes

The general outcomes of interest include a reduction in symptoms (e.g., number of incontinence episodes) and improvements in quality of life and cure rates. Treatment is for approximately 8 weeks, and follow-up is generally up to 6 months.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Review of Evidence

No studies were identified that evaluated magnetic PFS as a treatment of fecal incontinence.

Section Summary: Magnetic PFS for Fecal Incontinence

Current evidence is insufficient to draw conclusions about the efficacy of magnetic PFS to treat fecal incontinence.

Summary of Evidence

For individuals who have urinary incontinence who receive electrical pelvic floor stimulation (PFS), the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. Findings from systematic reviews have not found that electrical PFS used to treat urinary incontinence in women consistently improves the net health outcome compared with placebo or other conservative treatments. Moreover, meta-analyses of RCTs have not found a significant benefit of electrical PFS in men with postprostatectomy incontinence compared with a control intervention. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome. For individuals who have fecal incontinence who receive electrical PFS, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, change in disease status, quality of life,

and treatment-related morbidity. Among the RCTs that have evaluated electrical PFS as a treatment for fecal incontinence, only 1 trial was sham-controlled, and it did not find that electrical stimulation improved the net health outcome. Systematic reviews of RCTs have not found that electrical stimulation is superior to control interventions for treating fecal incontinence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have urinary incontinence who receive magnetic PFS, the evidence includes RCTs and 2 systematic reviews. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. A systematic review of RCTs on magnetic PFS for urinary incontinence in women concluded that the evidence was insufficient due to the following factors: a low number of trials with short-term follow-up, methodologic limitations, as well as heterogeneity in patient populations, interventions, and outcomes reported. One RCT evaluating magnetic stimulation for treating men with postprostatectomy urinary incontinence reported short-term results favoring magnetic PFS; however, the trial was small and lacked a sham comparator. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome. For individuals who have fecal incontinence who receive magnetic PFS, no relevant evidence was identified. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

The purpose of the following information is to provide reference material. Inclusion does not imply endorsement or alignment with the evidence review conclusions.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Gastroenterology

In 2021, the American College of Gastroenterology issued guidelines on the management of benign anorectal disorders.²⁶ In the section on fecal incontinence, pelvic floor stimulation (PFS) is not mentioned as a treatment option.

American Society of Colon and Rectal Surgeons

In 2023, the American Society of Colon and Rectal Surgeons updated an evidence-based guideline using GRADE methodology on treatment of fecal incontinence.²⁷ Dietary interventions and medical management are considered first-line treatments; PFS was not included in the recommendations.

American Urological Association et al

In 2024, the American Urological Association (AUA) and Society of Urodynamics, Female Pelvic Medicine and Urogenital Reconstruction (SUFU) updated guidelines on the diagnosis and management of overactive bladder.²⁸ Electromagnetic therapy is included as an example of non-invasive therapy. The recommendation states, "Clinicians may offer select non-invasive therapies to all patients with OAB." However, the guidelines also state, "While safety profiles are excellent across modalities, with few adverse effects and a high risk-benefit ratio, all non-invasive therapies do not have equivalent efficacy, and the evidence base is highly variable. Most non-invasive therapies require long-term patient compliance to maintain a durable effect, and patients should be counselled as such before embarking on a course of a potentially lifelong therapy." There is no additional information specific to PFS in the guidelines.

Joint guidelines issued in 2019 by the AUA and the Society of Urodynamics, Female Pelvic Medicine and Urogenital Reconstruction (SUFU) on management of post-prostatectomy urinary incontinence do not specifically address electrical or magnetic PFS as treatment options. Pelvic floor muscle training/exercise is recommended as first-line treatment for post-prostatectomy incontinence.²⁹

National Institute for Health and Care Excellence

In 2019, the NICE issued guidance on the management of urinary incontinence in women.³⁰ The NICE stated that electrical stimulation, alone or as an adjunct to pelvic floor muscle training, should not be routinely used to treat women with overactive bladder. The NICE guidance further stated: "electrical stimulation and/or biofeedback should be considered in women who cannot actively contract pelvic floor muscles in order to aid motivation and adherence to therapy." Magnetic PFS is not mentioned. In 2007, the NICE issued guidance on the management of fecal incontinence in adults.³¹ This guidance was last reviewed by NICE in 2018. The document stated that the evidence on electrical stimulation for treatment of fecal incontinence was inconclusive. The NICE recommended that patients who continue to have episodes of fecal incontinence after initial treatment be considered for specialized management, which may include electrical PFS. Magnetic PFS is not mentioned.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

The national coverage determination for Non-Implantable Pelvic Floor Electrical Stimulator (230.8) stated: "Pelvic floor electrical stimulation with a non-implantable stimulator is covered for the treatment of stress and/or urge urinary incontinence in cognitively intact patients who have failed a documented trial of pelvic muscle exercise (PME) training."³² The effective date was June 19, 2006.

The document did not mention fecal incontinence.

Ongoing and Unpublished Clinical Trials

Some currently ongoing and unpublished trials that might influence this review are listed in Table 1.

Table 1. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT05952258	Magnetic Stimulation as a Treatment for Stress urinary Incontinence	158	Jul 2026
<i>Unpublished</i>			
NCT04644614	Effectiveness of Magnetic Stimulation in Patients With Urinary Incontinence After Radical Prostatectomy: a Prospective Randomized Sham Controlled Clinical Study	40	Apr 2023

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

References

1. Gorina Y, Schappert S, Bercovitz A, et al. Prevalence of incontinence among older americans. Vital Health Stat 3. Jun 2014; (36): 1-33. PMID 24964267
2. Markland AD, Goode PS, Redden DT, et al. Prevalence of urinary incontinence in men: results from the national health and nutrition examination survey. J Urol. Sep 2010; 184(3): 1022-7. PMID 20643440
3. Abdelbary AM, El-Dessoukey AA, Massoud AM, et al. Combined Vaginal Pelvic Floor Electrical Stimulation (PFS) and Local Vaginal Estrogen for Treatment of Overactive Bladder (OAB) in

- Perimenopausal Females. Randomized Controlled Trial (RCT). *Urology*. Sep 2015; 86(3): 482-6. PMID 26135813
4. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Pelvic floor electrical stimulation in the treatment of urinary incontinence in adults. *TEC Assessments*. 2000;Volume 15:Tab 2.
 5. Leonardo K, Seno DH, Mirza H, et al. Biofeedback-assisted pelvic floor muscle training and pelvic electrical stimulation in women with overactive bladder: A systematic review and meta-analysis of randomized controlled trials. *Neurourol Urodyn*. Aug 2022; 41(6): 1258-1269. PMID 35686543
 6. Stewart F, Berghmans B, Bø K, et al. Electrical stimulation with non-implanted devices for stress urinary incontinence in women. *Cochrane Database Syst Rev*. Dec 22 2017; 12(12): CD012390. PMID 29271482
 7. Shamliyan T, Wyman J, Kane R. *Nonsurgical Treatments for Urinary Incontinence in Adult Women: Diagnosis and Comparative Effectiveness (Comparative Effectiveness Review 36)*. Rockville, MD: Agency for Healthcare Research and Quality; 2012.
 8. Moroni RM, Magnani PS, Haddad JM, et al. Conservative Treatment of Stress Urinary Incontinence: A Systematic Review with Meta-analysis of Randomized Controlled Trials. *Rev Bras Ginecol Obstet*. Feb 2016; 38(2): 97-111. PMID 26883864
 9. Sciarra A, Viscuso P, Arditi A, et al. A biofeedback-guided programme or pelvic floor muscle electric stimulation can improve early recovery of urinary continence after radical prostatectomy: A meta-analysis and systematic review. *Int J Clin Pract*. Oct 2021; 75(10): e14208. PMID 33811418
 10. Berghmans B, Hendriks E, Bernards A, et al. Electrical stimulation with non-implanted electrodes for urinary incontinence in men. *Cochrane Database Syst Rev*. Jun 06 2013; (6): CD001202. PMID 23740763
 11. Johnson EE, Mamoulakis C, Stoniute A, et al. Conservative interventions for managing urinary incontinence after prostate surgery. *Cochrane Database Syst Rev*. Apr 18 2023; 4(4): CD014799. PMID 37070660
 12. Zhu YP, Yao XD, Zhang SL, et al. Pelvic floor electrical stimulation for postprostatectomy urinary incontinence: a meta-analysis. *Urology*. Mar 2012; 79(3): 552-5. PMID 22386394
 13. Goode PS, Burgio KL, Johnson TM, et al. Behavioral therapy with or without biofeedback and pelvic floor electrical stimulation for persistent postprostatectomy incontinence: a randomized controlled trial. *JAMA*. Jan 12 2011; 305(2): 151-9. PMID 21224456
 14. Yamanishi T, Mizuno T, Watanabe M, et al. Randomized, placebo controlled study of electrical stimulation with pelvic floor muscle training for severe urinary incontinence after radical prostatectomy. *J Urol*. Nov 2010; 184(5): 2007-12. PMID 20850831
 15. Cohen-Zubary N, Gingold-Belfer R, Lambort I, et al. Home electrical stimulation for women with fecal incontinence: a preliminary randomized controlled trial. *Int J Colorectal Dis*. Apr 2015; 30(4): 521-8. PMID 25619464
 16. Norton C, Gibbs A, Kamm MA. Randomized, controlled trial of anal electrical stimulation for fecal incontinence. *Dis Colon Rectum*. Feb 2006; 49(2): 190-6. PMID 16362803
 17. Vonthein R, Heimerl T, Schwandner T, et al. Electrical stimulation and biofeedback for the treatment of fecal incontinence: a systematic review. *Int J Colorectal Dis*. Nov 2013; 28(11): 1567-77. PMID 23900652
 18. Schwandner T, König IR, Heimerl T, et al. Triple target treatment (3T) is more effective than biofeedback alone for anal incontinence: the 3T-AI study. *Dis Colon Rectum*. Jul 2010; 53(7): 1007-16. PMID 20551752
 19. Schwandner T, Hemmelmann C, Heimerl T, et al. Triple-target treatment versus low-frequency electrostimulation for anal incontinence: a randomized, controlled trial. *Dtsch Arztebl Int*. Sep 2011; 108(39): 653-60. PMID 22013492
 20. Hosker G, Cody JD, Norton CC. Electrical stimulation for faecal incontinence in adults. *Cochrane Database Syst Rev*. Jul 18 2007; 2007(3): CD001310. PMID 17636665
 21. Lim R, Lee SW, Tan PY, et al. Efficacy of electromagnetic therapy for urinary incontinence: A systematic review. *Neurourol Urodyn*. Nov 2015; 34(8): 713-22. PMID 25251335

22. Yamanishi T, Homma Y, Nishizawa O, et al. Multicenter, randomized, sham-controlled study on the efficacy of magnetic stimulation for women with urgency urinary incontinence. *Int J Urol. Apr 2014; 21(4): 395-400. PMID 24118165*
23. Gilling PJ, Wilson LC, Westenberg AM, et al. A double-blind randomized controlled trial of electromagnetic stimulation of the pelvic floor vs sham therapy in the treatment of women with stress urinary incontinence. *BJU Int. May 2009; 103(10): 1386-90. PMID 19154474*
24. Lim R, Liong ML, Leong WS, et al. Pulsed Magnetic Stimulation for Stress Urinary Incontinence: 1-Year Followup Results. *J Urol. May 2017; 197(5): 1302-1308. PMID 27871927*
25. Yokoyama T, Nishiguchi J, Watanabe T, et al. Comparative study of effects of extracorporeal magnetic innervation versus electrical stimulation for urinary incontinence after radical prostatectomy. *Urology. Feb 2004; 63(2): 264-7. PMID 14972468*
26. Wald A, Bharucha AE, Limketkai B, et al. ACG Clinical Guidelines: Management of Benign Anorectal Disorders. *Am J Gastroenterol. Oct 01 2021; 116(10): 1987-2008. PMID 34618700*
27. Bordeianou LG, Thorsen AJ, Keller DS, et al. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Fecal Incontinence. *Dis Colon Rectum. May 01 2023; 66(5): 647-661. PMID 36799739*
28. Cameron AP, Chung DE, Dielubanza EJ, et al. The AUA/SUFU Guideline on the Diagnosis and Treatment of Idiopathic Overactive Bladder. *J Urol. Jul 2024; 212(1): 11-20. PMID 38651651*
29. Sandhu JS, Breyer B, Comiter C, et al. Incontinence after Prostate Treatment: AUA/SUFU Guideline. *J Urol. Aug 2019; 202(2): 369-378. PMID 31059663*
30. National Institute for Health and Care Excellence (NICE) Guideline. Urinary Incontinence and Pelvic Organ Prolapse in Women: Management. NICE Guideline. 2019. <https://www.nice.org.uk/guidance/ng123>. Accessed June 11, 2024.
31. National Institute for Health and Care Excellence (NICE). Faecal incontinence in adults: management [CG49]. 2007; <https://www.nice.org.uk/guidance/cg49>. Accessed June 10, 2024.
32. Centers for Medicare & Medicaid Services (CMS). CMS Manual System: Pub 100-03 Medicare National Coverage Determinations; Transmittal 48. 2006; <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=231>. Accessed June 11, 2024.

Documentation for Clinical Review

- No records required

Coding

This Policy relates only to the services or supplies described herein. Benefits may vary according to product design; therefore, contract language should be reviewed before applying the terms of the Policy.

The following codes are included below for informational purposes. Inclusion or exclusion of a code(s) does not constitute or imply member coverage or provider reimbursement policy. Policy Statements are intended to provide member coverage information and may include the use of some codes for clarity. The Policy Guidelines section may also provide additional information for how to interpret the Policy Statements and to provide coding guidance in some cases.

Type	Code	Description
CPT®	97014	Application of a modality to 1 or more areas; electrical stimulation (unattended)
	97032	Application of a modality to 1 or more areas; electrical stimulation (manual), each 15 minutes
HCPCS	E0715	Intravaginal device intended to strengthen pelvic floor muscles during kegel exercises (Code effective 10/01/2024)

Type	Code	Description
	E0716	Supplies and accessories for intravaginal device intended to strengthen pelvic floor muscles during kegel exercises (<i>Code effective 10/01/2024</i>)
	E0740	Non-implanted pelvic floor electrical stimulator, complete system
	G0283	Electrical stimulation (unattended), to one or more areas for indication(s) other than wound care, as part of a therapy plan of care

Policy History

This section provides a chronological history of the activities, updates and changes that have occurred with this Medical Policy.

Effective Date	Action
02/27/2015	Policy title change from Urinary Incontinence Outpatient Treatment BCBSA Medical Policy adoption Policy revision with position change
04/30/2015	Coding update
07/31/2015	Policy revision without position change
12/01/2016	Policy revision without position change
10/01/2017	Policy revision without position change
10/01/2018	Policy revision without position change
12/16/2019	Policy revision without position change
04/01/2020	Annual review. No change to policy statement.
11/01/2020	No change to policy statement. Literature review updated.
10/01/2021	Annual Review. No change to policy statement. Literature review updated.
10/01/2022	Annual Review. No change to policy statement. Literature review updated.
10/01/2023	Annual Review. No change to policy statement. Literature review updated.
10/01/2024	Annual Review. No change to policy statement. Policy guidelines and literature review updated.
11/01/2024	Coding update.

Definitions of Decision Determinations

Medically Necessary: Services that are Medically Necessary include only those which have been established as safe and effective, are furnished under generally accepted professional standards to treat illness, injury or medical condition, and which, as determined by Blue Shield, are: (a) consistent with Blue Shield medical policy; (b) consistent with the symptoms or diagnosis; (c) not furnished primarily for the convenience of the patient, the attending Physician or other provider; (d) furnished at the most appropriate level which can be provided safely and effectively to the patient; and (e) not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of the Member's illness, injury, or disease.

Investigational/Experimental: A treatment, procedure, or drug is investigational when it has not been recognized as safe and effective for use in treating the particular condition in accordance with generally accepted professional medical standards. This includes services where approval by the federal or state governmental is required prior to use, but has not yet been granted.

Split Evaluation: Blue Shield of California/Blue Shield of California Life & Health Insurance Company (Blue Shield) policy review can result in a split evaluation, where a treatment, procedure, or drug will be considered to be investigational for certain indications or conditions, but will be deemed safe and

effective for other indications or conditions, and therefore potentially medically necessary in those instances.

Prior Authorization Requirements and Feedback (as applicable to your plan)

Within five days before the actual date of service, the provider must confirm with Blue Shield that the member's health plan coverage is still in effect. Blue Shield reserves the right to revoke an authorization prior to services being rendered based on cancellation of the member's eligibility. Final determination of benefits will be made after review of the claim for limitations or exclusions.

Questions regarding the applicability of this policy should be directed to the Prior Authorization Department at (800) 541-6652, or the Transplant Case Management Department at (800) 637-2066 ext. 3507708 or visit the provider portal at www.blueshieldca.com/provider.

We are interested in receiving feedback relative to developing, adopting, and reviewing criteria for medical policy. Any licensed practitioner who is contracted with Blue Shield of California or Blue Shield of California Promise Health Plan is welcome to provide comments, suggestions, or concerns. Our internal policy committees will receive and take your comments into consideration.

For utilization and medical policy feedback, please send comments to: MedPolicy@blueshieldca.com

Disclaimer: This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. Blue Shield of California may consider published peer-reviewed scientific literature, national guidelines, and local standards of practice in developing its medical policy. Federal and state law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over medical policy and must be considered first in determining covered services. Member contracts may differ in their benefits. Blue Shield reserves the right to review and update policies as appropriate.

Appendix A

POLICY STATEMENT (No changes)	
BEFORE	AFTER
<p>Pelvic Floor Stimulation as a Treatment of Urinary and Fecal Incontinence 1.01.17</p> <p>Policy Statement:</p> <ul style="list-style-type: none"> I. Electrical or magnetic stimulation of the pelvic floor muscles (pelvic floor stimulation) is considered investigational as a treatment for either of the following: <ul style="list-style-type: none"> A. Urinary incontinence B. Fecal incontinence 	<p>Pelvic Floor Stimulation as a Treatment of Urinary and Fecal Incontinence 1.01.17</p> <p>Policy Statement:</p> <ul style="list-style-type: none"> I. Electrical or magnetic stimulation of the pelvic floor muscles (pelvic floor stimulation) is considered investigational as a treatment for either of the following: <ul style="list-style-type: none"> A. Urinary incontinence B. Fecal incontinence